



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

The impact of major disease outbreaks in the third millennium on adolescent and youth sexual and reproductive health and rights in LMICs: a systematic scoping review protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051216
Article Type:	Protocol
Date Submitted by the Author:	13-Mar-2021
Complete List of Authors:	<p>Akbarialiabad, Hossein; Shiraz University of Medical Sciences, Student research committee, Shiraz school of medicine, Shiraz University of Medical Sciences, Shiraz, Iran</p> <p>Shidhaye, Rahul ; Pravara Institute of Medical Sciences, Senior Research Scientist and Associate Professor of Psychiatry, Pravara Institute of Medical Sciences, Loni, India.</p> <p>Shidhaye, Pallavi; ICMR-National AIDS Research Institute, Scientist B, Division of Clinical Sciences, ICMR-National AIDS Research Institute Pune, India</p> <p>Cuijpers, Pim; VU University Amsterdam, Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research institute, and WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam</p> <p>Weaver, Marcia; University of Washington, Departments of Health Metrics Sciences and Global Health University of Washington</p> <p>Kiburi, Sarah; Department of psychiatry, Mbagathi Hospital, Nairobi Metropolitan Service, Nairobi, Kenya</p> <p>Njuguna, Irene; Kenyatta National Hospital, Research & Programs</p> <p>Taghrir, Mohammad Hossein ; Department of community medicine, Shiraz School of medicine, Shiraz University of medical sciences, Shiraz, Iran</p> <p>KUMAR, MANASI; University College London, Department of Psychology, University College London, London, United Kingdom ; Department of Psychiatry, University of Nairobi, Nairobi, Kenya</p>
Keywords:	Sexual and gender disorders < PSYCHIATRY, SEXUAL MEDICINE, REPRODUCTIVE MEDICINE, Child & adolescent psychiatry < PSYCHIATRY, PUBLIC HEALTH

SCHOLARONE™
Manuscripts

The impact of major disease outbreaks in the third millennium on adolescent and youth sexual and reproductive health and rights in LMICs: a systematic scoping review protocol

Hossein Akbarialiabad¹, Rahul Shidhaye², Pallavi Shidhaye³, Pim Cuijpers⁴, Marcia R Weaver⁵, Sarah Kanana⁶, Irene Njuguna⁷, Mohammad Hossein Taghrir⁸, Manasi Kumar^{9,10*}

¹- Student research committee, Shiraz school of medicine, Shiraz University of Medical Sciences, Shiraz, Iran

²- Research Scientist and Associate Professor of Psychiatry, Pravara Institute of Medical Sciences, Loni, India.

³- Scientist B, Division of Clinical Sciences, ICMR-National AIDS Research Institute, Pune, India

⁴-Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research Institute, and WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam, The Netherlands

⁵- Research Professor, Departments of Health Metrics Sciences and Global Health, University of Washington, Seattle Washington, USA

⁶-Department of psychiatry, Mbagathi Hospital, Nairobi Metropolitan Service, Nairobi, Kenya

⁷- Kenyatta National Hospital, Research and Programs, Nairobi, Kenya

⁸- Department of community medicine, Shiraz School of medicine, Shiraz University of medical sciences, Shiraz, Iran

⁹- Department of Psychiatry, University of Nairobi, Nairobi, Kenya

¹⁰- Department of Psychology, University College London, London, United Kingdom

Corresponding author:

Dr. Manasi Kumar

Department of Psychiatry, University of Nairobi, Nairobi, Kenya

Department of Psychology, University College London, London, United Kingdom

Email: Email: m.kumar@ucl.ac.uk / Phone number: +254717379687/ Fax: +254-20-2723719

Word Count : 3596

Keyword: Sexual health, SARS-COV-2, Zika virus, Ebolavirus, H1N1 Virus

Abstract

Introduction

Adolescent sexual and reproductive health (SRHR) continues to present a high burden and remains underinvested. This is more so in lower and middle-income countries (LMICs), where empirical evidence reveals disruption of SRHR maintenance, need for enhancement of programs, resources, and services during pandemics. Despite the importance of the subject, there is no published review yet combining recent disease outbreaks such as (H1N1/09, Zika, Ebola, and SARS-COV-2) to assess their impact on adolescents and youth SRHR in LMICs.

Method

We will adopt a four-step search to reach the maximum possible number of studies. In the first step, we had a limited/preliminary investigation in databases for reaching relevant keywords (appendix1). Secondly, we will search in four databases: Pubmed, Cochrane Library, Embase, and PsycINFO. The search would begin from the inception of the first major outbreak in 2009 (H1N1/09) up to the date of publication of the protocol in early 2021. We will search databases using related keywords, screen title/abstract, and review full texts of the selected titles to arrive at the list of eligible studies. In the third stage, we will go to the included article's reference list to check their eligibility. In the fourth stage, we will check the citation of included papers in phase 2 to complete our study selection. We will include all types of original studies and without any language restriction in our final synthesis. Our review results would be charted for each pandemic separately and include the details related to authors, year, country, region of the study, study design, participants (disaggregated by age and gender), purpose, and report associated SRHR outcomes. The review will adhere to the PRISMA-ScR guideline.

Ethics and dissemination

Ethical assessment is not required for this study. The results of the study will be presented in peer-reviewed publications and conferences on adolescent SRHR.

Strengths and limitations of our study

Strengths

- ✓ we are targeting an underserved area of research adolescent SRHR and how select disease outbreaks impact the adolescent outcomes
- ✓ our systematic scoping review protocol minimizes the possibility of duplications, and we have sought to engage a stringent peer-review process to arrive at meaningful search outcomes around SRHR impacts.
- ✓ our strategy will include four-step of searching and sifting through studies, which maximize our effort in reaching the all possible eligible studies
- ✓ using ICPD as our guiding framework to elicit key outcomes, we hope to arrive at studies that are tracking adolescent SRHR in a rigorous and insightful manner

Limitations

- There could be other diseases outbreaks to consider; however, we took a broad sweep at significant outbreaks to document here

Introduction

The beginning of sexual activity differs based on sex, country, culture, religion, and context (1). Meanwhile, more people are currently reaching puberty earlier and engaging later in sexual activities and marriage. So, before marriage, they are sexually active longer than their parents (2). The risk of neglecting the adolescent sexual and reproductive health (ASRH) could impose a life-lasting impact on them. For females, adolescent pregnancy and maternity may be mentally and physically challenging (3, 4). It may also impede further academic/educational achievements and economic potential (5). In both sexes, especially for girls, there is a high risk of sexually transmitted infections (STIs), including HIV, reproductive coercion, and violence (6). Approximately one-ninth (11%) of the annual birth rates are related to mothers in the age of 15-19. Rates differ by region, but it is estimated that the percentage of adolescents' pregnancy to all births ranges from 50% in Sub-Saharan Africa to 2% in China (3). In 2019, in LMICs, those between 15-19 gave birth to about 21 million. The number of unintended pregnancies in those between 15-19 was about ten million, and about 55% of these unintended pregnancies resulted in abortion (7, 8). Nearly 70% of the total abortion attempt at the ages of 15-19 leads to unsafe abortions, which can have catastrophic health effects (8). In the GBD 2015 report, the maternal mortality rate (MMR) was highest in 10-14 years old girls, as high as 278 (95% UI 229–339) (9). Additionally, pregnancy and delivery-related complications cause more mortality in 15- 19 years old girls (10).

Value of the proposed review: The already missed adolescents and young SRHR need to develop and become more complicated during the disasters and outbreaks (11). Besides, they have less life experience and theoretically do not have evolved decision making, coping, and judgment capabilities to navigate such complex circumstances. These crises offer a belated opportunity for the governments to fill the gaps in sex education and providing access to contraceptives, safer motherhood, safer abortion, and empowerment programs to impede discrimination and protect adolescents living with disabilities and special conditions. If governments do not take such opportunities, future outbreaks can be a more significant public health issue and exacerbate existing mental health issues, gender inequalities, and social injustice. We know that these social determinants of adolescent health are tied together, and this review will help understand the existing evidence and gaps. We have prioritized studying four significant outbreaks since 2000: swine flu virus (H1N1/09), Ebola virus, Zika virus, and SARS-COV-2 and their impacts on the SRHR of young and adolescents in LMICs.

A) Influenza (H1N1/09) :

On first July 2009, the world health organization acknowledged that the flu outbreak has become universal and called it the Pandemic H1N1/09 virus, becoming popular as “swine flu.” Further studies showed that pregnant women are in danger of this virus (12). Pregnant women were considered a high-risk group for H1N1, as there was a higher rate of spontaneous abortion, preterm birth, low birth weight, fetal disease, and maternal death in those who had the virus (13, 14). In a study at the outset of the pandemic in Australia and New Zealand, around 9% of patients admitted to the Intensive Care Units (ICUs) were pregnant (comparing to 1% in the general population). Moreover, the mortality rate was 11% among those pregnant women (15). In flu pandemic, a global pooled analysis revealed that pregnant women have a higher chance of hospitalization than non-pregnant women of childbearing age, with a relative risk of 3.5-25.3 (16). The virus's vertical transmission during pregnancy has been reported

1
2
3 rarely(14). Due to the high mortality and morbidity rate of the virus for pregnant women, vaccination of
4 pregnant women was prioritized to protect them. However, studies showed that mothers do not show
5 compliance to use the vaccine as they had concerns regarding vaccines' adverse effects (17, 18).
6 Fortunately, multiple studies showed that vaccines are safe and have no significant threat to pregnancy,
7 including spontaneous abortion, still birth and congenital malformations(19-21).
8

9
10 **B) Zika Virus :**

11 In March 2015, a group of patients was admitted in Brazil with rash, fatigue, and arthralgia, caused by
12 Zika Virus. In October that year, a few reports showed that the mother-newborn transmission of the virus
13 (22) might lead to fetal loss, dead fetus, preterm delivery, and microcephalic (small head) babies,
14 cumulatively known as congenital Zika syndrome (CZS) (23). Reports showed that the viral load is
15 around 10,000 times more in semen than in the blood (24, 25). The transmission of the virus in human
16 cycles is sexual contact, blood transfusion, and organ transplantation (26-28). Combining the sexual
17 transmission of virus between partners and vertical transmission to the fetus, the WHO recommended
18 safer practice of sexual activities or abstinence during pregnancy. Furthermore, in the case of a pregnancy
19 probability, emergency pills has been recommended (22). To date, a total of 86 countries have reported
20 evidence of Zika infection, considering the outbreak of 2015-2016 as a pandemic (29).
21
22

23 In mid-January 2016, the health ministers of multiple Latin American countries recommended
24 postponing pregnancy for at least 6-24 months in facing the Zika virus. This seemed too idealistic, as
25 more than half of the region's pregnancies are unintended (30). In addition to that, inadequate sex
26 education, difficulty accessing contraception, high rape prevalence, and local cultural constraints resulted
27 in many women not adopting proper advice (31). Also, Latin American countries have a relatively wide
28 range of laws in SRHR. In El-Salvador, a self-induced abortion may lead to a penalty as high as 40 years
29 of incarceration (32).
30

31 On the other hand, in response to the Zika virus, the Columbian Ministry of health declared that women
32 have the right to have self-induced miscarriages. Nevertheless, due to the poor education and scarcity of
33 information, women are not well informed concerning the risk and their abortion rights (31). This was
34 reflected by the fact that the rate of contraception use did not change in Columbia during the outbreak in
35 2015-2016 (33).
36
37

38
39
40 **C) Ebola virus :**

41 Ebola hemorrhagic fever (EHF) is one of the fatal outbreaks within history. By the end of 2016, a study
42 showed that about 11310 deaths resulted from 28618 definite cases, a staggering 40% (34). Of 10
43 patients, almost nine of them died in those with critical clinical pictures. The transmission route was via
44 exposure to infected patients with each of the following blood, direct contact (mucus membrane and
45 injured cutaneous tissue), and secretions like semen. It has shown that, even after recovery, the virus may
46 exist in the semen of inflicted cases (35). Studies also showed that Burial ceremonies that require direct
47 contact with the dead body are also contributing to Ebola transmission(36). The latter endangers females
48 as they are usually doing significant roles in the ceremony. Besides, pregnant women can infect their
49 babies as their milk is possibly contaminated with the virus(37-39).
50
51

52 In the past century, multiple outbreaks of the Ebola virus occurred; the first one was in Zaire (currently
53 known as the Democratic Republic of the Congo) and Sudan in 1976. The second outbreak occurred in
54 Susan around 1979. The third epidemic was in Gabon,1996. Coming to this century, Northern Uganda
55
56
57
58
59
60

faced the fourth epidemic in the fall of 2000. The 2014-2016 epidemic of the Ebola virus was the largest outbreak of the Ebola, considered a pandemic (40, 41). The outset was from Guinea and then moved to other countries such as Liberia and Sierra Leone. The current outbreak (2018- now) in the eastern Democratic Republic of Congo (DRC) is the last reported outbreak(42).

WHO Advisory Group on the Ebola Virus Disease Response recommended that the male who survived the EVD should practice hygiene and safer sex until one year from symptom beginning, or after two negative semen specimen for the Ebola virus. Moreover, in some patients after recovery, Ebola Virus can persist in several tissues, such as testicles, the eye, and the central nervous system (43). In women who have contracted the disease during pregnancy, the virus remains in the placenta, amniotic fluid, and fetus. In those women who have been infected during breastfeeding, the virus could persist in breast milk(37, 44-49).

The Ebola has just more uncovered many problems in adolescence SRHR in Africa (50). It is well-established that in humanitarian crises and disasters, like Ebola, women and girls are more vulnerable to gender-based violence (GBV) (51, 52). Multiple studies showed that in Sierra Leone, in the Ebola outbreak, the rate of adolescent pregnancy, rape, the sexual and gender-based violation was increased (50, 53, 54). UNDP estimates that the teenage pregnancy rate increased by 65% during the outbreak (55). More girls have been forced into prostitution/transactional sex due to family members' loss and financial insecurity(56). This makes the girls twice victim, as pregnant girls are forbidden to attend school in Liberia and Sierra Leone (57). In Sierra-Leone, the Ebola increased the fear and affected the newborn and maternal and care by disrupting the health services. This led to an estimated 3600 death of stillbirth, maternal and newborn death (58). The studies showed that even after pandemics, the level of contraceptive care and family planning had returned or exceeded the baseline level after 6 (59) to 24 months (60). After six months of the outbreak in Guinea, the number of family planning sessions and prenatal care visits did not reverse to the prior level, harshly influencing the already insufficient care level (59).

D) SARS COV-2 :

In late 2019, multiple patients were diagnosed with a pneumonia-causing virus. Numerous subsequent studies showed a beta-coronidae family virus named "SARS-COV-2" by WHO to be the causative agent. In March 2020, the WHO declared the pandemic of the SARS-COV-2 and called this condition "coronavirus disease of 2019", or simply "COVID-19". Currently, there is no substantial evidence that the virus can transmit during sexual activity or breastfeeding of a newborn. The possibility of vertical transmission is controversial. However, a recently published systematic review and meta-analysis suggest that the minority of pregnant cases can transmit the virus to newborns (61).

As the COVID-19 escalated, many countries adopted a high level of lockdown to mitigate the virus spread. The pandemic disrupted the supply chain of key contraceptive commodities (62). Besides, in practice, most governments categorized SRH services as non-essential and forced them to close. This led to the closure of many clinics and other related health care facilities for adolescents. In late April 2020, UNFPA predicted around 7 million unintended pregnancies during the crisis, with possibly thousands of maternal and newborn maternal mortality and morbidity(63). This is because of the lack of providing SRH services like contraception and safe abortion care, time-sensitive, and potentially lifesaving services (64).

COVID-19 has forced nearly 1.4 billion child and youths out of school/university, including 743 million girls globally (65), raising concerns around long-term impacts on their lives, who, in addition to increasing poverty levels, might also experience increased sexual and gender-based violence, furthering the need not only for remediation and support services but also access to emergency contraception and other reproductive health services (66). COVID-19 has impacted health systems and services severely globally. It has also led to educational institutions' closure, public places for recreation, leisure, and impacted movement (67). The protracted closures across the world have led to increasing concerns around mental well-being and the availability/ accessibility of critical sexual and reproductive health services for adolescents in LMICs (68).

The Guttmacher institute predicted that with a hypothetical one-year 10% decline in the use of short and long-acting contraceptives in LMICs during the COVID-19 pandemic, 49 million women would lose their access to health services. They estimated that around 15 million unintended pregnancies might occur. This number can also lead to an additional 1.7 million delivery, 2.6 million newborns with significant complications, 168000 newborn deaths, and 28000 maternal mortality. The second hypothesis was that with a yearly 10% shift of safe abortion to unsafe abortion, an additional 3.3 unsafe abortions might occur in LMICs, and 1000 other maternal deaths in expected (62). We know that the impact is possibly more than a 10% decline in such services, as frontline partners have predicted a slip could be up to 80% (69). In India, by March 2020, compared to December 2019, a 36% decrease in injectable contraceptives and a 21% reduction in IUD insertion rate were reported. Simultaneously, the distribution of the condoms and oral cycle pills dipped 23% and 15%, respectively. The COVID-19 has interrupted the prevention programs and impacted the household economic status, resulting in 13 million child marriages and two million female genital mutilations in the next decade(53). The health record analysis showed a 68% reduction in HPV vaccination from February to early April 2020(70).

We aim to conduct a scoping review to map the range, extent, and nature of articles related to the effects of H1N1, Ebola virus, Zika virus, and SARS-COV-2 on sexual health and rights (SRHR) adolescents and youth in LMICs. Our goal is to identify current evidence in the literature, identify research gaps, and suggest future applicable types of research. The details pertaining to population, exposure, and outcomes are provided in table 1.

Objectives

The key research questions are:

- What is the existing evidence regarding the impact of the Zika virus outbreak on adolescents and youth SRHR in LMICs?
- What is the existing evidence regarding the impact of the H1N1/09 virus outbreak on adolescents and youth SRHR in LMICs?
- What is the existing evidence regarding the impact of the Ebola virus outbreak on adolescents and youth SRHR in LMICs?
- What is the existing evidence regarding the impact of the SARS-COV-2 virus outbreak on adolescents and youth SRHR in LMICs?

Table1. PICOS format in this review

PICOS framework for Systematic Reviews	
Population	Adolescents and youths living in LMICs ages 10-24 years
Intervention/exposure	SARS-COV-2, Zika, H1N1, and Ebola-related disruptions
Comparative /Control intervention	n/a
Outcome	SRHR (sexual well-being, sexual health and illness outcomes, reproductive health and illnesses, sexual and reproductive health services outcomes)
Study designs	All types of original articles

Method

We will use *PRISMA*-ScR (Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews) Statement to report the findings of our review (71).

Eligibility Criteria

Study designs:

We will include cross-sectional studies, case-control, cohort studies, clinical trials, and qualitative studies.

Participants/Population:

In the current review for full inclusiveness, we consider those aged 10-24 as adolescents and youth. Studies about youth and adolescents' SRHR) in LMICs are being investigated to study the impact of the outbreak of the Zika virus, Ebola virus, H1N1, and SARS-COV-2 on this target population.

Exposure/Intervention:

This study is designed to exclusively assess the impact of the Ebola virus, Zika virus, H1N1, and SARS-COV-2 on adolescents and youth SRHR in LMICs. The papers published since 2009 will be included.

Control:

There is no control group in this review.

Outcome

We will focus on the SRHR of adolescents and youth in LMICs. SRHR would consist of sexual well-being, sexual health, and illness outcomes, reproductive health, rights and illnesses, sexual and reproductive health services outcomes.

Our proposed outcome indicators inspired by the International Conference on Population and Development (ICPD) Programme of Action and sustainable development goal (SDG) 2030 (72, 73) are summarized in table 2.

Table2. The outcomes in our review

Category of outcome	Sub-category
A) Contraception	<ul style="list-style-type: none">- The proportion of adolescent and youth have access to contraceptive agent or procedures during the outbreaks
B) SRH Service Availability	<ul style="list-style-type: none">- The proportion of facilities providing SRHR services for adolescents and youth during the outbreaks- The proportion of health facilities providing post-abortion care and post-partum care for adolescent and youth and also services for those who have contracted HIV in our target group during the outbreaks
C) Knowledge About SRHR	<ul style="list-style-type: none">- The proportion of adolescent and youth have basic knowledge on SRHR during the pandemics
D) Adolescent Fertility	<ul style="list-style-type: none">- The adolescents and youth’s birth rate during the outbreaks- The proportion of pregnancies related to those 24 years old and below during the outbreaks
E) Quality of Care, Including Respect for Rights	<ul style="list-style-type: none">- The proportion of females are knowledgeable about side effects of their contraceptive method of choice and how to deal with adverse effects and also are familiar with other methods of contraception during the outbreaks- The universal access to contraceptives and SRHR information during outbreaks in national policy actions during the outbreaks- An indicator reflective of respectful care and human rights in the provision of SRH information and services during the outbreaks
F) Prevention of Sexually Transmitted Infections	<ul style="list-style-type: none">- The rate of HPV vaccination during the epidemics
G) Abortions:	<ul style="list-style-type: none">- Number of unsafe abortions during outbreaks- Number of facilities provide safe abortions services; or if it’s illegal, the number of facilities providing the services when unsafe abortions become complicated during the outbreaks
H) Comprehensive Sexuality Education	<ul style="list-style-type: none">- The proportion of school/facilities providing comprehensive sexual education during the outbreaks
I) Gender Equality in SRHR	<ul style="list-style-type: none">- whether adolescents and youth’s sexual autonomy within marriage is respected during the outbreaks

Timing:

The first significant outbreak of the third millennium was the H1N1/09 (swine flu) in 2009; hence, all the papers published since 2009 until 2021 will be included.

Setting:

Our study will focus on LMICs. Low- and middle-income countries (LMICs) were defined according to the 2020-2021 World Bank classification(74).

Language:

Our review compiles articles with no language restriction. We will use Google Translate for the primary translation of abstracts of non-English articles. In case the abstract of the article fits the scope of this review, we will request the experts in SRHR familiar with that specific language, e.g., Mandarin, to fully evaluate the paper based on the inclusion/exclusion criteria and finally extract the data. Alternatively, we will request official language translation centers to make the English version of the article for our evaluation.

Exclusion criteria

- all non-original papers will not be included.
- Those irrelevant to the SRHR
- Studies that do not include adolescents and youth (10-24 years old)
- Studies from countries not categorized as LMICs
- Those not relevant to Zika virus, Ebola Virus, H1N1/09, and SARS-COV-2 outbreaks.
- Full text is not available for evaluation
- If the sample size of a study would be a mixture of other ages besides 10-24, we will include it if data is disaggregated by age group or sub-group analysis on age 10-24. If they have included vivid results about this age group, we will consist of their study.
- Animal studies

Information Sources:

We will look at four databases in our review: Pubmed, Cochrane Library, Embase, and PsycINFO.

Search:

Table 3 (appendix 1) shows our proposed search strategy in Pubmed. Table 4-6 (appendix 1) is our search strategy for each of the other target databases (Cochrane Library, Embase, PsycINFO)

Selection of sources of evidence:

A four-step search strategy is being followed in this study. First, the authors have done a preliminary (and limited) search in several databases (Pubmed and Google Scholar) on this topic. We assessed the keywords, abstract, and several full texts to reach our study's final keywords. We discussed our search strategy multiple times until a consensus was reached by all authors reflected in the table of 3-8.

Following the duplication removal, title/abstract, and full-text screening of studies using RAYYAN (75), we will choose our relevant evidence considering the inclusion/exclusion criteria.

In the third step, we will check the articles' reference list to find more relevant studies. Finally, we will look for studies that have cited or included studies and check them for their eligibility. Any discrepancy would be addressed via discussion until consensus is reached in each stage. Two authors worked separately in all searching and extraction stages, and discrepancies would be resolved through discussion.

Data Charting Process:

After choosing the eligible papers, two authors will separately extract the required items based on the extraction excel sheet form designed and approved by researchers for the process.

Data items:

Following the selected papers' primary selection and inclusion and extracting the results, the tabulation phase begins. We will categorize the extracted data based on the items mentioned before. We will extract the following data: authors, year, country and continent (region of the study), study design, participants (differentiated by age and gender), purpose, and main findings. Any disagreements in the process of data extraction will be resolved through discussion until consensus. We will also indicate the country's income level where studies are being done (low, lower-middle, and higher-middle income).

Critical appraisal of individual sources of evidence and synthesis of result

Based on the nature of scoping review, we will neither synthesize the findings/results, nor critically appraise the papers, which is more applicable by performing a systematic review(76). However, publications and the mentioned data would be summarized and categorized.

Ethics and dissemination

As the nature of the work is a scoping review of the currently published papers in peer review journals, ethical approval is not required for this study. The results of the study will be presented in a peer-reviewed publication.

Authors' contribution: HA and MK developed the concept and wrote the early draft. RS,PS,MHT,PC,MRW,SK,IN edited the early draft. All authors approved the final version.

Funding: The review received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interest: none

References

1. Chandra-Mouli V, McCarraher DR, Phillips SJ, Williamson NE, Hainsworth G. Contraception for adolescents in low and middle income countries: needs, barriers, and access. *Reproductive health*. 2014;11(1):1.
 2. Blanc AK, Tsui AO, Croft TN, Trevitt JLIpos, health r. Patterns and trends in adolescents' contraceptive use and discontinuation in developing countries and comparisons with adult women. 2009:63-71.
 3. WHO. Adolescent pregnancy. Available at <https://www.who.int/en/news-room/fact-sheets/detail/adolescent-pregnancy> . Accessed 27th Dec 2020. .
 4. Hodgkinson SC, Colantuoni E, Roberts D, Berg-Cross L, Belcher HMJJop, gynecology a. Depressive symptoms and birth outcomes among pregnant teenagers. 2010;23(1):16-22.
 5. Raj A, Boehmer UJVaw. Girl child marriage and its association with national rates of HIV, maternal health, and infant mortality across 97 countries. 2013;19(4):536-51.
 6. Organization WH. Global standards for quality health-care services for adolescents: a guide to implement a standards-driven approach to improve the quality of health care services for adolescents. 2015.
 7. Sully E. Adding it up: investing in sexual and reproductive health 2019. New York: Guttmacher Institute; 2020. 2019.
 8. Darroch JE, Woog V, Bankole A, Ashford LS, Points KJGI. Costs and benefits of meeting the contraceptive needs of adolescents. 2016.
 9. Gore FM, Bloem PJ, Patton GC, Ferguson J, Joseph V, Coffey C, et al. Global burden of disease in young people aged 10–24 years: a systematic analysis. 2011;377(9783):2093-102.
 10. Neal S, Matthews Z, Frost M, Fogstad H, Camacho AV, Laski LJAoegS. Childbearing in adolescents aged 12–15 years in low resource countries: a neglected issue. New estimates from demographic and household surveys in 42 countries. 2012;91(9):1114-8.
 11. Guessoum SB, Lachal J, Radjack R, Carretier E, Minassian S, Benoit L, et al. Adolescent psychiatric disorders during the COVID-19 pandemic and lockdown. 2020:113264.
 12. Briand S, editor Epidemiology and illness severity of pandemic (H1N1) 09 virus. World Health Organization Strategic Advisory Group of Experts on Immunization Extraordinary Meeting, Geneva, Switzerland; 2009.
 13. WHO Guidelines Approved by the Guidelines Review Committee. Pregnancy and Pandemic Influenza A (H1N1) 2009: Information for Programme Managers and Clinicians. Geneva: World Health Organization
- Copyright © World Health Organization 2010.; 2010.
14. Somerville LK, Basile K, Dwyer DE, Kok JJFm. The impact of influenza virus infection in pregnancy. 2018;13(2):263-74.
 15. Medicine AIJNEJo. Critical care services and 2009 H1N1 influenza in Australia and New Zealand. 2009;361(20):1925-34.
 16. Van Kerkhove MD, Vandemaele KA, Shinde V, Jaramillo-Gutierrez G, Koukounari A, Donnelly CA, et al. Risk factors for severe outcomes following 2009 influenza A (H1N1) infection: a global pooled analysis. 2011;8(7):e1001053.
 17. Drees M, Johnson O, Wong E, Stewart A, Ferisin S, Silverman PR, et al. Acceptance of 2009 H1N1 influenza vaccine among pregnant women in Delaware. 2012;29(04):289-94.
 18. McCarthy EA, Pollock WE, Nolan T, Hay S, McDonald SJA, Obstetrics NZJo, et al. Improving influenza vaccination coverage in pregnancy in Melbourne 2010–2011. 2012;52(4):334-41.

19. McMillan M, Porritt K, Kralik D, Costi L, Marshall H. Influenza vaccination during pregnancy: a systematic review of fetal death, spontaneous abortion, and congenital malformation safety outcomes. *Vaccine*. 2015;33(18):2108-17.

20. Goldfarb I, Panda B, Wylie B, Riley LJAjoo, gynecology. Uptake of influenza vaccine in pregnant women during the 2009 H1N1 influenza pandemic. 2011;204(6):S112-S5.

21. Conlin AMS, Bukowinski AT, Seveck CJ, DeScisciolo C, Crum-Cianflone NFJO, Gynecology. Safety of the pandemic H1N1 influenza vaccine among pregnant US military women and their newborns. 2013;121(3):511-8.

22. WHO. Zika Virus. Available from <https://www.who.int/news-room/fact-sheets/detail/zika-virus> . Accessed 28th Dec 2020. .

23. Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Da Fonseca EB, et al. Characterizing the pattern of anomalies in congenital Zika syndrome for pediatric clinicians. 2017;171(3):288-95.

24. Mansuy JM, Dutertre M, Mengelle C, Fourcade C, Marchou B, Delobel P, et al. Zika virus: high infectious viral load in semen, a new sexually transmitted pathogen? 2016;16(4):405.

25. Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau V-MJ. Potential sexual transmission of Zika virus. 2015;21(2):359.

26. Brent CJMM, report mw. Preliminary findings from an investigation of Zika virus infection in a patient with no known risk factors—Utah, 2016. 2016;65.

27. Moghadam SRJ, Bayrami S, Moghadam SJ, Golrokhi R, Pahlaviani FG, SeyedAlinaghi SJAPJoTB. Zika virus: A review of literature. 2016;6(12):989-94.

28. Katz LM, Rossmann SNJAop, medicine I. Zika and the blood supply: a work in progress. 2017;141(1):85-92.

29. Chapter 6 - Comparing the Zika Virus Disease Pandemic to Other Disease Pandemics. In: Qureshi AI, editor. *Zika Virus Disease*: Academic Press; 2018. p. 85-96.

30. Sedgh G, Singh S, Hussain RJSifp. Intended and unintended pregnancies worldwide in 2012 and recent trends. 2014;45(3):301-14.

31. Roa MJTL. Zika virus outbreak: reproductive health and rights in Latin America. 2016;387(10021):843.

32. Viterna J, Bautista JSGJH, Rights H. Pregnancy and the 40-year prison sentence: how “abortion is murder” became institutionalized in the Salvadoran judicial system. 2017;19(1):81.

33. Machado-Alba JE, Machado-Duque ME, Gaviria-Mendoza A, Orozco-Giraldo VAJTL. Hormonal contraceptive prescriptions in Colombia and Zika virus. 2016;387(10032):1993.

34. Coltart CE, Lindsey B, Ghinai I, Johnson AM, Heymann DLJPTotRSBBS. The Ebola outbreak, 2013–2016: old lessons for new epidemics. 2017;372(1721):20160297.

35. Ebola Haemorrhagic Fever. Retrieved from <https://www.msf.org/who-fact-sheet-ebola-haemorrhagic-fever> . Accessed in 29th Dec 2020.

36. Nkangu MN, Olatunde OA, Yaya SJIdop. The perspective of gender on the Ebola virus using a risk management and population health framework: a scoping review. 2017;6(1):135.

37. Medina-Rivera M, Centeno-Tablante E, Finkelstein JL, Rayco-Solon P, Peña-Rosas JP, Garcia-Casal MN, et al. Presence of Ebola virus in breast milk and risk of mother-to-child transmission: synthesis of evidence. 2020.

38. Nordenstedt H, Bah EI, de la Vega M-A, Barry M, N’Faly M, Barry M, et al. Ebola virus in breast milk in an Ebola virus–positive mother with twin babies, Guinea, 2015. 2016;22(4):759.

39. Sissoko D, Keita M, Diallo B, Aliabadi N, Fitter DL, Dahl BA, et al. Ebola virus persistence in breast milk after no reported illness: a likely source of virus transmission from mother to child. 2017;64(4):513-6.

40. Ifediora OF, Aning KJGGARoM, Organizations I. West Africa’s Ebola Pandemic: Toward Effective Multilateral Responses to Health Crises. 2017;23(2):225-44.

41. Zhong Y, Xu J, Li T, Yu X, Sheng MJSCLS. Potential clinical treatment for Ebola pandemic. 2014;57(10):982.
42. WHO. Ebola virus . Retrived from <https://www.who.int/news-room/fact-sheets/detail/ebola-virus-disease>. Accessed 29th DEC 2020. .
43. Racine T, Kobinger GP. Viral pathogenesis: Unlocking Ebola persistence. Nature Microbiology. 2017;2(9):17124.
44. Chughtai A, Barnes M, Macintyre CJE, Infection. Persistence of Ebola virus in various body fluids during convalescence: evidence and implications for disease transmission and control. 2016;144(8):1652-60.
45. Bausch DG, Towner JS, Dowell SF, Kaducu F, Lukwiya M, Sanchez A, et al. Assessment of the risk of Ebola virus transmission from bodily fluids and fomites. 2007;196(Supplement_2):S142-S7.
46. Baggi F, Taybi A, Kurth A, Van Herp M, Di Caro A, Wölfel R, et al. Management of pregnant women infected with Ebola virus in a treatment centre in Guinea, June 2014. 2014;19(49):20983.
47. Caluwaerts S, Fautsch T, Lagrou D, Moreau M, Modet Camara A, Günther S, et al. Dilemmas in managing pregnant women with Ebola: 2 case reports. 2016;62(7):903-5.
48. Oduyebo T, Pineda D, Lamin M, Leung A, Corbett C, Jamieson DJJO, et al. A pregnant patient with Ebola virus disease. 2015;126(6):1273-5.
49. Muehlenbachs A, de la Rosa Vázquez O, Bausch DG, Schafer IJ, Paddock CD, Nyakio JP, et al. Ebola virus disease in pregnancy: clinical, histopathologic, and immunohistochemical findings. 2017;215(1):64-9.
50. Denney L, Gordon R, Ibrahim AJLODI. Teenage Pregnancy after Ebola in Sierra Leone.'. 2015.
51. Hynes M, Ward J, Robertson K, Crouse CJD. A determination of the prevalence of gender-based violence among conflict-affected populations in East Timor. 2004;28(3):294-321.
52. Kofman YB, Garfin DRJPTT, Research, Practice,, Policy. Home is not always a haven: The domestic violence crisis amid the COVID-19 pandemic. 2020.
53. PAHO .Adolescent Pregnancy in Latin America and the Caribbean. Retrived from https://pancap.org/pc/pcc/media/pancap_event/Technical-brief-Adolescent-Pregnancy-in-Latin-America-and-the-Caribbean.pdf . Accessed 30th Dec 2020. . .
54. Elston J, Moosa A, Moses F, Walker G, Dotta N, Waldman RJ, et al. Impact of the Ebola outbreak on health systems and population health in Sierra Leone. 2016;38(4):673-8.
55. Werber CJQA. How Ebola led to more teenage pregnancy in West Africa. 2015.
56. Minor OMJAiA. Ebola and accusation: Gender dimensions of stigma in Sierra Leone's Ebola response. 2017;24(2):25-35.
57. Walsh S, Mulhern E. A Step in the Rights' Direction: Advocacy, Negotiation, and Money as Tools for Realising the Right to Education for Pregnant Girls in Sierra Leone During the Ebola Epidemic. Pregnant in the Time of Ebola: Springer; 2019. p. 399-416.
58. Sochas L, Channon AA, Nam SJHp, planning. Counting indirect crisis-related deaths in the context of a low-resilience health system: the case of maternal and neonatal health during the Ebola epidemic in Sierra Leone. 2017;32(suppl_3):iii32-iii9.
59. Camara BS, Delamou A, Diro E, Béavogui AH, El Ayadi AM, Sidibé S, et al. Effect of the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: an ecological study. 2017;111(1):22-9.
60. Bietsch K, Williamson J, Reeves MJSiFP. Family planning during and after the West African Ebola crisis. 2020;51(1):71-86.
61. Yang Z, Liu YJAjop. Vertical transmission of severe acute respiratory syndrome coronavirus 2: a systematic review. 2020;37(10):1055.
62. Riley T, Sully E, Ahmed Z, Biddlecom AJIPSRH. Estimates of the potential impact of the COVID-19 pandemic on sexual and reproductive health in low-and middle-income countries. 2020;46:46.

63. Impact of the COVID-19 Pandemic on Family Planning and Ending Gender-based Violence, Female Genital Mutilation and Child Marriage . UNFPA. Retrieved from <https://www.unfpa.org/resources/impact-covid-19-pandemic-family-planning-and-ending-gender-based-violence-female-genital>. Accessed 30 Dec 2020 . .

64. Cousins SJTL. COVID-19 has “devastating” effect on women and girls. 2020;396(10247):301-2.

65. COVID-19 School Closures Around the World Will Hit Girls Hardest . available from <https://plan-international.org/blog/2020/03/covid-19-school-closures-hit-girls-hardest> . 30th DEC 2020.

66. Nanda P, Tandon S, Khanna A. Virtual and essential – adolescent SRHR in the time of COVID-19. Sexual and Reproductive Health Matters. 2020;28(1):1831136.

67. Lee JJTLC, Health A. Mental health effects of school closures during COVID-19. 2020;4(6):421.

68. Mayurasakorn K, Pinsawas B, Mongkolsucharitkul P, Sranacharoenpong K, Damapong SnJJoP, Health C. School closure, COVID-19 and lunch programme: Unprecedented undernutrition crisis in low-middle income countries. 2020;56(7):1013-7.

69. Marie Stopes International, Methodology for calculating impact of COVID-19, 2020, <https://www.mariestopes.org/resources/methodology-for-calculating-impact-of-covid-19>. Accessed 30 Dec 2020.

70. Vaccine Rates Drop Dangerously as Parents Avoid Doctor’s Visits . <https://www.nytimes.com/2020/04/23/health/coronavirus-measles-vaccines.html> . Accessed 30 DEC 2020.

71. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. 2018;169(7):467-73.

72. International Conference on Population and Development Programme of Action . 2014 . available in : <https://www.unfpa.org/publications/international-conference-population-and-development-programme-action> . accessed date March 9th 2021. .

73. Barot S, Cohen S, Darroch J, Galati A, Polis C, Singh S, et al. Sexual and Reproductive Health and Rights Indicators for the SDGs. New York: Guttmacher Institute; 2015.

74. World Bank. Low and middle income countries. Retrived from <https://data.worldbank.org/country/XO> . Accessed 28th Dec 2020.

75. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. Systematic reviews. 2016;5(1):1-10.

76. Askarian M, Khakpour M, Taghrir MH, Akbarialiabad H, Borazjani RJJes. Investigating the epidemiology of the methanol poisoning outbreaks in the third millennium: a scoping review protocol. 2020.



PRISMA 2009 Flow Diagram for scoping reviews




Diagram1. Flow chart of the review phases

Table 2. data charting table

Authors/year	Country and income level	Continent (geographical region)	Sample size and Median Participant age (10-14), (15-19), (20-24), and the gender	Study design	Purpose	Main Findings

Table 3. Search in PUBMED/MEDLINE (*Title/Abstract for all options*)

PUBMED	<i>Search terms in Pubmed</i>
<i>Population</i>	<p> <i>("adolescen*"[Title/Abstract] OR "young"[Title/Abstract] OR "youth"[Title/Abstract] OR "teen"[Title/Abstract] OR "teenager"[Title/Abstract] OR "child"[Title/Abstract] OR "children"[Title/Abstract] OR "paediatr*"[Title/Abstract] OR "pediatr*"[Title/Abstract] OR "juvenil*"[Title/Abstract] OR "boy"[Title/Abstract] OR "boys"[Title/Abstract] OR "girl"[Title/Abstract] OR "girls"[Title/Abstract] OR "pubert*"[Title/Abstract] OR "pubescen*"[Title/Abstract] OR "Young Adult"[Title/Abstract] OR "Young Adults"[Title/Abstract] OR "Adolescent"[MeSH Terms] OR "Young Adult"[MeSH Terms])</i> </p>
	<i>AND</i>
<i>Exposure</i>	<p> <i>(((((("swine flu"[Title/Abstract] OR "H1N1"[Title/Abstract] OR "flu pandemic"[Title/Abstract] OR "Zika"[Title/Abstract] OR "Ebola"[Title/Abstract] OR "EVD"[Title/Abstract] OR "EHF"[Title/Abstract] OR "Ebola virus disease"[Title/Abstract] OR "Ebola hemorrhagic fever"[Title/Abstract] OR "COVID-19"[Title/Abstract] OR "SARS-COV-2"[Title/Abstract] OR "Novel Coronavirus"[Title/Abstract] OR "2019- nCov"[Title/Abstract])) OR ("SARS-CoV-2"[MeSH Terms])) OR ("hemorrhagic fever, ebola"[MeSH Terms])) OR (ebolavirus[MeSH Terms])) OR ("influenza a virus, h1n1 subtype"[MeSH Terms])) OR ("Zika virus"[MeSH Terms])</i> </p>
	<i>AND</i>
<i>Outcome (in SRHR)</i>	<p> <i>((("ASRH"[Title/Abstract] OR "SRH"[Title/Abstract] OR "SRHR"[Title/Abstract] OR "human right*"[Title/Abstract] OR "pregnan*"[Title/Abstract] OR "abort*"[Title/Abstract] OR "termination"[Title/Abstract] OR "antenatal"[Title/Abstract] OR</i> </p>

	<p><i>"postnatal"[Title/Abstract] OR</i> <i>"perinatal"[Title/Abstract] OR</i> <i>"couple*"[Title/Abstract] OR "abuse"[Title/Abstract]</i> <i>OR "intercourse"[Title/Abstract] OR</i> <i>"rape"[Title/Abstract] OR "coerci*"[Title/Abstract]</i> <i>OR "violence"[Title/Abstract] OR</i> <i>"violation"[Title/Abstract] OR "incest"[Title/Abstract]</i> <i>OR "emergency pills"[Title/Abstract] OR "Long-acting</i> <i>reversible contraceptive"[Title/Abstract] OR</i> <i>"LARC"[Title/Abstract] OR</i> <i>"Misoprostol"[Title/Abstract] OR</i> <i>"mifepristone"[Title/Abstract] OR</i> <i>"methotrexate"[Title/Abstract] OR</i> <i>"meteneprost"[Title/Abstract] OR</i> <i>"prostaglandin"[Title/Abstract] OR</i> <i>"lilopristone"[Title/Abstract] OR</i> <i>"onapristone"[Title/Abstract] OR "oxytocin"</i> <i>[Title/Abstract] OR "sulprostone"[Title/Abstract] OR</i> <i>"family planning"[Title/Abstract] OR "sex</i> <i>education"[Title/Abstract] OR "STI"[Title/Abstract]</i> <i>OR "sexually transmitted infection"[Title/Abstract] OR</i> <i>"HPV vaccination"[Title/Abstract] OR "intimate</i> <i>partner violence"[Title/Abstract] OR "female genital</i> <i>mutilation"[Title/Abstract] OR "female genital</i> <i>cutting"[Title/Abstract] OR "FGM"[Title/Abstract] OR</i> <i>"maternal death"[Title/Abstract] OR "maternal</i> <i>morbidity"[Title/Abstract] OR "maternal</i> <i>mortality"[Title/Abstract] OR "newborn</i> <i>morbidity"[Title/Abstract] OR "newborn</i> <i>mortality"[Title/Abstract] OR "maternal</i> <i>care"[Title/Abstract] OR "newborn</i> <i>care"[Title/Abstract] OR "infant"[Title/Abstract] OR</i> <i>"infancy"[Title/Abstract] OR "stigma"[Title/Abstract]</i> <i>OR "taboo"[Title/Abstract] OR</i> <i>"Discrimination"[Title/Abstract] OR</i> <i>"GBV"[Title/Abstract] OR "gender"[Title/Abstract]</i> <i>OR "VAWG"[Title/Abstract] OR (sexual and gender</i> <i>disorders[MeSH Terms])) OR (sexual and gender</i> <i>minorities[MeSH Terms])) OR ("Adolescent sexual</i> <i>and reproductive health" OR "sexual and reproductive</i> <i>health" OR "Menstruation-Inducing Agents")</i></p>
	<p>AND</p>

Low and middle-income countries (96)

((("low income countr*" [Title/Abstract] OR "middle income countr*" [Title/Abstract] OR "low middle income countr*" [Title/Abstract] OR "developing countr*" [Title/Abstract] OR "middle income countr*" [Title/Abstract] OR "LMIC" [Title/Abstract] OR "upper middle income countr*" [Title/Abstract] OR "Albania" [Title/Abstract] OR "American Samoa" [Title/Abstract] OR "Argentina" [Title/Abstract] OR "Armenia" [Title/Abstract] OR "Azerbaijan" [Title/Abstract] OR "Belarus" [Title/Abstract] OR "Belize" [Title/Abstract] OR "bosnia" [Title/Abstract] OR "Herzegovina" [Title/Abstract] OR "Botswana" [Title/Abstract] OR "Brazil" [Title/Abstract] OR "Bulgaria" [Title/Abstract] OR "China" [Title/Abstract] OR "Colombia" [Title/Abstract] OR "Costa Rica" [Title/Abstract] OR "Cuba" [Title/Abstract] OR "Dominica" [Title/Abstract] OR "Dominican Republic" [Title/Abstract] OR "Ecuador" [Title/Abstract] OR "Equatorial Guinea" [Title/Abstract] OR "Fiji" [Title/Abstract] OR "Gabon" [Title/Abstract] OR "Georgia" [Title/Abstract] OR "Grenada" [Title/Abstract] OR "Guatemala" [Title/Abstract] OR "Guyana" [Title/Abstract] OR "Indonesia" [Title/Abstract] OR "Iran" [Title/Abstract] OR "Iraq" [Title/Abstract] OR "Jamaica" [Title/Abstract] OR "Jordan" [Title/Abstract] OR "Kazakhstan" [Title/Abstract] OR "Kosovo" [Title/Abstract] OR "Lebanon" [Title/Abstract] OR "Libya" [Title/Abstract] OR "Malaysia" [Title/Abstract] OR "Maldives" [Title/Abstract] OR "Marshall Islands" [Title/Abstract] OR "Mexico" [Title/Abstract] OR "Montenegro" [Title/Abstract] OR "Namibia" [Title/Abstract] OR "North Macedonia" [Title/Abstract] OR "Paraguay" [Title/Abstract] OR "Peru" [Title/Abstract] OR "Russia" [Title/Abstract] OR "Samoa" [Title/Abstract] OR "Serbia" [Title/Abstract] OR "South Africa" [Title/Abstract] OR "Saint Lucia" [Title/Abstract] OR "saint Vincent" [Title/Abstract] OR "grenadines" [Title/Abstract] OR

	<p>"Suriname"[Title/Abstract] OR "Thailand"[Title/Abstract] OR "Tonga"[Title/Abstract] OR "Turkey"[Title/Abstract] OR "Turkmenistan"[Title/Abstract] OR "Tuvalu"[Title/Abstract] OR "Venezuela"[Title/Abstract] OR "lower middle income countr*"[Title/Abstract] OR "Algeria"[Title/Abstract] OR "Angola"[Title/Abstract] OR "Bangladesh"[Title/Abstract] OR "Benin"[Title/Abstract] OR "Bhutan"[Title/Abstract] OR "Bolivia"[Title/Abstract] OR "Cabo Verde"[Title/Abstract] OR "Cambodia"[Title/Abstract] OR "Cameroon"[Title/Abstract] OR "Comoros"[Title/Abstract] OR "Congo"[Title/Abstract] OR "Cote d'Ivoire"[Title/Abstract] OR "Djibouti"[Title/Abstract] OR "Egypt"[Title/Abstract] OR "El Salvador"[Title/Abstract] OR "Eswatini"[Title/Abstract] OR "Ghana"[Title/Abstract] OR "Honduras"[Title/Abstract] OR "India"[Title/Abstract] OR "Kenya"[Title/Abstract] OR "Kiribati"[Title/Abstract] OR "Kyrgyz"[Title/Abstract] OR "Laos"[Title/Abstract] OR "Lesotho"[Title/Abstract] OR "Mauritania"[Title/Abstract] OR "Micronesia"[Title/Abstract] OR "Moldova"[Title/Abstract] OR "Mongolia"[Title/Abstract] OR "Morocco"[Title/Abstract] OR "Myanmar"[Title/Abstract] OR "Nepal"[Title/Abstract] OR "Nicaragua"[Title/Abstract] OR "Nigeria"[Title/Abstract] OR "Pakistan"[Title/Abstract] OR "Papua New Guinea"[Title/Abstract] OR "Philippines"[Title/Abstract] OR "Sao Tome"[Title/Abstract] OR "Principe"[Title/Abstract] OR "Senegal"[Title/Abstract] OR "Solomon Islands"[Title/Abstract] OR "Sri Lanka"[Title/Abstract] OR "Tanzania"[Title/Abstract] OR "Timor-Leste"[Title/Abstract] OR "Tunisia"[Title/Abstract] OR "Ukraine"[Title/Abstract] OR "Uzbekistan"[Title/Abstract] OR "Vanuatu"[Title/Abstract] OR "Vietnam"[Title/Abstract] OR "West</p>
--	--

Bank"[Title/Abstract] OR "Gaza"[Title/Abstract] OR "Zambia"[Title/Abstract] OR "Zimbabwe"[Title/Abstract] OR "low-income country"[Title/Abstract] OR "Afghanistan"[Title/Abstract] OR "Burkina Faso"[Title/Abstract] OR "Burundi"[Title/Abstract] OR "Central African Republic"[Title/Abstract] OR "Chad"[Title/Abstract] OR "Congo"[Title/Abstract] OR "Eritrea"[Title/Abstract] OR "Ethiopia"[Title/Abstract] OR "Gambia"[Title/Abstract] OR "Guinea"[Title/Abstract] OR "Guinea-Bissau"[Title/Abstract] OR "North Korea"[Title/Abstract] OR "Korea Democratic People's Republic"[Title/Abstract] OR "Haiti"[Title/Abstract] OR "Liberia"[Title/Abstract] OR "Madagascar"[Title/Abstract] OR "Malawi"[Title/Abstract] OR "Mali"[Title/Abstract] OR "Mozambique"[Title/Abstract] OR "Niger"[Title/Abstract] OR "Rwanda"[Title/Abstract] OR "Sierra Leone"[Title/Abstract] OR "Somalia"[Title/Abstract] OR "South Sudan"[Title/Abstract] OR "Sudan"[Title/Abstract] OR "Syrian Arab Republic"[Title/Abstract] OR "Syria"[Title/Abstract] OR "Tajikistan"[Title/Abstract] OR "Togo"[Title/Abstract] OR "Yemen"[Title/Abstract]) OR ("africa"[MeSH Terms])) OR ("central america"[MeSH Terms])) OR ("middle east"[MeSH Terms]))

Table 4. terms used in Embase
Abstract, title, and keywords for all option

Search	Search terms
Population	'adolescen*':ti,ab,kw OR 'young':ti,ab,kw OR 'youth':ti,ab,kw OR 'teen':ti,ab,kw OR 'teens':ti,ab,kw OR 'teenag*':ti,ab,kw OR 'child*':ti,ab,kw OR 'paediatr*':ti,ab,kw OR 'pediatr*':ti,ab,kw OR 'juvenil*':ti,ab,kw OR 'boy':ti,ab,kw OR 'boys':ti,ab,kw OR 'girl':ti,ab,kw OR 'girls':ti,ab,kw OR 'pubert*':ti,ab,kw OR 'pubescen*':ti,ab,kw
	AND
Exposure	'swine flu':ti,ab,kw OR 'h1n1':ti,ab,kw OR 'flu pandemic':ti,ab,kw OR 'zika':ti,ab,kw OR 'ebola':ti,ab,kw OR 'evd':ti,ab,kw OR 'ehf':ti,ab,kw OR 'ebola virus disease':ti,ab,kw OR 'ebola hemorrhagic fever':ti,ab,kw OR 'covid-19':ti,ab,kw OR 'sars-cov-2':ti,ab,kw OR 'novel coronavirus':ti,ab,kw OR '2019-ncov':ti,ab,kw
	AND
Outcome (in SRHR)	asrh:ti,ab,kw OR 'srh':ti,ab,kw OR 'srhr':ti,ab,kw OR 'human right*':ti,ab,kw OR 'pregnan*':ti,ab,kw OR 'abort*':ti,ab,kw OR 'termination':ti,ab,kw OR 'antenatal':ti,ab,kw OR 'postnatal':ti,ab,kw OR 'perinatal':ti,ab,kw OR 'couple*':ti,ab,kw OR 'abuse':ti,ab,kw OR 'intercourse':ti,ab,kw OR 'rape':ti,ab,kw OR 'coerci*':ti,ab,kw OR 'violence':ti,ab,kw OR 'violation':ti,ab,kw OR 'incest':ti,ab,kw OR 'emergency pills':ti,ab,kw OR 'long-acting reversible contraceptive':ti,ab,kw OR 'larc':ti,ab,kw OR 'misoprostol':ti,ab,kw OR 'mifepristone':ti,ab,kw OR 'methotrexate':ti,ab,kw OR 'meteneprost':ti,ab,kw OR 'prostaglandin':ti,ab,kw OR 'lilopristone':ti,ab,kw OR 'onapristone':ti,ab,kw OR oxytocin:ti,ab,kw OR 'sulprostone':ti,ab,kw OR 'family planning':ti,ab,kw OR 'sex education':ti,ab,kw OR 'sti':ti,ab,kw OR 'sexually transmitted infection':ti,ab,kw OR 'hvp vaccination':ti,ab,kw OR 'intimate partner violence':ti,ab,kw OR 'female genital mutilation':ti,ab,kw OR 'female genital cutting':ti,ab,kw OR 'fgm':ti,ab,kw OR 'maternal death':ti,ab,kw OR 'maternal morbidity':ti,ab,kw OR

	'maternal mortality':ti,ab,kw OR 'newborn morbidity':ti,ab,kw OR 'newborn mortality':ti,ab,kw OR 'maternal care':ti,ab,kw OR 'newborn care':ti,ab,kw OR 'infant':ti,ab,kw OR 'infancy':ti,ab,kw OR 'stigma':ti,ab,kw OR 'taboo':ti,ab,kw OR 'discrimination':ti,ab,kw OR 'gbv':ti,ab,kw OR 'gender':ti,ab,kw
AND	
<i>Low and middle-income countries (96)</i>	'low income countr*':ti,ab,kw OR 'low middle income countr*':ti,ab,kw OR 'developing countr*':ti,ab,kw OR 'middle income countr*':ti,ab,kw OR 'lmic':ti,ab,kw OR 'upper middle income countr*':ti,ab,kw OR 'albania':ti,ab,kw OR 'american samoa':ti,ab,kw OR 'argentina':ti,ab,kw OR 'armenia':ti,ab,kw OR 'azerbaijan':ti,ab,kw OR 'belarus':ti,ab,kw OR 'belize':ti,ab,kw OR 'bosnia':ti,ab,kw OR 'herzegovina':ti,ab,kw OR 'botswana':ti,ab,kw OR 'brazil':ti,ab,kw OR 'bulgaria':ti,ab,kw OR 'china':ti,ab,kw OR 'colombia':ti,ab,kw OR 'costa rica':ti,ab,kw OR 'cuba':ti,ab,kw OR 'dominica':ti,ab,kw OR 'dominican republic':ti,ab,kw OR 'ecuador':ti,ab,kw OR 'equatorial guinea':ti,ab,kw OR 'fiji':ti,ab,kw OR 'gabon':ti,ab,kw OR 'georgia':ti,ab,kw OR 'grenada':ti,ab,kw OR 'guatemala':ti,ab,kw OR 'guyana':ti,ab,kw OR 'indonesia':ti,ab,kw OR 'iran':ti,ab,kw OR 'iraq':ti,ab,kw OR 'jamaica':ti,ab,kw OR 'jordan':ti,ab,kw OR 'kazakhstan':ti,ab,kw OR 'kosovo':ti,ab,kw OR 'lebanon':ti,ab,kw OR 'libya':ti,ab,kw OR 'malaysia':ti,ab,kw OR 'maldives':ti,ab,kw OR 'marshall islands':ti,ab,kw OR 'mexico':ti,ab,kw OR 'montenegro':ti,ab,kw OR 'namibia':ti,ab,kw OR 'north macedonia':ti,ab,kw OR 'paraguay':ti,ab,kw OR 'peru':ti,ab,kw OR 'russia':ti,ab,kw OR 'samoa':ti,ab,kw OR 'serbia':ti,ab,kw OR 'south africa':ti,ab,kw OR 'saint lucia':ti,ab,kw OR 'saint vincent':ti,ab,kw OR 'grenadines':ti,ab,kw OR 'suriname':ti,ab,kw OR 'thailand':ti,ab,kw OR 'tonga':ti,ab,kw OR 'turkey':ti,ab,kw OR 'turkmenistan':ti,ab,kw OR 'tuvalu':ti,ab,kw OR 'venezuela':ti,ab,kw OR 'lower middle income countr*':ti,ab,kw OR 'algeria':ti,ab,kw

	OR 'angola':ti,ab,kw OR 'bangladesh':ti,ab,kw OR 'benin':ti,ab,kw OR 'bhutan':ti,ab,kw OR 'bolivia':ti,ab,kw OR 'cabo verde':ti,ab,kw OR cambodia:ti,ab,kw OR 'cameroon':ti,ab,kw OR 'comoros':ti,ab,kw OR 'cote divoire':ti,ab,kw OR 'Ivory Coast ':ti,ab,kw OR 'djibouti':ti,ab,kw OR 'egypt':ti,ab,kw OR 'el salvador':ti,ab,kw OR 'eswatini':ti,ab,kw OR 'ghana':ti,ab,kw OR 'honduras':ti,ab,kw OR 'india':ti,ab,kw OR 'kenya':ti,ab,kw OR 'kiribati':ti,ab,kw OR 'kyrgyz':ti,ab,kw OR 'laos':ti,ab,kw OR 'lesotho':ti,ab,kw OR 'mauritania':ti,ab,kw OR 'micronesia':ti,ab,kw OR 'moldova':ti,ab,kw OR 'mongolia':ti,ab,kw OR 'morocco':ti,ab,kw OR 'myanmar':ti,ab,kw OR 'nepal':ti,ab,kw OR 'nicaragua':ti,ab,kw OR 'nigeria':ti,ab,kw OR 'pakistan':ti,ab,kw OR 'papua new guinea':ti,ab,kw OR 'philippines':ti,ab,kw OR 'sao tome':ti,ab,kw OR 'principe':ti,ab,kw OR 'senegal':ti,ab,kw OR 'solomon islands':ti,ab,kw OR 'sri lanka':ti,ab,kw OR 'tanzania':ti,ab,kw OR 'timor-leste':ti,ab,kw OR 'tunisia':ti,ab,kw OR 'ukraine':ti,ab,kw OR 'uzbekistan':ti,ab,kw OR 'vanuatu':ti,ab,kw OR 'vietnam':ti,ab,kw OR 'west bank':ti,ab,kw OR 'gaza':ti,ab,kw OR 'zambia':ti,ab,kw OR 'zimbabwe':ti,ab,kw OR 'low-income country':ti,ab,kw OR 'afghanistan':ti,ab,kw OR 'burkina faso':ti,ab,kw OR 'burundi':ti,ab,kw OR 'central african republic':ti,ab,kw OR 'chad':ti,ab,kw OR 'congo':ti,ab,kw OR 'eritrea':ti,ab,kw OR 'ethiopia':ti,ab,kw OR 'gambia':ti,ab,kw OR 'guinea':ti,ab,kw OR 'guinea-bissau':ti,ab,kw OR 'north korea':ti,ab,kw OR 'korea democratic people republic':ti,ab,kw OR 'haiti':ti,ab,kw OR 'liberia':ti,ab,kw OR 'madagascar':ti,ab,kw OR 'malawi':ti,ab,kw OR 'mali':ti,ab,kw OR 'mozambique':ti,ab,kw OR 'niger':ti,ab,kw OR 'rwanda':ti,ab,kw OR 'sierra leone':ti,ab,kw OR 'somalia':ti,ab,kw OR 'south sudan':ti,ab,kw OR 'sudan':ti,ab,kw OR 'syrian arab republic':ti,ab,kw OR 'syria':ti,ab,kw OR 'tajikistan':ti,ab,kw OR 'togo':ti,ab,kw OR 'yemen':ti,ab,kw
--	---

**Table 5. terms used in PsycINFO
abstract for all options**

<i>Search</i>	<i>Search terms</i>
<i>Population</i>	ab(("Adolescen*" OR "young" OR "youth" OR "teen" OR "teenag*" OR "child" OR "children" OR "paediatr*" OR "pediatr*" OR "juvenil*" OR "boy*" OR "girl*" OR "pubert*" OR "pubescen*"))
	AND
<i>Exposure</i>	ab(("swine flu" OR "H1N1" OR "flu pandemic" OR "Zika" OR "Ebola" OR "EVD" OR "EHF" OR "Ebola virus disease" OR "Ebola hemorrhagic fever" OR "COVID-19" OR "SARS-COV-2" OR "Novel Coronavirus" OR "2019-nCov"))
	AND
<i>Outcome (in SRHR)</i>	ab(("ASRH" or "SRH" or "SRHR" or "human right*" or "pregnan*" or "abort*" or "termination" or "antenatal" or "postnatal" or "perinatal" or "couple*" or "abuse" or "intercourse" or "rape" or "coerci*" or "violence" or "violation" or "incest" or "emergency pills" or "Long-acting reversible contraceptive" or "LARC" or "Misoprostol" or "mifepristone" or "methotrexate" or "meteneprost" or "prostaglandin" or "lilopristone" or "onapristone" or "oxytocin" or "sulprostone" or "family planning" or "sex education" or "STI" or "sexually transmitted infection" or "HPV vaccination" or "intimate partner violence" or "female genital mutilation" or "female genital cutting" or "FGM" or "maternal death" or "maternal morbidity" or "maternal mortality" or "newborn morbidity" or "newborn mortality" or "maternal care" or "newborn care" or "infant" or "infancy" or "stigma" or "taboo" or "Discrimination" or "GBV" or "gender"
	AND

<i>Low and middle-income countries (96)</i>	ab(("low income countr*" OR "middle income countr*" OR "low middle income countr*" OR "developing countr*" OR "middle income countr*" OR "LMIC" OR "upper middle income countr*" OR "Albania" OR "American Samoa" OR "Argentina" OR "Armenia" OR "Azerbaijan" OR "Belarus" OR "Belize" OR "bosnia" OR "Herzegovina" OR "Botswana" OR "Brazil" OR "Bulgaria" OR "China" OR "Colombia" OR "Costa Rica" OR "Cuba" OR "Dominica" OR "Dominican Republic" OR "Ecuador" OR "Equatorial Guinea" OR "Fiji" OR "Gabon" OR "Georgia" OR "Grenada" OR "Guatemala" OR "Guyana" OR "Indonesia" OR "Iran" OR "Iraq" OR "Jamaica" OR "Jordan" OR "Kazakhstan" OR "Kosovo" OR "Lebanon" OR "Libya" OR "Malaysia" OR "Maldives" OR "Marshall Islands" OR "Mexico" OR "Montenegro" OR "Namibia" OR "North Macedonia" OR "Paraguay" OR "Peru" OR "Russia" OR "Samoa" OR "Serbia" OR "South Africa" OR "Saint Lucia" OR "saint Vincent" OR "grenadines" OR "Suriname" OR "Thailand" OR "Tonga" OR "Turkey" OR "Turkmenistan" OR "Tuvalu" OR "Venezuela" OR "lower middle income countr*" OR "Algeria" OR "Angola" OR "Bangladesh" OR "Benin" OR "Bhutan" OR "Bolivia" OR "Cabo Verde" OR "Cambodia" OR "Cameroon" OR "Comoros" OR "Congo" OR "Cote d'Ivoire" OR "Ivory Coast" OR "Djibouti" OR "Egypt" OR "El Salvador" OR "Eswatini" OR "Ghana" OR "Honduras" OR "India" OR "Kenya" OR "Kiribati" OR "Kyrgyz" OR "Laos" OR "Lesotho" OR "Mauritania" OR "Micronesia" OR "Moldova" OR "Mongolia" OR "Morocco" OR "Myanmar" OR "Nepal" OR "Nicaragua" OR "Nigeria" OR "Pakistan" OR "Papua New Guinea" OR "Philippines" OR "Sao Tome" OR "Principe" OR "Senegal" OR "Solomon Islands" OR "Sri Lanka" OR "Tanzania" OR "Timor-Leste" OR "Tunisia" OR "Ukraine" OR "Uzbekistan" OR "Vanuatu" OR "Vietnam" OR "West Bank" OR "Gaza" OR "Zambia" OR "Zimbabwe" OR "low-income country" OR "Afghanistan" OR "Burkina Faso" OR "Burundi" OR "Central African Republic" OR "Chad" OR "Congo" OR "Eritrea" OR "Ethiopia" OR "Gambia" OR "Guinea" OR "Guinea-Bissau" OR "North Korea" OR "Korea Democratic People's Republic" OR "Haiti" OR "Liberia" OR "Madagascar"
---	---

OR "Malawi" OR "Mali" OR "Mozambique" OR
"Niger" OR "Rwanda" OR "Sierra Leone" OR
"Somalia" OR "South Sudan" OR "Sudan" OR "Syrian
Arab Republic" OR "Syria" OR "Tajikistan" OR
"Togo" OR "Yemen"))

Table 6. terms used in Cochrane library search
abstract, title and keywords for all options

Search	Search terms
Population	((“Adolescen*” OR “young” OR “youth” OR “teen” OR “teenag*” OR “child” OR “children” OR “paediatr*” OR “pediatr*” OR “juvenil*” OR “boy*” OR “girl*” OR “pubert*” OR “pubescen*”)):ti,ab,kw
AND	
Exposure	(“swine flu” or “H1N1” or “flu pandemic” or “Zika” or “Ebola” or “EVD” or “EHF” or “Ebola virus disease” or “Ebola hemorrhagic fever” or “COVID19” or “SARSCOV2” or “Novel Coronavirus” or “2019nCov”):ti,ab,kw
AND	
Outcome (in SRHR)	(“ASRH” or “SRH” or “SRHR” or “human right*” or “pregnan*” or “abort*” or “termination” or “antenatal” or “postnatal” or “perinatal” or “couple*” or “abuse” or “intercourse” or “rape” or “coerci*” or “violence” or “violation” or “incest” or “emergency pills” or “Long acting reversible contraceptive” or “LARC” or “Misoprostol” or “mifepristone” or “methotrexate” or “meteneprost” or “prostaglandin” or “lilopristone” or “onapristone” or “oxytocin” or “sulprostone” or “family planning” or “sex education” or “STI” or “sexually transmitted infection” or “HPV vaccination” or “intimate partner violence” or “female genital mutilation” or “female genital cutting” or “FGM” or “maternal death” or “maternal morbidity” or “maternal mortality” or “newborn morbidity” or “newborn mortality” or “maternal care” or “newborn care” or “infant” or “infancy” or “Discrimination” or “GBV” or “gender”) :ti,ab,kw
AND	
Low and middle-income countries (96)	(“low income countr*” or “middle income countr*” or “low middle income countr*” or “developing countr*”

or “middle income countr*” or “LMIC” or “upper middle income countr*” or “Albania” or “American Samoa” or “Argentina” or “Armenia” or “Azerbaijan” or “Belarus” or “Belize” or “bosnia” or “Herzegovina” or “Botswana” or “Brazil” or “Bulgaria” or “China” or “Colombia” or “Costa Rica” or “Cuba” or “Dominica” or “Dominican Republic” or “Ecuador” or “Equatorial Guinea” or “Fiji” or “Gabon” or “Georgia” or “Grenada” or “Guatemala” or “Guyana” or “Indonesia” or “Iran” or “Iraq” or “Jamaica” or “Jordan” or “Kazakhstan” or “Kosovo” or “Lebanon” or “Libya” or “Malaysia” or “Maldives” or “Marshall Islands” or “Mexico” or “Montenegro” or “Namibia” or “North Macedonia” or “Paraguay” or “Peru” or “Russia” or “Samoa” or “Serbia” or “South Africa” or “Saint Lucia” or “saint Vincent” or “grenadines” or “Suriname” or “Thailand” or “Tonga” or “Turkey” or “Turkmenistan” or “Tuvalu” or “Venezuela” or “lower middle income countr*” or “Algeria” or “Angola” or “Bangladesh” or “Benin” or “Bhutan” or “Bolivia” or “Cabo Verde” or “Cambodia” or “Cameroon” or “Comoros” or “Congo” or “Cote d’Ivoire” or “Ivory Coast” or “Djibouti” or “Egypt” or “El Salvador” or “Eswatini” or “Ghana” or “Honduras” or “India” or “Kenya” or “Kiribati” or “Kyrgyz” or “Laos” or “Lesotho” or “Mauritania” or “Micronesia” or “Moldova” or “Mongolia” or “Morocco” or “Myanmar” or “Nepal” or “Nicaragua” or “Nigeria” or “Pakistan” or “Papua New Guinea” or “Philippines” or “Sao Tome” or “Principe” or “Senegal” or “Solomon Islands” or “Sri Lanka” or “Tanzania” or “Timor Leste” or “Tunisia” or “Ukraine” or “Uzbekistan” or “Vanuatu” or “Vietnam” or “West Bank” or “Gaza” or “Zambia” or “Zimbabwe” or “low income country” or “Afghanistan” or “Burkina Faso” or “Burundi” or “Central African Republic” or “Chad” or “Congo” or “Eritrea” or “Ethiopia” or “Gambia” or “Guinea” or “Guinea Bissau” or “North Korea” or “Korea Democratic People’s Republic” or “Haiti” or “Liberia” or “Madagascar” or “Malawi” or “Mali” or “Mozambique” or “Niger” or “Rwanda” or “Sierra Leone” or “Somalia” or “South Sudan” or “Sudan” or “Syrian Arab Republic” or “Syria” or “Tajikistan” or “Togo” or “Yemen”):ti,ab,kw

BMJ Open

The impact of major disease outbreaks in the third millennium on adolescent and youth sexual and reproductive health and rights in Low and/or Middle Income Countries: a systematic scoping review protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051216.R1
Article Type:	Protocol
Date Submitted by the Author:	10-Feb-2022
Complete List of Authors:	<p>Akbarialiabad, Hossein; Shiraz University of Medical Sciences, Student research committee, Shiraz school of medicine, Shiraz University of Medical Sciences, Shiraz, Iran</p> <p>Shidhaye, Rahul ; Pravara Institute of Medical Sciences, Senior Research Scientist and Associate Professor of Psychiatry, Pravara Institute of Medical Sciences, Loni, India.</p> <p>Shidhaye , Pallavi; ICMR-National AIDS Research Institute, Scientist B, Division of Clinical Sciences, ICMR-National AIDS Research Institute Pune, India</p> <p>Cuijpers, Pim; VU University Amsterdam, Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research institute, and WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam</p> <p>Weaver, Marcia; University of Washington, Departments of Health Metrics Sciences and Global Health University of Washington</p> <p>Bahrololoom, Mina; Shiraz University of Medical Sciences</p> <p>Kiburi, Sarah; Nairobi Metropolitan Services, Kenya, Department of psychiatry, Mbagathi Hospital, Nairobi Metropolitan Service, Nairobi, Kenya</p> <p>Njuguna, Irene; Kenyatta National Hospital, Research & Programs</p> <p>Taghrir, Mohammad Hossein ; Shiraz University of Medical Sciences, Department of community medicine, Shiraz School of medicine, Shiraz University of medical sciences, Shiraz, Iran</p> <p>KUMAR, MANASI; University College London, Department of Psychology, University College London, London, United Kingdom ; Department of Psychiatry, University of Nairobi, Nairobi, Kenya</p>
Primary Subject Heading:	Sexual health
Secondary Subject Heading:	Reproductive medicine
Keywords:	Sexual and gender disorders < PSYCHIATRY, SEXUAL MEDICINE, REPRODUCTIVE MEDICINE, PUBLIC HEALTH, Child & adolescent psychiatry < PSYCHIATRY

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



The impact of major disease outbreaks in the third millennium on adolescent and youth sexual and reproductive health and rights in Low and/or Middle Income Countries: a systematic scoping review protocol

Hossein Akbarialiabad¹, Rahul Shidhaye², Pallavi Shidhaye³, Pim Cuijpers⁴, Marcia R Weaver⁵, Mina Bahrololoom¹, Sarah Kanana⁶, Irene Njuguna⁷, Mohammad Hossein Taghrir⁸, Manasi Kumar^{9,10*}

¹- Student research committee, Shiraz school of medicine, Shiraz University of Medical Sciences, Shiraz, Iran

²Research Scientist and Associate Professor of Psychiatry, Pravara Institute of Medical Sciences, Loni, India.

³Scientist B, Division of Clinical Sciences, ICMR-National AIDS Research Institute, Pune, India

⁴Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research Institute, and WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam, The Netherlands

⁵Research Professor, Departments of Health Metrics Sciences and Global Health, University of Washington, Seattle Washington, USA

⁶Department of psychiatry, Mbagathi Hospital, Nairobi Metropolitan Service, Nairobi, Kenya

⁷Kenyatta National Hospital, Research and Programs, Nairobi, Kenya

⁸Department of community medicine, Shiraz School of medicine, Shiraz University of medical sciences, Shiraz, Iran

⁹ Department of Psychiatry, University of Nairobi, Nairobi, Kenya

¹⁰Department of Psychology, University College London, London, United Kingdom

Corresponding author:

Dr. Manasi Kumar

Department of Psychiatry, University of Nairobi, Nairobi, Kenya

Department of Psychology, University College London, London, United Kingdom

Email: Email: m.kumar@ucl.ac.uk / Phone number: +254717379687/ Fax: +254-20-2723719

Word Count : 3596

Keyword: Sexual health, SARS-COV-2, Zika virus, Ebolavirus, H1N1 Virus

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Introduction

Sexual and Reproductive Health and Rights (SRHR) of young people continue to present a high burden and remain underinvested. This is more so in lower and middle-income countries (LMICs), where empirical evidence reveals disruption of SRHR maintenance, need for enhancement of programs, resources, and services during pandemics. Despite the importance of the subject, there is no published review yet combining recent disease outbreaks such as (H1N1/09, Zika, Ebola, and SARS-COV-2) to assess their impact on adolescents and youth SRHR in LMICs.

Methods and Analysis

We will adopt a four-step search to reach the maximum possible number of studies. In the first step, we had a limited/preliminary investigation in databases for getting relevant keywords (appendix1). Secondly, we will search in four databases: Pubmed, Cochrane Library, Embase, and PsycINFO. The search would begin from the inception of the first major outbreak in 2009 (H1N1/09) up to the date of publication of the protocol in early 2022. We will search databases using related keywords, screen title/abstract, and review full texts of the selected titles to arrive at the list of eligible studies. In the third stage, we will check their eligibility to the included article's reference list. In the fourth stage, we will check the citations of included papers in phase 2 to complete our study selection. We will include all types of original studies and without any language restriction in our final synthesis. Our review results will be charted for each pandemic separately and include the details related to authors, year, country, region of the study, study design, participants (disaggregated by age and gender), purpose, and report associated SRHR outcomes. The review will adhere to the PRISMA-ScR guideline.

Patient and public involvement

Patients or public were not involved in this study.

Ethics and dissemination

Ethical assessment is not required for this study. The results of the study will be presented in peer-reviewed publications and conferences on adolescent SRHR.

Strengths and limitations of our study

Strengths

- ✓ Our systematic scoping review protocol minimizes the possibility of duplications, and engages a stringent peer-review process to arrive at meaningful search outcomes around SRHR impacts.
- ✓ Our strategy will include four steps of searching and sifting through studies, maximizing our effort to reach all eligible studies.
- ✓ Using International Conference on Population and Development (ICPD) as our guiding framework to elicit key outcomes, we will to study adolescent SRHR rigorously and insightfully.

Limitations

- There could be other localized and globally impacting diseases outbreaks to consider other than what we focused on that merit further scrutiny.

86 Introduction

88 The beginning of sexual maturation differs based on sex, country, culture and religious context
89 (1). We know that adolescents are accessing services earlier, but the use of these services tends
90 to be inconsistent (2). The risk of neglecting the adolescent sexual and reproductive health
91 (ASRH) could impose a life-lasting impact on them. For females, adolescent pregnancy and
92 motherhood may be mentally and physically challenging (3, 4). It may also impede further
93 academic and educational achievements and economic potential (5). There is a high risk of
94 sexually transmitted infections (STIs), including HIV, reproductive coercion, and violence in
95 both sexes, with severe impacts seen especially for girls.(6). In 2016, it was found that 15-19
96 years old adolescents in developing countries had over 21 million pregnancies, and
97 approximately than half of them (12 million) resulted in child delivery (7). The number of
98 unintended pregnancies in girls between 15-19 years were around ten million, and about 55% of
99 these unintended pregnancies resulted in abortion (8, 9). Nearly 70% of the total abortion
100 attempts at the ages of 15-19 leads to unsafe abortions that can trigger catastrophic health effects
101 (9). In the Global Burden of Disease (GBD) 2015 report, the maternal mortality rate (MMR) was
102 highest in 10-14 years old girls. The ratio was as high as 278/100,000 (95% UI 229–339) (10).
103 Additionally, pregnancy and delivery-related complications cause more mortality in 15-19 years
104 old girls than other age groups (11).

106 Research question:

107 What is the existing evidence regarding the impact of the Zika virus, H1N1/09, Ebola and
108 SARS- COV-2outbreaks on adolescents and youth SRHR in LMICs?

110 **Value of the proposed review:** Adolescents and young people have SRHR needs that become
111 neglected and more complex during the disasters and outbreaks (12). Besides this, we have to
112 underscore that their life experience is limited, and their decision-making abilities are still
113 evolving. Consequently, their coping mechanisms and judgment capabilities to navigate complex
114 circumstances triggered by these extraordinary disease outbreaks are also limited. Despite this
115 developmental challenge, these crises offer a belated opportunity for Governments to fill the
116 gaps in sex education and provide access to contraceptives, safer motherhood, safer abortion, and
117 empowerment programs to address inequities in SRHR and protect vulnerable adolescents in
118 time. If governments do not take such opportunities, future outbreaks can further compound the
119 public health burden, including exacerbation of mental health issues, gender inequalities, and
120 social injustices in society. We know that these social determinants of adolescent health are tied
121 together, and this review will help understand the existing evidence and gaps. We have
122 prioritized studying four significant outbreaks since 2000: swine flu virus (H1N1/09), Ebola
123 virus, Zika virus, and SARS-COV-2 and their impacts on the SRHR of young and adolescents in
124 LMICs.

Influenza (H1N1/09) :

On the first July 2009, the World Health Organization acknowledged that the influenza outbreak had become pandemic and called it the Pandemic H1N1/09 virus, popularly known as “swine flu”. Pregnant women were considered a high-risk group for H1N1, as there was a higher rate of spontaneous abortion, preterm birth, low birth weight, fetal disease, as well as increased hospitalization and maternal death (13,14, 15). In a study at the outset of the pandemic in Australia and New Zealand, 9% of patients admitted to the Intensive Care Units (ICUs) were pregnant, a high proportion, given that 1% of the Australasian population were pregnant (16). A global pooled analysis revealed that pregnant women had a 3.5-25.3 fold higher risk of hospitalization compared to non-pregnant women of childbearing age(17). Due to the high mortality and morbidity rate in for pregnant women, vaccination of pregnant women was prioritized. However, studies show that poor vaccine uptake in pregnant women possibly a result of concerns regarding adverse effects (18, 19). Despite multiple studies showing that influenza vaccines are safe and have no significant threat to pregnancy, including spontaneous abortion, still birth and congenital malformations (20-22).

Zika Virus:

In March 2015, a group of patients were admitted in Brazil with rash, fatigue, and arthralgia, caused by the Zika Virus. In October that year, a few reports showed that the mother-newborn transmission of the virus (23) might lead to fetal loss, preterm delivery, and microcephalic (small head) babies, cumulatively known as congenital Zika syndrome (CZS) (24). Case reports showed high semen viral load, 10,000 times higher in semen than in blood in one case (25, 26). The transmission of the virus in humans is via sexual contact, blood transfusion, and organ transplantation (27-29). Combining the sexual transmission of the virus between partners and vertical transmission to the fetus, the WHO recommendations were adoption of safer sex practices, abstinence during pregnancy and emergency contraception (23). To date, a total of 86 countries have reported evidence of Zika infection (30).

In mid-January 2016, the health ministers of multiple Latin American countries recommended postponing pregnancy for at least 6-24 months. This seemed too idealistic and unrealistic, as more than half of the region's pregnancies have been unintended (31). In addition, inadequate sex education, difficulty accessing contraception, high rape prevalence, and local cultural constraints resulted in many women not adopting proper advice (32). Latin American countries have a relatively wide range of restrictive laws in SRHR. In El-Salvador, a self-induced abortion may lead to a penalty as high as 40 years of incarceration (33). On the other hand, in response to the Zika virus, the Columbian Ministry of Health declared that women have the right to have self-induced miscarriages. Nevertheless, due to poor education and scarcity of information, women are not well informed about the risk and their abortion rights(32). This was reflected by the fact that the rate of contraception usage did not change in Columbia during the outbreak in 2015-2016 (34).

Ebola virus:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Ebola hemorrhagic fever (EHF) is one of the fatal outbreaks within history. By the end of 2016, an estimated 11310 deaths resulted from 28618 definite cases, a staggering 40% mortality (35). The transmission route is through contact with infected patient’s blood or secretions and direct contact (mucus membrane and injured cutaneous tissue). Even after recovery, the virus may exist in the semen of cases for up to 7 weeks later (36). Burial ceremonies that require direct contact with the dead bodies contribute to Ebola transmission (37). The latter endangers females as they have significant roles in the ceremony. Pregnant women can also transmit the virus to their babies either in-utero or through breastmilk (38-40).

In the past century, multiple outbreaks of the Ebola virus occurred; the first one was in Zaire (currently known as the Democratic Republic of the Congo) and Sudan in 1976. The second outbreak occurred in Sudan around 1979. The third epidemic was in Gabon,1996. In this century, Northern Uganda faced the fourth epidemic in the fall of 2000. The 2014-2016 Ebola virus epidemic was the largest outbreak of the Ebola, considered a pandemic (41, 42). The outset was from Guinea and then moved to other countries such as Liberia and Sierra Leone. The current outbreak (2018- now) in the eastern Democratic Republic of Congo (DRC) is the last reported outbreak(43).

WHO Advisory Group on the Ebola Virus Disease Response recommended that the male who survived the Ebola virus disease (EVD) practice hygiene and safer sex until one year from symptom beginning or after two negative semen specimens for the Ebola virus. In some patients after recovery, Ebola Virus can persist in several tissues, such as testicles, the eye, and the central nervous system (44). In women who have contracted the disease during pregnancy, the virus remains in the placenta, amniotic fluid, and fetus. In those women who have been infected during breastfeeding, the virus could persist in breast milk (38, 45-50).

Ebola revealed many problems in adolescence SRHR in Africa (51). It is well-established that in humanitarian crises and disasters, women and girls are more vulnerable to gender-based violence (GBV) (52, 53). Multiple studies showed that in Sierra Leone, during the Ebola outbreak, the rate of adolescent pregnancy, rape, sexual and gender-based violation increased (51, 54, 55). United Nations Department of Political Affairs (UNDP) estimates that the teenage pregnancy rate increased by 65% during the outbreak (56). More girls were forced into prostitution/transactional sex due to family members' loss and financial insecurity(57). This makes the girls twice victims, as pregnant girls are forbidden to attend school in Libera and Sierra Leone (58). In Sierra-Leone, Ebola increased the fear of clinic attendance and impacted newborn and maternal care by disrupting the health services. This led to 549 maternal deaths, 2161 neonatal deaths and 883 still births in the year 2014-15 (59). The studies showed that even after pandemics in Liberia and Sierra-Leone, the level of contraceptive care and family planning had returned or exceeded the baseline level after 6 months (60) to 24 months (61). After six months of the outbreak in Guinea, the number of family planning sessions and prenatal care visits did not reverse to the prior level, harshly influencing the already insufficient care level (60).

SARS COV-2 :

In late 2019, multiple patients were diagnosed with a pneumonia-causing virus. Numerous subsequent studies showed a beta-coronidae family virus named “SARS-COV-2” by WHO to be the causative agent. In March 2020, the WHO declared SARS-COV-2 a pandemic and called the disease “coronavirus disease of 2019”, or simply “COVID-19”. Currently, there is no substantial evidence that the virus is transmitted sexually or through breastfeeding. The possibility of vertical transmission is controversial. However, a recently published systematic review and meta-analysis suggest that a minority of pregnant women can transmit the virus to newborns (62).

As the COVID-19 escalated, many countries adopted lockdown measures to mitigate the virus spread. The pandemic disrupted the supply chain of key contraceptive commodities (63). In practice, most Governments categorized sexual and reproductive health (SRH) services as non-essential and forced them to close. In late April 2020, United Nations Population Fund (UNFPA) predicted an estimated 7 million unintended pregnancies would occur (64), as a result of lack of access to SRH services like contraception and safe abortion care and time-sensitive potentially lifesaving services (65).

COVID-19 has forced nearly 1.4 billion children and youths out of school/university, including 743 million girls globally (66). This has raised concerns around long-term impact on their lives. As in addition to increasing poverty levels they might also experience increased sexual and gender-based violence, furthering the need not only for remediation and support services but also access to emergency contraception and other reproductive health services (67). COVID-19 has impacted health systems and services severely globally. It has also led to closure of educational institutions', public places for recreation and leisure, and has impacted movement (68). The protracted closures across the world have increased concerns around mental well-being and the availability/ accessibility of critical sexual and reproductive health services for adolescents in LMICs (69).

The Guttmacher institute predicted that with a hypothetical one-year 10% decline in the use of short and long-acting contraceptives in LMICs during the COVID-19 pandemic, 49 million women would lose their access to contraception of their choice. They estimated that around 15 million unintended pregnancies might occur, leading to an additional 1.7 million deliveries, 2.6 million newborns with significant complications, 168000 newborn deaths, and 28000 maternal deaths. The second hypothesis was that with a yearly 10% shift of safe abortion to unsafe abortion, an additional 3.3 unsafe abortions might occur in LMICs, and 1000 maternal deaths are expected (63). We know that the impact is possibly more than a 10% decline in such services, as frontline partners have predicted a slip could be up to 80% (70). In India, by March 2020, compared to December 2019, a 36% decrease in injectable contraceptives use and a 21% reduction in IUD insertion were reported. Simultaneously, the distribution of the condoms and oral contraceptive pills dipped 23% and 15%, respectively. The COVID-19 has interrupted the prevention programs and impacted the household economic status, and is expected to result to 13 million child marriages and two million female genital mutilations in the next decade(54). The health record analysis showed a 68% reduction in HPV vaccination from February to early April 2020(71).

We aim to conduct a scoping review to map the range, extent, and nature of effects of H1N1, Ebola virus, Zika virus, and SARS-COV-2 on SRHR among adolescents and youth in LMICs. Our goal is to identify current evidence in the literature, identify research gaps, and suggest future research. The details pertaining to population, exposure, and outcomes are provided in Table 1. Our data charting table is also available in appendix1 (Table A-1).

Objective

What is the existing evidence regarding the impact of theH1N1/09, Ebola, Zika virus and SARS-COV-2 outbreaks on adolescents and youth SRHR in LMICs?

Table1. PICOS format in this review

PICOS framework for Systematic Reviews	
Population	Adolescents and youths living in LMICs ages 10-24 years
Intervention/exposure	SARS-COV-2, Zika, H1N1, and Ebola-related disruptions
Comparative /Control intervention	n/a
Outcome	SRHR (sexual well-being, sexual health and illness outcomes, reproductive health and illnesses, sexual and reproductive health services outcomes)
Study designs	All types of original articles

Method

We will use *PRISMA*-ScR (Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews) Statement ,showed in appendix figure 1, to report the findings of our review (72).

Patient and public involvement

Patients or public were not involved in this study.

Eligibility Criteria Study designs:

We will include cross-sectional studies, case-control, cohort studies, clinical trials, and qualitative studies.

Participants/Population:

As per WHO definition of young people we are considering ages 10-24 to denote adolescents and youth.

Exposure/Intervention:

This study is designed to assess the impact of the H1N1/0, Ebola virus and Zika virus, and

Category of outcome	Sub-category
A) Contraception	- The proportion of adolescent and youth have access to contraceptive agent or procedures during the outbreaks
B) SRH Service Availability and Access	- The proportion of facilities providing SRHR services for adolescents and youth during the outbreaks - The proportion of health facilities providing post-abortion care and post-partum care for adolescent and youth and also services for those who have contracted HIV in our target group during the outbreaks
C) Knowledge About SRHR	- The proportion of adolescent and youth have basic knowledge on SRHR during the pandemics
D) Adolescent Fertility	- The adolescents and youth's birth rate during the outbreaks - The proportion of pregnancies related to those 24 years old and below during the

SARS-COV-2 on adolescents and youth SRHR in LMICs. The review will include work published after 2009

Control:

There is no control group in this review.

Outcome

We will focus on the SRHR of adolescents and youth in LMICs. SRHR would consist of sexual and reproductive health access and services outcomes, sexual health and well-being, and illness outcomes, reproductive health, reproductive health, rights and illnesses, sexual and reproductive health access and services outcomes.

Our proposed outcome indicators inspired by the International Conference on Population and Development (ICPD) Programme of Action and sustainable development goal (SDG) 2030 (73, 74) are summarized in table 2.

Table 2. The outcomes in our review

	outbreaks
E) Quality of Care, Including Respect for Rights	<ul style="list-style-type: none">- The proportion of females are knowledgeable about side effects of their contraceptive method of choice and how to deal with adverse effects and also are familiar with other methods of contraception during the outbreaks- The universal access to contraceptives and SRHR information during outbreaks in national policy actions during the outbreaks- An indicator reflective of respectful care and human rights in the provision of SRH information and services during the outbreaks
F) Prevention of Sexually Transmitted Infections	<ul style="list-style-type: none">- The rate of HPV vaccination during the epidemics
G) Abortions:	<ul style="list-style-type: none">- Number of unsafe abortions during outbreaks- Number of facilities provide safe abortions services; or if it's illegal, the number of facilities providing the services when unsafe abortions become complicated during the outbreaks
H) Comprehensive Sexuality Education	<ul style="list-style-type: none">- The proportion of school/facilities providing comprehensive sexual education during the outbreaks
I) Gender Equality in SRHR	<ul style="list-style-type: none">- whether adolescents and youth's sexual autonomy within marriage is respected during the outbreaks

Timing:

The first significant outbreak of the third millennium was the H1N1/09 (swine flu) in 2009; hence, all the papers published from 2009 until 2021 will be included.

Setting:

Our study will focus on LMICs. Low- and middle-income countries (LMICs) were defined according to the 2020-2021 World Bank classification(75).

Language:

Our review compiles articles with no language restriction. We will use Google Translate for the primary translation of abstracts of non-English articles. In case the abstract of the article fits the scope of this review, we will request the experts in SRHR familiar with that specific language, to evaluate the paper based on the inclusion/exclusion criteria and extract the data. Alternatively, we will request official language translation centers to make the English version of the article for our evaluation.

Exclusion criteria

- all non-original papers
- Those irrelevant to the SRHR
- Studies that do not include adolescents and youth (10-24 years old)
- Studies from countries not categorized as LMICs
- Those not relevant to Zika virus, Ebola Virus, H1N1/09, and SARS-COV-2 outbreaks.
- Full text is not available for evaluation
- If the population comprises of other ages besides 10-24, we will include it if data is disaggregated by age group or sub-group analysis on age 10-24. If they have included vivid results about this age group, we will consist of their study.
- Animal studies

Information Sources:

We will look at four databases in our review: Pubmed, Cochrane Library, Embase, and PsycINFO.

Search:

Table A-2 (appendix 1) shows our proposed search strategy in Pubmed. Tables A-3 to A-5 (appendix 1) is our search strategy for each of the other target databases (Cochrane Library, Embase, PsycINFO)

Selection of sources of evidence:

A four-step search strategy is being followed in this study. First, the authors have done a preliminary (and limited) search in several databases (Pubmed and Google Scholar) on this topic. We assessed the keywords, abstract, and several full texts to reach our study's final keywords. We discussed our search strategy multiple times until all authors reached a consensus reflected in Tables 3-8. Following the duplication removal, title/abstract, and full-text screening of studies using RAYYAN (76), we will choose our relevant evidence considering the inclusion/exclusion criteria.

In the third step, we will check the articles' reference lists to find more relevant studies. Finally, we will look for studies that have cited or included studies and check them for their eligibility. Any discrepancy would be addressed via discussion until consensus is reached in each stage.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Two authors worked separately in all searching and extraction stages, and discrepancies would be resolved through discussion.

Data Charting Process:

After choosing the eligible papers, two authors will separately extract the required items based on the extraction excel sheet form designed and approved by researchers for the process.

Data items:

Following the selected papers' primary selection and inclusion and extracting the results, the tabulation phase begins. We will categorize the extracted data based on the items mentioned before. We will extract the following data: authors, year, country and continent (region of the study), study design, participants (differentiated by age and gender), purpose, and main findings. Any disagreements in the process of data extraction will be resolved through discussion until consensus. We will also indicate the country's income level where studies are being done (low, lower-middle, and higher-middle income).

Critical appraisal of individual sources of evidence and synthesis of result

Based on the nature of scoping review, we will neither synthesize the findings/results, nor critically appraise the papers, which is more applicable by performing a systematic review(77). However, publications and the mentioned data would be summarized and categorized.

Ethics and dissemination

As the nature of the work is a scoping review of the currently published papers in peer review journals, ethical approval is not required for this study. The results of the study will be presented in a peer-reviewed publication.

Authors' contribution: HA and MK developed the concept and wrote the early draft. RS, PS ,MHT, PC, MRW, SK, IN and MB edited the early draft. All authors approved the final version.

Funding: The review received no specific grant from any funding agency in public, commercial or not-for-profit sectors.

Competing interest: none

References

1. Chandra-Mouli V, McCarraher DR, Phillips SJ, Williamson NE, Hainsworth G. Contraception for adolescents in low and middle income countries: needs, barriers, and access. *Reproductive health*. 2014;11(1):1.
2. Blanc AK, Tsui AO, Croft TN, Trevitt LJ. Patterns and trends in adolescents' contraceptive use and discontinuation in developing countries and comparisons with adult women. 2009:63-71.
3. WHO. Adolescent pregnancy. Available at <https://www.who.int/en/news-room/fact-sheets/detail/adolescent-pregnancy> . Accessed 27th Dec 2020. .
4. Hodgkinson SC, Colantuoni E, Roberts D, Berg-Cross L, Belcher HM. Depressive symptoms and birth outcomes among pregnant teenagers. 2010;23(1):16-22.
5. Raj A, Boehmer U. Girl child marriage and its association with national rates of HIV, maternal health, and infant mortality across 97 countries. 2013;19(4):536-51.
6. Organization WH. Global standards for quality health-care services for adolescents: a guide to implement a standards-driven approach to improve the quality of health care services for adolescents. 2015.
7. Darroch JE, Woog V, Bankole A, Ashford LS. Adding it up: costs and benefits of meeting the contraceptive needs of adolescents. 2016.
8. Sully E. Adding it up: investing in sexual and reproductive health 2019. New York: Guttmacher Institute; 2020. 2019.
9. Darroch JE, Woog V, Bankole A, Ashford LS, Points KJ. Costs and benefits of meeting the contraceptive needs of adolescents. 2016.
10. Gore FM, Bloem PJ, Patton GC, Ferguson J, Joseph V, Coffey C, et al. Global burden of disease in young people aged 10–24 years: a systematic analysis. 2011;377(9783):2093-102.
11. Neal S, Matthews Z, Frost M, Fogstad H, Camacho AV, Laski L. Childbearing in adolescents aged 12–15 years in low resource countries: a neglected issue. New estimates from demographic and household surveys in 42 countries. 2012;91(9):1114-8.
12. Guessoum SB, Lachal J, Radjack R, Carretier E, Minassian S, Benoit L, et al. Adolescent psychiatric disorders during the COVID-19 pandemic and lockdown. 2020:113264.
13. Briand S, editor Epidemiology and illness severity of pandemic (H1N1) 09 virus. World Health Organization Strategic Advisory Group of Experts on Immunization Extraordinary Meeting, Geneva, Switzerland; 2009.
14. WHO Guidelines Approved by the Guidelines Review Committee. Pregnancy and Pandemic Influenza A (H1N1) 2009: Information for Programme Managers and Clinicians. Geneva: World Health Organization
- Copyright © World Health Organization 2010.; 2010.
15. Somerville LK, Basile K, Dwyer DE, Kok JF. The impact of influenza virus infection in pregnancy. 2018;13(2):263-74.
16. Medicine A. Critical care services and 2009 H1N1 influenza in Australia and New Zealand. 2009;361(20):1925-34.
17. Van Kerkhove MD, Vandemaële KA, Shinde V, Jaramillo-Gutierrez G, Koukounari A, Donnelly CA, et al. Risk factors for severe outcomes following 2009 influenza A (H1N1) infection: a global pooled analysis. 2011;8(7):e1001053.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

18. Drees M, Johnson O, Wong E, Stewart A, Ferisin S, Silverman PR, et al. Acceptance of 2009 H1N1 influenza vaccine among pregnant women in Delaware. 2012;29(04):289-94.

19. McCarthy EA, Pollock WE, Nolan T, Hay S, McDonald SJA, Obstetrics NZJo, et al. Improving influenza vaccination coverage in pregnancy in Melbourne 2010–2011. 2012;52(4):334-41.

20. McMillan M, Porritt K, Kralik D, Costi L, Marshall H. Influenza vaccination during pregnancy: a systematic review of fetal death, spontaneous abortion, and congenital malformation safety outcomes. Vaccine. 2015;33(18):2108-17.

21. Goldfarb I, Panda B, Wylie B, Riley LJAjoo, gynecology. Uptake of influenza vaccine in pregnant women during the 2009 H1N1 influenza pandemic. 2011;204(6):S112-S5.

22. Conlin AMS, Bukowinski AT, Sevic CJ, DeScisciolo C, Crum-Cianflone NFJO, Gynecology. Safety of the pandemic H1N1 influenza vaccine among pregnant US military women and their newborns. 2013;121(3):511-8.

23. WHO. Zika Virus. Available from <https://www.who.int/news-room/fact-sheets/detail/zika-virus> . Accessed 28th Dec 2020. .

24. Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Da Fonseca EB, et al. Characterizing the pattern of anomalies in congenital Zika syndrome for pediatric clinicians. 2017;171(3):288-95.

25. Mansuy JM, Dutertre M, Mengelle C, Fourcade C, Marchou B, Delobel P, et al. Zika virus: high infectious viral load in semen, a new sexually transmitted pathogen? 2016;16(4):405.

26. Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau V-MJEid. Potential sexual transmission of Zika virus. 2015;21(2):359.

27. Brent CJMM, report mw. Preliminary findings from an investigation of Zika virus infection in a patient with no known risk factors—Utah, 2016. 2016;65.

28. Moghadam SRJ, Bayrami S, Moghadam SJ, Golrokhi R, Pahlaviani FG, SeyedAlinaghi SJAPJoTB. Zika virus: A review of literature. 2016;6(12):989-94.

29. Katz LM, Rossmann SNJAop, medicine I. Zika and the blood supply: a work in progress. 2017;141(1):85-92.

30. Chapter 6 - Comparing the Zika Virus Disease Pandemic to Other Disease Pandemics. In: Qureshi AI, editor. Zika Virus Disease: Academic Press; 2018. p. 85-96.

31. Sedgh G, Singh S, Hussain RJSifp. Intended and unintended pregnancies worldwide in 2012 and recent trends. 2014;45(3):301-14.

32. Roa MJTL. Zika virus outbreak: reproductive health and rights in Latin America. 2016;387(10021):843.

33. Viterna J, Bautista JSGJH, Rights H. Pregnancy and the 40-year prison sentence: how “abortion is murder” became institutionalized in the Salvadoran judicial system. 2017;19(1):81.

34. Machado-Alba JE, Machado-Duque ME, Gaviria-Mendoza A, Orozco-Giraldo VAJTL. Hormonal contraceptive prescriptions in Colombia and Zika virus. 2016;387(10032):1993.

35. Coltart CE, Lindsey B, Ghinai I, Johnson AM, Heymann DLJPTotRSBBS. The Ebola outbreak, 2013–2016: old lessons for new epidemics. 2017;372(1721):20160297.

36. Ebola Haemorrhagic Fever. Retrieved from <https://www.msf.org/who-fact-sheet-ebola-haemorrhagic-fever> . Accessed in 29th Dec 2020.

37. Nkangu MN, Olatunde OA, Yaya SIIdop. The perspective of gender on the Ebola virus using a risk management and population health framework: a scoping review. 2017;6(1):135.

38. Medina-Rivera M, Centeno-Tablante E, Finkelstein JL, Rayco-Solon P, Peña-Rosas JP, Garcia-Casal MN, et al. Presence of Ebola virus in breast milk and risk of mother-to-child transmission: synthesis of evidence. 2020.
39. Nordenstedt H, Bah EI, de la Vega M-A, Barry M, N'Faly M, Barry M, et al. Ebola virus in breast milk in an Ebola virus-positive mother with twin babies, Guinea, 2015. 2016;22(4):759.
40. Sissoko D, Keita M, Diallo B, Aliabadi N, Fitter DL, Dahl BA, et al. Ebola virus persistence in breast milk after no reported illness: a likely source of virus transmission from mother to child. 2017;64(4):513-6.
41. Ifediora OF, Aning KJGGARoM, Organizations I. West Africa's Ebola Pandemic: Toward Effective Multilateral Responses to Health Crises. 2017;23(2):225-44.
42. Zhong Y, Xu J, Li T, Yu X, Sheng MJSCLS. Potential clinical treatment for Ebola pandemic. 2014;57(10):982.
43. WHO. Ebola virus . Retrived from <https://www.who.int/news-room/fact-sheets/detail/ebola-virus-disease>. Accessed 29th DEC 2020. .
44. Racine T, Kobinger GP. Viral pathogenesis: Unlocking Ebola persistence. Nature Microbiology. 2017;2(9):17124.
45. Chughtai A, Barnes M, Macintyre CJE, Infection. Persistence of Ebola virus in various body fluids during convalescence: evidence and implications for disease transmission and control. 2016;144(8):1652-60.
46. Bausch DG, Towner JS, Dowell SF, Kaducu F, Lukwiya M, Sanchez A, et al. Assessment of the risk of Ebola virus transmission from bodily fluids and fomites. 2007;196(Supplement_2):S142-S7.
47. Baggi F, Taybi A, Kurth A, Van Herp M, Di Caro A, Wölfel R, et al. Management of pregnant women infected with Ebola virus in a treatment centre in Guinea, June 2014. 2014;19(49):20983.
48. Caluwaerts S, Fautsch T, Lagrou D, Moreau M, Modet Camara A, Günther S, et al. Dilemmas in managing pregnant women with Ebola: 2 case reports. 2016;62(7):903-5.
49. Oduyebo T, Pineda D, Lamin M, Leung A, Corbett C, Jamieson DJJO, et al. A pregnant patient with Ebola virus disease. 2015;126(6):1273-5.
50. Muehlenbachs A, de la Rosa Vázquez O, Bausch DG, Schafer IJ, Paddock CD, Nyakio JP, et al. Ebola virus disease in pregnancy: clinical, histopathologic, and immunohistochemical findings. 2017;215(1):64-9.
51. Denney L, Gordon R, Ibrahim AJLODI. Teenage Pregnancy after Ebola in Sierra Leone.'. 2015.
52. Hynes M, Ward J, Robertson K, Crouse CJD. A determination of the prevalence of gender-based violence among conflict-affected populations in East Timor. 2004;28(3):294-321.
53. Kofman YB, Garfin DRJPTT, Research, Practice,, Policy. Home is not always a haven: The domestic violence crisis amid the COVID-19 pandemic. 2020.
54. PAHO .Adolescent Pregnancy in Latin America and the Caribbean. Retrived from https://pancap.org/pc/pcc/media/pancap_event/Technical-brief-Adolescent-Pregnancy-in-Latin-America-and-the-Caribbean.pdf . Accessed 30th Dec 2020. .
55. Elston J, Moosa A, Moses F, Walker G, Dotta N, Waldman RJ, et al. Impact of the Ebola outbreak on health systems and population health in Sierra Leone. 2016;38(4):673-8.
56. Werber CJQA. How Ebola led to more teenage pregnancy in West Africa. 2015.

1
2
3 490 57. Minor OMJAiA. Ebola and accusation: Gender dimensions of stigma in Sierra Leone's
4 491 Ebola response. 2017;24(2):25-35.
5 492 58. Walsh S, Mulhern E. A Step in the Rights' Direction: Advocacy, Negotiation, and Money
6 493 as Tools for Realising the Right to Education for Pregnant Girls in Sierra Leone During the
7 494 Ebola Epidemic. *Pregnant in the Time of Ebola*: Springer; 2019. p. 399-416.
8 495 59. Sochas L, Channon AA, Nam SJHp, planning. Counting indirect crisis-related deaths in
9 496 the context of a low-resilience health system: the case of maternal and neonatal health during the
10 497 Ebola epidemic in Sierra Leone. 2017;32(suppl_3):iii32-iii9.
11 498 60. Camara BS, Delamou A, Diro E, Béavogui AH, El Ayadi AM, Sidibé S, et al. Effect of
12 499 the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: an
13 500 ecological study. 2017;111(1):22-9.
14 501 61. Bietsch K, Williamson J, Reeves MJSiFP. Family planning during and after the West
15 502 African Ebola crisis. 2020;51(1):71-86.
16 503 62. Yang Z, Liu YJAjop. Vertical transmission of severe acute respiratory syndrome
17 504 coronavirus 2: a systematic review. 2020;37(10):1055.
18 505 63. Riley T, Sully E, Ahmed Z, Biddlecom AJIPSRH. Estimates of the potential impact of
19 506 the COVID-19 pandemic on sexual and reproductive health in low-and middle-income countries.
20 507 2020;46:46.
21 508 64. Impact of the COVID-19 Pandemic on Family Planning and Ending Gender-based
22 509 Violence, Female Genital Mutilation and Child Marriage . UNFPA. Retrieved from
23 510 [https://www.unfpa.org/resources/impact-covid-19-pandemic-family-planning-and-ending-](https://www.unfpa.org/resources/impact-covid-19-pandemic-family-planning-and-ending-gender-based-violence-female-genital)
24 511 [gender-based-violence-female-genital](https://www.unfpa.org/resources/impact-covid-19-pandemic-family-planning-and-ending-gender-based-violence-female-genital). Accessed 30 Dec 2020 . .
25 512 65. Cousins SJTL. COVID-19 has “devastating” effect on women and girls.
26 513 2020;396(10247):301-2.
27 514 66. COVID-19 School Closures Around the World Will Hit Girls Hardest . available from
28 515 <https://plan-international.org/blog/2020/03/covid-19-school-closures-hit-girls-hardest> . 30th DEC
29 516 2020.
30 517 67. Nanda P, Tandon S, Khanna A. Virtual and essential – adolescent SRHR in the time of
31 518 COVID-19. *Sexual and Reproductive Health Matters*. 2020;28(1):1831136.
32 519 68. Lee JJTLC, Health A. Mental health effects of school closures during COVID-19.
33 520 2020;4(6):421.
34 521 69. Mayurasakorn K, Pinsawas B, Mongkolsucharitkul P, Sranacharoenpong K, Damapong
35 522 SnJJoP, Health C. School closure, COVID-19 and lunch programme: Unprecedented
36 523 undernutrition crisis in low-middle income countries. 2020;56(7):1013-7.
37 524 70. Marie Stopes International, Methodology for calculating impact of COVID-19, 2020,
38 525 <https://www.mariestopes.org/resources/methodology-for-calculating-impact-of-covid-19>.
39 526 Accessed 30 Dec 2020.
40 527 71. Vaccine Rates Drop Dangerously as Parents Avoid Doctor's Visits .
41 528 <https://www.nytimes.com/2020/04/23/health/coronavirus-measles-vaccines.html> . Accessed 30
42 529 DEC 2020.
43 530 72. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA
44 531 extension for scoping reviews (PRISMA-ScR): checklist and explanation. 2018;169(7):467-73.
45 532 73. International Conference on Population and Development Programme of Action . 2014 .
46 533 available in : [https://www.unfpa.org/publications/international-conference-population-and-](https://www.unfpa.org/publications/international-conference-population-and-development-programme-action)
47 534 [development-programme-action](https://www.unfpa.org/publications/international-conference-population-and-development-programme-action) . accessed date March 9th 2021. .

74. Barot S, Cohen S, Darroch J, Galati A, Polis C, Singh S, et al. Sexual and Reproductive Health and Rights Indicators for the SDGs. New York: Guttmacher Institute; 2015.
75. World Bank. Low and middle income countries. Retrived from <https://data.worldbank.org/country/XO> . Accessed 28th Dec 2020.
76. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Systematic reviews*. 2016;5(1):1-10.
77. Askarian M, Khakpour M, Taghrir MH, Akbarialiabad H, Borazjani RJJes. Investigating the epidemiology of the methanol poisoning outbreaks in the third millennium: a scoping review protocol. 2020.

Figure 1 PRISMA flow chart



PRISMA 2009 Flow Diagram for scoping reviews

Identification

Screening

Eligibility

Included




Diagram1. Flow chart of the review phases (98)

Table A-1. data charting table

Authors/year	Country and income level	Continent (geographical region)	Sample size and Median Participant age (10-14), (15-19), (20-24), and the gender	Study design	Purpose	Main Findings

TableA-2. Search in PUBMED/MEDLINE (*Title/Abstract for all options*)

<i>PUBMED</i>	<i>Search terms in Pubmed</i>
<i>Population</i>	("adolescen*" [Title/Abstract] OR "young" [Title/Abstract] OR "youth" [Title/Abstract] OR "teen" [Title/Abstract] OR "teenager" [Title/Abstract] OR "child" [Title/Abstract] OR "children" [Title/Abstract] OR "paediatr*" [Title/Abstract] OR "pediatr*" [Title/Abstract] OR "juvenil*" [Title/Abstract] OR "boy" [Title/Abstract] OR "boys" [Title/Abstract] OR "girl" [Title/Abstract] OR "girls" [Title/Abstract] OR "pubert*" [Title/Abstract] OR "pubescen*" [Title/Abstract] OR "Young Adult" [Title/Abstract] OR "Young Adults" [Title/Abstract] OR "Adolescent" [MeSH Terms] OR "Young Adult" [MeSH Terms])
	AND
<i>Exposure</i>	((("swine flu" [Title/Abstract] OR "H1N1" [Title/Abstract] OR "flu pandemic" [Title/Abstract] OR "Zika" [Title/Abstract] OR "Ebola" [Title/Abstract] OR "EVD" [Title/Abstract] OR "EHF" [Title/Abstract] OR "Ebola virus disease" [Title/Abstract] OR "Ebola hemorrhagic fever" [Title/Abstract] OR "COVID-19" [Title/Abstract] OR "SARS-COV-2" [Title/Abstract] OR "Novel Coronavirus" [Title/Abstract] OR "2019-nCov" [Title/Abstract]) OR ("SARS-CoV-2" [MeSH Terms])) OR ("hemorrhagic fever, ebola" [MeSH Terms])) OR (ebolavirus [MeSH Terms])) OR ("influenza a virus, h1n1 subtype" [MeSH Terms])) OR ("Zika virus" [MeSH Terms])
	AND
<i>Outcome (in SRHR)</i>	((("ASRH" [Title/Abstract] OR "SRH" [Title/Abstract] OR "SRHR" [Title/Abstract] OR "human right*" [Title/Abstract] OR "pregnan*" [Title/Abstract] OR "abort*" [Title/Abstract] OR "termination" [Title/Abstract] OR "antenatal" [Title/Abstract] OR

	<p> <i>"postnatal"[Title/Abstract] OR</i> <i>"perinatal"[Title/Abstract] OR</i> <i>"couple*"[Title/Abstract] OR "abuse"[Title/Abstract]</i> <i>OR "intercourse"[Title/Abstract] OR</i> <i>"rape"[Title/Abstract] OR "coerci*"[Title/Abstract]</i> <i>OR "violence"[Title/Abstract] OR</i> <i>"violation"[Title/Abstract] OR "incest"[Title/Abstract]</i> <i>OR "emergency pills"[Title/Abstract] OR "Long-acting</i> <i>reversible contraceptive"[Title/Abstract] OR</i> <i>"LARC"[Title/Abstract] OR</i> <i>"Misoprostol"[Title/Abstract] OR</i> <i>"mifepristone"[Title/Abstract] OR</i> <i>"methotrexate"[Title/Abstract] OR</i> <i>"meteneprost"[Title/Abstract] OR</i> <i>"prostaglandin"[Title/Abstract] OR</i> <i>"lilopristone"[Title/Abstract] OR</i> <i>"onapristone"[Title/Abstract] OR "oxytocin"</i> <i>[Title/Abstract] OR "sulprostone"[Title/Abstract] OR</i> <i>"family planning"[Title/Abstract] OR "sex</i> <i>education"[Title/Abstract] OR "STI"[Title/Abstract]</i> <i>OR "sexually transmitted infection"[Title/Abstract] OR</i> <i>"HPV vaccination"[Title/Abstract] OR "intimate</i> <i>partner violence"[Title/Abstract] OR "female genital</i> <i>mutilation"[Title/Abstract] OR "female genital</i> <i>cutting"[Title/Abstract] OR "FGM"[Title/Abstract] OR</i> <i>"maternal death"[Title/Abstract] OR "maternal</i> <i>morbidity"[Title/Abstract] OR "maternal</i> <i>mortality"[Title/Abstract] OR "newborn</i> <i>morbidity"[Title/Abstract] OR "newborn</i> <i>mortality"[Title/Abstract] OR "maternal</i> <i>care"[Title/Abstract] OR "newborn</i> <i>care"[Title/Abstract] OR "infant"[Title/Abstract] OR</i> <i>"infancy"[Title/Abstract] OR "stigma"[Title/Abstract]</i> <i>OR "taboo"[Title/Abstract] OR</i> <i>"Discrimination"[Title/Abstract] OR</i> <i>"GBV"[Title/Abstract] OR "gender"[Title/Abstract]</i> <i>OR "VAWG"[Title/Abstract] OR (sexual and gender</i> <i>disorders[MeSH Terms])) OR (sexual and gender</i> <i>minorities[MeSH Terms])) OR ("Adolescent sexual</i> <i>and reproductive health" OR "sexual and reproductive</i> <i>health" OR "Menstruation-Inducing Agents")</i> </p>
	<p>AND</p>

Low and middle-income countries (96)	((("low income countr*" [Title/Abstract] OR "middle income countr*" [Title/Abstract] OR "low middle income countr*" [Title/Abstract] OR "developing countr*" [Title/Abstract] OR "middle income countr*" [Title/Abstract] OR "LMIC" [Title/Abstract] OR "upper middle income countr*" [Title/Abstract] OR "Albania" [Title/Abstract] OR "American Samoa" [Title/Abstract] OR "Argentina" [Title/Abstract] OR "Armenia" [Title/Abstract] OR "Azerbaijan" [Title/Abstract] OR "Belarus" [Title/Abstract] OR "Belize" [Title/Abstract] OR "bosnia" [Title/Abstract] OR "Herzegovina" [Title/Abstract] OR "Botswana" [Title/Abstract] OR "Brazil" [Title/Abstract] OR "Bulgaria" [Title/Abstract] OR "China" [Title/Abstract] OR "Colombia" [Title/Abstract] OR "Costa Rica" [Title/Abstract] OR "Cuba" [Title/Abstract] OR "Dominica" [Title/Abstract] OR "Dominican Republic" [Title/Abstract] OR "Ecuador" [Title/Abstract] OR "Equatorial Guinea" [Title/Abstract] OR "Fiji" [Title/Abstract] OR "Gabon" [Title/Abstract] OR "Georgia" [Title/Abstract] OR "Grenada" [Title/Abstract] OR "Guatemala" [Title/Abstract] OR "Guyana" [Title/Abstract] OR "Indonesia" [Title/Abstract] OR "Iran" [Title/Abstract] OR "Iraq" [Title/Abstract] OR "Jamaica" [Title/Abstract] OR "Jordan" [Title/Abstract] OR "Kazakhstan" [Title/Abstract] OR "Kosovo" [Title/Abstract] OR "Lebanon" [Title/Abstract] OR "Libya" [Title/Abstract] OR "Malaysia" [Title/Abstract] OR "Maldives" [Title/Abstract] OR "Marshall Islands" [Title/Abstract] OR "Mexico" [Title/Abstract] OR "Montenegro" [Title/Abstract] OR "Namibia" [Title/Abstract] OR "North Macedonia" [Title/Abstract] OR "Paraguay" [Title/Abstract] OR "Peru" [Title/Abstract] OR "Russia" [Title/Abstract] OR "Samoa" [Title/Abstract] OR "Serbia" [Title/Abstract] OR "South Africa" [Title/Abstract] OR "Saint Lucia" [Title/Abstract] OR "saint Vincent" [Title/Abstract] OR "grenadines" [Title/Abstract] OR
--------------------------------------	---

"Suriname"[Title/Abstract] OR
 "Thailand"[Title/Abstract] OR
 "Tonga"[Title/Abstract] OR "Turkey"[Title/Abstract]
 OR "Turkmenistan"[Title/Abstract] OR
 "Tuvalu"[Title/Abstract] OR
 "Venezuela"[Title/Abstract] OR "lower middle income
 countr*"[Title/Abstract] OR "Algeria"[Title/Abstract]
 OR "Angola"[Title/Abstract] OR
 "Bangladesh"[Title/Abstract] OR
 "Benin"[Title/Abstract] OR "Bhutan"[Title/Abstract]
 OR "Bolivia"[Title/Abstract] OR "Cabo
 Verde"[Title/Abstract] OR "Cambodia"[Title/Abstract]
 OR "Cameroon"[Title/Abstract] OR
 "Comoros"[Title/Abstract] OR
 "Congo"[Title/Abstract] OR "Cote
 d'Ivoire"[Title/Abstract] OR "Djibouti"[Title/Abstract]
 OR "Egypt"[Title/Abstract] OR "El
 Salvador"[Title/Abstract] OR
 "Eswatini"[Title/Abstract] OR
 "Ghana"[Title/Abstract] OR
 "Honduras"[Title/Abstract] OR "India"[Title/Abstract]
 OR "Kenya"[Title/Abstract] OR
 "Kiribati"[Title/Abstract] OR "Kyrgyz"[Title/Abstract]
 OR "Laos"[Title/Abstract] OR
 "Lesotho"[Title/Abstract] OR
 "Mauritania"[Title/Abstract] OR
 "Micronesia"[Title/Abstract] OR
 "Moldova"[Title/Abstract] OR
 "Mongolia"[Title/Abstract] OR
 "Morocco"[Title/Abstract] OR
 "Myanmar"[Title/Abstract] OR
 "Nepal"[Title/Abstract] OR
 "Nicaragua"[Title/Abstract] OR
 "Nigeria"[Title/Abstract] OR
 "Pakistan"[Title/Abstract] OR "Papua New
 Guinea"[Title/Abstract] OR
 "Philippines"[Title/Abstract] OR "Sao
 Tome"[Title/Abstract] OR "Principe"[Title/Abstract]
 OR "Senegal"[Title/Abstract] OR "Solomon
 Islands"[Title/Abstract] OR "Sri
 Lanka"[Title/Abstract] OR "Tanzania"[Title/Abstract]
 OR "Timor-Leste"[Title/Abstract] OR
 "Tunisia"[Title/Abstract] OR
 "Ukraine"[Title/Abstract] OR
 "Uzbekistan"[Title/Abstract] OR
 "Vanuatu"[Title/Abstract] OR
 "Vietnam"[Title/Abstract] OR "West

Bank"[Title/Abstract] OR "Gaza"[Title/Abstract] OR "Zambia"[Title/Abstract] OR "Zimbabwe"[Title/Abstract] OR "low-income country"[Title/Abstract] OR "Afghanistan"[Title/Abstract] OR "Burkina Faso"[Title/Abstract] OR "Burundi"[Title/Abstract] OR "Central African Republic"[Title/Abstract] OR "Chad"[Title/Abstract] OR "Congo"[Title/Abstract] OR "Eritrea"[Title/Abstract] OR "Ethiopia"[Title/Abstract] OR "Gambia"[Title/Abstract] OR "Guinea"[Title/Abstract] OR "Guinea-Bissau"[Title/Abstract] OR "North Korea"[Title/Abstract] OR "Korea Democratic People's Republic"[Title/Abstract] OR "Haiti"[Title/Abstract] OR "Liberia"[Title/Abstract] OR "Madagascar"[Title/Abstract] OR "Malawi"[Title/Abstract] OR "Mali"[Title/Abstract] OR "Mozambique"[Title/Abstract] OR "Niger"[Title/Abstract] OR "Rwanda"[Title/Abstract] OR "Sierra Leone"[Title/Abstract] OR "Somalia"[Title/Abstract] OR "South Sudan"[Title/Abstract] OR "Sudan"[Title/Abstract] OR "Syrian Arab Republic"[Title/Abstract] OR "Syria"[Title/Abstract] OR "Tajikistan"[Title/Abstract] OR "Togo"[Title/Abstract] OR "Yemen"[Title/Abstract]) OR ("africa"[MeSH Terms])) OR ("central america"[MeSH Terms])) OR ("middle east"[MeSH Terms]))

Table A-3. terms used in Embase

Abstract, title, and keywords for all option

Search	Search terms
Population	'adolescen*':ti,ab,kw OR 'young':ti,ab,kw OR 'youth':ti,ab,kw OR 'teen':ti,ab,kw OR 'teens':ti,ab,kw OR 'teenag*':ti,ab,kw OR 'child*':ti,ab,kw OR 'paediatr*':ti,ab,kw OR 'pediatr*':ti,ab,kw OR 'juvenil*':ti,ab,kw OR 'boy':ti,ab,kw OR 'boys':ti,ab,kw OR 'girl':ti,ab,kw OR 'girls':ti,ab,kw OR 'pubert*':ti,ab,kw OR 'pubescen*':ti,ab,kw
	AND
Exposure	'swine flu':ti,ab,kw OR 'h1n1':ti,ab,kw OR 'flu pandemic':ti,ab,kw OR 'zika':ti,ab,kw OR 'ebola':ti,ab,kw OR 'evd':ti,ab,kw OR 'ehf':ti,ab,kw OR 'ebola virus disease':ti,ab,kw OR 'ebola hemorrhagic fever':ti,ab,kw OR 'covid-19':ti,ab,kw OR 'sars-cov-2':ti,ab,kw OR 'novel coronavirus':ti,ab,kw OR '2019-ncov':ti,ab,kw
	AND
Outcome (in SRHR)	asrh:ti,ab,kw OR 'srh':ti,ab,kw OR 'srhr':ti,ab,kw OR 'human right*':ti,ab,kw OR 'pregnan*':ti,ab,kw OR 'abort*':ti,ab,kw OR 'termination':ti,ab,kw OR 'antenatal':ti,ab,kw OR 'postnatal':ti,ab,kw OR 'perinatal':ti,ab,kw OR 'couple*':ti,ab,kw OR 'abuse':ti,ab,kw OR 'intercourse':ti,ab,kw OR 'rape':ti,ab,kw OR 'coerci*':ti,ab,kw OR 'violence':ti,ab,kw OR 'violation':ti,ab,kw OR 'incest':ti,ab,kw OR 'emergency pills':ti,ab,kw OR 'long-acting reversible contraceptive':ti,ab,kw OR 'larc':ti,ab,kw OR 'misoprostol':ti,ab,kw OR 'mifepristone':ti,ab,kw OR 'methotrexate':ti,ab,kw OR 'meteneprost':ti,ab,kw OR 'prostaglandin':ti,ab,kw OR 'lilopristone':ti,ab,kw OR 'onapristone':ti,ab,kw OR oxytocin:ti,ab,kw OR 'sulprostone':ti,ab,kw OR 'family planning':ti,ab,kw OR 'sex education':ti,ab,kw OR 'sti':ti,ab,kw OR 'sexually transmitted infection':ti,ab,kw OR 'hvp vaccination':ti,ab,kw OR 'intimate partner violence':ti,ab,kw OR 'female genital mutilation':ti,ab,kw OR 'female genital cutting':ti,ab,kw OR 'fgm':ti,ab,kw OR 'maternal death':ti,ab,kw OR 'maternal morbidity':ti,ab,kw OR

	'maternal mortality':ti,ab,kw OR 'newborn morbidity':ti,ab,kw OR 'newborn mortality':ti,ab,kw OR 'maternal care':ti,ab,kw OR 'newborn care':ti,ab,kw OR 'infant':ti,ab,kw OR 'infancy':ti,ab,kw OR 'stigma':ti,ab,kw OR 'taboo':ti,ab,kw OR 'discrimination':ti,ab,kw OR 'gbv':ti,ab,kw OR 'gender':ti,ab,kw
AND	
<i>Low and middle-income countries (96)</i>	'low income countr*':ti,ab,kw OR 'low middle income countr*':ti,ab,kw OR 'developing countr*':ti,ab,kw OR 'middle income countr*':ti,ab,kw OR 'lmic':ti,ab,kw OR 'upper middle income countr*':ti,ab,kw OR 'albania':ti,ab,kw OR 'american samoa':ti,ab,kw OR 'argentina':ti,ab,kw OR 'armenia':ti,ab,kw OR 'azerbaijan':ti,ab,kw OR 'belarus':ti,ab,kw OR 'belize':ti,ab,kw OR 'bosnia':ti,ab,kw OR 'herzegovina':ti,ab,kw OR 'botswana':ti,ab,kw OR 'brazil':ti,ab,kw OR 'bulgaria':ti,ab,kw OR 'china':ti,ab,kw OR 'colombia':ti,ab,kw OR 'costa rica':ti,ab,kw OR 'cuba':ti,ab,kw OR 'dominica':ti,ab,kw OR 'dominican republic':ti,ab,kw OR 'ecuador':ti,ab,kw OR 'equatorial guinea':ti,ab,kw OR 'fiji':ti,ab,kw OR 'gabon':ti,ab,kw OR 'georgia':ti,ab,kw OR 'grenada':ti,ab,kw OR 'guatemala':ti,ab,kw OR 'guyana':ti,ab,kw OR 'indonesia':ti,ab,kw OR 'iran':ti,ab,kw OR 'iraq':ti,ab,kw OR 'jamaica':ti,ab,kw OR 'jordan':ti,ab,kw OR 'kazakhstan':ti,ab,kw OR 'kosovo':ti,ab,kw OR 'lebanon':ti,ab,kw OR 'libya':ti,ab,kw OR 'malaysia':ti,ab,kw OR 'maldives':ti,ab,kw OR 'marshall islands':ti,ab,kw OR 'mexico':ti,ab,kw OR 'montenegro':ti,ab,kw OR 'namibia':ti,ab,kw OR 'north macedonia':ti,ab,kw OR 'paraguay':ti,ab,kw OR 'peru':ti,ab,kw OR 'russia':ti,ab,kw OR 'samoa':ti,ab,kw OR 'serbia':ti,ab,kw OR 'south africa':ti,ab,kw OR 'saint lucia':ti,ab,kw OR 'saint vincent':ti,ab,kw OR 'grenadines':ti,ab,kw OR 'suriname':ti,ab,kw OR 'thailand':ti,ab,kw OR 'tonga':ti,ab,kw OR 'turkey':ti,ab,kw OR 'turkmenistan':ti,ab,kw OR 'tuvalu':ti,ab,kw OR 'venezuela':ti,ab,kw OR 'lower middle income countr*':ti,ab,kw OR 'algeria':ti,ab,kw

	<p>OR 'angola':ti,ab,kw OR 'bangladesh':ti,ab,kw OR 'benin':ti,ab,kw OR 'bhutan':ti,ab,kw OR 'bolivia':ti,ab,kw OR 'cabo verde':ti,ab,kw OR 'cambodia':ti,ab,kw OR 'cameroon':ti,ab,kw OR 'comoros':ti,ab,kw OR 'cote d'ivoire':ti,ab,kw OR 'Ivory Coast':ti,ab,kw OR 'djibouti':ti,ab,kw OR 'egypt':ti,ab,kw OR 'el salvador':ti,ab,kw OR 'eswatini':ti,ab,kw OR 'ghana':ti,ab,kw OR 'honduras':ti,ab,kw OR 'india':ti,ab,kw OR 'kenya':ti,ab,kw OR 'kiribati':ti,ab,kw OR 'kyrgyz':ti,ab,kw OR 'laos':ti,ab,kw OR 'lesotho':ti,ab,kw OR 'mauritania':ti,ab,kw OR 'micronesia':ti,ab,kw OR 'moldova':ti,ab,kw OR 'mongolia':ti,ab,kw OR 'morocco':ti,ab,kw OR 'myanmar':ti,ab,kw OR 'nepal':ti,ab,kw OR 'nicaragua':ti,ab,kw OR 'nigeria':ti,ab,kw OR 'pakistan':ti,ab,kw OR 'papua new guinea':ti,ab,kw OR 'philippines':ti,ab,kw OR 'sao tome':ti,ab,kw OR 'principe':ti,ab,kw OR 'senegal':ti,ab,kw OR 'solomon islands':ti,ab,kw OR 'sri lanka':ti,ab,kw OR 'tanzania':ti,ab,kw OR 'timor-leste':ti,ab,kw OR 'tunisia':ti,ab,kw OR 'ukraine':ti,ab,kw OR 'uzbekistan':ti,ab,kw OR 'vanuatu':ti,ab,kw OR 'vietnam':ti,ab,kw OR 'west bank':ti,ab,kw OR 'gaza':ti,ab,kw OR 'zambia':ti,ab,kw OR 'zimbabwe':ti,ab,kw OR 'low-income country':ti,ab,kw OR 'afghanistan':ti,ab,kw OR 'burkina faso':ti,ab,kw OR 'burundi':ti,ab,kw OR 'central african republic':ti,ab,kw OR 'chad':ti,ab,kw OR 'congo':ti,ab,kw OR 'eritrea':ti,ab,kw OR 'ethiopia':ti,ab,kw OR 'gambia':ti,ab,kw OR 'guinea':ti,ab,kw OR 'guinea-bissau':ti,ab,kw OR 'north korea':ti,ab,kw OR 'korea democratic people republic':ti,ab,kw OR 'haiti':ti,ab,kw OR 'liberia':ti,ab,kw OR 'madagascar':ti,ab,kw OR 'malawi':ti,ab,kw OR 'mali':ti,ab,kw OR 'mozambique':ti,ab,kw OR 'niger':ti,ab,kw OR 'rwanda':ti,ab,kw OR 'sierra leone':ti,ab,kw OR 'somalia':ti,ab,kw OR 'south sudan':ti,ab,kw OR 'sudan':ti,ab,kw OR 'syrian arab republic':ti,ab,kw OR 'syria':ti,ab,kw OR 'tajikistan':ti,ab,kw OR 'togo':ti,ab,kw OR 'yemen':ti,ab,kw</p>
--	--

**Table A-4. terms used in PsycINFO
abstract for all options**

Search	Search terms
Population	ab(("Adolescen*" OR "young" OR "youth" OR "teen" OR "teenag*" OR "child" OR "children" OR "paediatr*" OR "pediatr*" OR "juvenil*" OR "boy*" OR "girl*" OR "pubert*" OR "pubescen*"))
AND	
Exposure	ab(("swine flu" OR "H1N1" OR "flu pandemic" OR "Zika" OR "Ebola" OR "EVD" OR "EHF" OR "Ebola virus disease" OR "Ebola hemorrhagic fever" OR "COVID-19" OR "SARS-COV-2" OR "Novel Coronavirus" OR "2019-nCov"))
AND	
Outcome (in SRHR)	ab(("ASRH" or "SRH" or "SRHR" or "human right*" or "pregnan*" or "abort*" or "termination" or "antenatal" or "postnatal" or "perinatal" or "couple*" or "abuse" or "intercourse" or "rape" or "coerci*" or "violence" or "violation" or "incest" or "emergency pills" or "Long-acting reversible contraceptive" or "LARC" or "Misoprostol" or "mifepristone" or "methotrexate" or "meteneprost" or "prostaglandin" or "lilopristone" or "onapristone" or "oxytocin" or "sulprostone" or "family planning" or "sex education" or "STI" or "sexually transmitted infection" or "HPV vaccination" or "intimate partner violence" or "female genital mutilation" or "female genital cutting" or "FGM" or "maternal death" or "maternal morbidity" or "maternal mortality" or "newborn morbidity" or "newborn mortality" or "maternal care" or "newborn care" or "infant" or "infancy" or "stigma" or "taboo" or "Discrimination" or "GBV" or "gender"
AND	

<i>Low and middle-income countries (96)</i>	<p>ab(("low income countr*" OR "middle income countr*" OR "low middle income countr*" OR "developing countr*" OR "middle income countr*" OR "LMIC" OR "upper middle income countr*" OR "Albania" OR "American Samoa" OR "Argentina" OR "Armenia" OR "Azerbaijan" OR "Belarus" OR "Belize" OR "bosnia" OR "Herzegovina" OR "Botswana" OR "Brazil" OR "Bulgaria" OR "China" OR "Colombia" OR "Costa Rica" OR "Cuba" OR "Dominica" OR "Dominican Republic" OR "Ecuador" OR "Equatorial Guinea" OR "Fiji" OR "Gabon" OR "Georgia" OR "Grenada" OR "Guatemala" OR "Guyana" OR "Indonesia" OR "Iran" OR "Iraq" OR "Jamaica" OR "Jordan" OR "Kazakhstan" OR "Kosovo" OR "Lebanon" OR "Libya" OR "Malaysia" OR "Maldives" OR "Marshall Islands" OR "Mexico" OR "Montenegro" OR "Namibia" OR "North Macedonia" OR "Paraguay" OR "Peru" OR "Russia" OR "Samoa" OR "Serbia" OR "South Africa" OR "Saint Lucia" OR "saint Vincent" OR "grenadines" OR "Suriname" OR "Thailand" OR "Tonga" OR "Turkey" OR "Turkmenistan" OR "Tuvalu" OR "Venezuela" OR "lower middle income countr*" OR "Algeria" OR "Angola" OR "Bangladesh" OR "Benin" OR "Bhutan" OR "Bolivia" OR "Cabo Verde" OR "Cambodia" OR "Cameroon" OR "Comoros" OR "Congo" OR "Cote d'Ivoire" OR "Ivory Coast" OR "Djibouti" OR "Egypt" OR "El Salvador" OR "Eswatini" OR "Ghana" OR "Honduras" OR "India" OR "Kenya" OR "Kiribati" OR "Kyrgyz" OR "Laos" OR "Lesotho" OR "Mauritania" OR "Micronesia" OR "Moldova" OR "Mongolia" OR "Morocco" OR "Myanmar" OR "Nepal" OR "Nicaragua" OR "Nigeria" OR "Pakistan" OR "Papua New Guinea" OR "Philippines" OR "Sao Tome" OR "Principe" OR "Senegal" OR "Solomon Islands" OR "Sri Lanka" OR "Tanzania" OR "Timor-Leste" OR "Tunisia" OR "Ukraine" OR "Uzbekistan" OR "Vanuatu" OR "Vietnam" OR "West Bank" OR "Gaza" OR "Zambia" OR "Zimbabwe" OR "low-income country" OR "Afghanistan" OR "Burkina Faso" OR "Burundi" OR "Central African Republic" OR "Chad" OR "Congo" OR "Eritrea" OR "Ethiopia" OR "Gambia" OR "Guinea" OR "Guinea-Bissau" OR "North Korea" OR "Korea Democratic People's Republic" OR "Haiti" OR "Liberia" OR "Madagascar"</p>
---	--

	OR "Malawi" OR "Mali" OR "Mozambique" OR "Niger" OR "Rwanda" OR "Sierra Leone" OR "Somalia" OR "South Sudan" OR "Sudan" OR "Syrian Arab Republic" OR "Syria" OR "Tajikistan" OR "Togo" OR "Yemen"))
--	---

For peer review only

Table A-5. terms used in Cochrane library search
abstract, title and keywords for all options

<i>Search</i>	<i>Search terms</i>
<i>Population</i>	((“Adolescen*” OR “young” OR “youth” OR “teen” OR “teenag*” OR “child” OR “children” OR “paediatr*” OR “pediatr*” OR “juvenil*” OR “boy*” OR “girl*” OR “pubert*” OR “pubescen*”)):ti,ab,kw
AND	
<i>Exposure</i>	(“swine flu” or “H1N1” or “flu pandemic” or “Zika” or “Ebola” or “EVD” or “EHF” or “Ebola virus disease” or “Ebola hemorrhagic fever” or “COVID19” or “SARSCOV2” or “Novel Coronavirus” or “2019nCov”):ti,ab,kw
AND	
<i>Outcome (in SRHR)</i>	(“ASRH” or “SRH” or “SRHR” or “human right*” or “pregnan*” or “abort*” or “termination” or “antenatal” or “postnatal” or “perinatal” or “couple*” or “abuse” or “intercourse” or “rape” or “coerci*” or “violence” or “violation” or “incest” or “emergency pills” or “Long acting reversible contraceptive” or “LARC” or “Misoprostol” or “mifepristone” or “methotrexate” or “meteneprost” or “prostaglandin” or “lilopristone” or “onapristone” or “oxytocin” or “sulprostone” or “family planning” or “sex education” or “STI” or “sexually transmitted infection” or “HPV vaccination” or “intimate partner violence” or “female genital mutilation” or “female genital cutting” or “FGM” or “maternal death” or “maternal morbidity” or “maternal mortality” or “newborn morbidity” or “newborn mortality” or “maternal care” or “newborn care” or “infant” or “infancy” or “Discrimination” or “GBV” or “gender”) :ti,ab,kw
AND	
<i>Low and middle-income countries (96)</i>	(“low income countr*” or “middle income countr*” or “low middle income countr*” or “developing countr*”

	or “middle income countr*” or “LMIC” or “upper middle income countr*” or “Albania” or “American Samoa” or “Argentina” or “Armenia” or “Azerbaijan” or “Belarus” or “Belize” or “bosnia” or “Herzegovina” or “Botswana” or “Brazil” or “Bulgaria” or “China” or “Colombia” or “Costa Rica” or “Cuba” or “Dominica” or “Dominican Republic” or “Ecuador” or “Equatorial Guinea” or “Fiji” or “Gabon” or “Georgia” or “Grenada” or “Guatemala” or “Guyana” or “Indonesia” or “Iran” or “Iraq” or “Jamaica” or “Jordan” or “Kazakhstan” or “Kosovo” or “Lebanon” or “Libya” or “Malaysia” or “Maldives” or “Marshall Islands” or “Mexico” or “Montenegro” or “Namibia” or “North Macedonia” or “Paraguay” or “Peru” or “Russia” or “Samoa” or “Serbia” or “South Africa” or “Saint Lucia” or “saint Vincent” or “grenadines” or “Suriname” or “Thailand” or “Tonga” or “Turkey” or “Turkmenistan” or “Tuvalu” or “Venezuela” or “lower middle income countr*” or “Algeria” or “Angola” or “Bangladesh” or “Benin” or “Bhutan” or “Bolivia” or “Cabo Verde” or “Cambodia” or “Cameroon” or “Comoros” or “Congo” or “Cote d’Ivoire” or “Ivory Coast” or “Djibouti” or “Egypt” or “El Salvador” or “Eswatini” or “Ghana” or “Honduras” or “India” or “Kenya” or “Kiribati” or “Kyrgyz” or “Laos” or “Lesotho” or “Mauritania” or “Micronesia” or “Moldova” or “Mongolia” or “Morocco” or “Myanmar” or “Nepal” or “Nicaragua” or “Nigeria” or “Pakistan” or “Papua New Guinea” or “Philippines” or “Sao Tome” or “Principe” or “Senegal” or “Solomon Islands” or “Sri Lanka” or “Tanzania” or “Timor Leste” or “Tunisia” or “Ukraine” or “Uzbekistan” or “Vanuatu” or “Vietnam” or “West Bank” or “Gaza” or “Zambia” or “Zimbabwe” or “low income country” or “Afghanistan” or “Burkina Faso” or “Burundi” or “Central African Republic” or “Chad” or “Congo” or “Eritrea” or “Ethiopia” or “Gambia” or “Guinea” or “Guinea Bissau” or “North Korea” or “Korea Democratic People’s Republic” or “Haiti” or “Liberia” or “Madagascar” or “Malawi” or “Mali” or “Mozambique” or “Niger” or “Rwanda” or “Sierra Leone” or “Somalia” or “South Sudan” or “Sudan” or “Syrian Arab Republic” or “Syria” or “Tajikistan” or “Togo” or “Yemen”):ti,ab,kw
--	--

Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
TITLE			
Title	1	Identify the report as a scoping review.	Page 1
ABSTRACT			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	Page 2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	Page 4
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	Page 8
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	Page 14
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	Page 9 and 13
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	Page 13
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	Page 13
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	Page 13
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	Page 13
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	Page 14
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	Page 14

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	Page 14
RESULTS			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	Page 13
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	Page 13
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	Page 14
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	Page 10 and 11
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	Page 14
DISCUSSION			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	Page 11
Limitations	20	Discuss the limitations of the scoping review process.	Page 3
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	Click here to enter text.
FUNDING			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	Page 14

JB1 = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.

* Where *sources of evidence* (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.

† A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with *information sources* (see first footnote).

‡ The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JB1 guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.

§ The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

From: Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. Ann Intern Med. 2018;169:467–473. doi: 10.7326/M18-0850.