

BMJ Open Study protocol for investigating racial disparities in pain care: a comprehensive integration of patient-level and provider-level mechanisms with dyadic communication processes using a mixed-methods research design

Nao Hagiwara ¹, Emily Rivet,² Brian A Eiler,³ Christen Edwards,⁴ Nadia Harika,⁵ Shawn C T Jones,⁶ Amelia C Grover,² Peter Mende-Siedlecki⁷

To cite: Hagiwara N, Rivet E, Eiler BA, *et al.* Study protocol for investigating racial disparities in pain care: a comprehensive integration of patient-level and provider-level mechanisms with dyadic communication processes using a mixed-methods research design. *BMJ Open* 2025;**15**:e090365. doi:10.1136/bmjopen-2024-090365

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-090365>).

Received 24 June 2024
Accepted 12 March 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Professor Nao Hagiwara;
nh6ty@virginia.edu

ABSTRACT

Introduction Although many efforts have been made to reduce racial pain disparities over decades, the pain of black patients is still undertreated. Previous work has identified a host of patient and provider factors that contribute to racial disparities in healthcare in general, and consequently, may contribute to disparities in pain care in particular. That said, there has been limited clinically meaningful progress in eliminating these disparities. This lack of progress is likely because prior research has investigated the influence of patient and provider factors in isolation, rather than examining their interaction. Successful pain care requires constructive patient-provider communication, and constructive communication is both dyadic and dynamic. One well-accepted operationalisation of such dyadic processes is behavioural coordination. We hypothesise that the pain of black patients continues to be undertreated because black patients are more likely than white patients to participate in racially discordant medical interactions (ie, seeing other-race providers) and experience disruptions in behavioural coordination. We further hypothesise that disruptions in behavioural coordination will reflect patient and provider factors identified in prior research. We propose to test these hypotheses in the planned surgical context.

Methods and analysis Using a convergent mixed methods research design, we will collect data from at least 15 surgeons and their 150 patients (approximately equal number of black and white patients per surgeon). The data sources will include one surgeon survey, four patient surveys, video- and/or audio-recordings of preoperative consultations and medical chart reviews. The recorded preoperative consultations will be analysed both qualitatively and quantitatively to assess the magnitude and pattern of behavioural coordination between patients and surgeons. Those data will be linked to survey data and data from medical chart reviews to test our hypotheses.

Ethics and dissemination Ethical approval has been obtained from the Virginia Commonwealth University Institutional Review Board (HM20023574). Findings will be disseminated through presentations at scientific

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This research examines the role of dyadic and dynamic communication processes, specifically behavioural coordination, in influencing patient/provider factors and racial disparities in pain care and outcomes.
- ⇒ This study uses a convergent mixed methods research design to comprehensively analyse the nature of behavioural coordination during preoperative consultations.
- ⇒ We will use cross-recurrence quantification analysis to quantify the magnitude of coordination between the patient and surgeon at the non-verbal, paraverbal and verbal behavioural levels.
- ⇒ We will code transcribed preoperative consultations to analyse the context of behavioural coordination.

conferences, publications in peer-reviewed journals and speaking engagements with clinician stakeholders. We will also share the main findings from this project with patients via a newsletter on completion of the entire project.

INTRODUCTION

Racial disparities in pain care have broad and deep consequences for patients' lives. Untreated and persistent pain is a robust predictor of a diminished quality of life,¹⁻⁴ including, but not limited to, higher risk of experiencing depression and anxiety,⁴⁻⁷ challenges in resuming work^{8 9} and increased healthcare demands and burdens.⁹⁻¹¹ Although many efforts have been made to reduce racial pain disparities over decades,^{12 13} the pain of black patients is still undertreated. For example, a recent meta-analysis on the treatment of acute pain in emergency departments from 1990 to 2018

demonstrates that black patients were at least 60% less likely than white patients to receive analgesia.¹⁴ Furthermore, racial disparities exist across care stages (ie, pain assessment, treatment and management), contexts (eg, from the emergency room to postoperative care) and pain types (eg, acute, chronic, cancer),¹⁵ controlling for age, gender and pain intensity.¹⁶

Prior research, including our work, has identified both patient factors (eg, medical mistrust, perceived discrimination)^{17–24} and provider factors (eg, implicit prejudice, explicit stereotyping, pain perception bias)^{25–30} fuelling racial disparities in healthcare in general, and thus likely contributing to disparities in pain care in particular. In spite of the comprehensive catalogue of patient and provider factors examined in this literature, there has been limited clinically meaningful progress in reducing these disparities. We argue that this is mainly because of the fragmented approach that is typically employed within this literature, where individual-level factors predicting racial disparities in pain care are identified in isolation. This approach has grossly limited our ability to gain a comprehensive understanding of racial pain care disparities.³¹

According to widely used frameworks like the Social Communication Model of pain, interpersonal processes are fundamental to how people experience and express pain.^{32–33} Therefore, decisions regarding pain care are not made by a patient or a provider independently; they are a unilateral process. This suggests successful pain care requires constructive patient-provider communication, which is characterised as both dyadic and dynamic. Specifically, patients express their pain-related concerns, providers interpret patients' pain experience and respond

accordingly and this ongoing exchange lasts until they reach a mutual agreement about the optimal approach to pain management for the patient. Based on these premises, we developed a conceptual model that underscores these dyadic and dynamic communication processes as the crucial factor underlying the racial disparities in pain care and pain outcomes (figure 1).³¹

Our conceptual model³¹ posits that patient and provider factors predict patients' and providers' communication behaviours, respectively (Paths A and B). Our model diverges from prior research here and highlights the mutual influences between patients' and providers' communication behaviours (Path C). One well-accepted operationalisation of these dyadic processes in social psychology research is behavioural coordination. Behavioural coordination is characterised by the spatial and/or temporal matching in the rhythms or patterns of behaviours across individuals in an interaction (eg, patients and providers), such as synchrony, leader-and-follower dynamics and turn-taking.^{34–36} Coordination might manifest at the verbal (eg, word use, the contents of speech),^{34–37} paraverbal (eg, pitch, tone, speed of speech),^{37–38} or non-verbal levels (eg, posture, body movements, gaze)^{39–40} and could be cross-modal (eg, verbal-paraverbal coordination).^{34–36–37–41} Critically, coordination in dyads is thought to reflect meaningful social and cognitive connections between individuals. For example, behavioural coordination facilitates empathy and shared understanding—indicators of more constructive communication.^{42–44} However, dissimilarity or social distance between dyadic partners disrupts coordination.

In the context of patient-provider communication, patients in racially concordant medical interactions (ie,




Figure 1 Our new conceptual model that highlights the central role of dyadic and dynamic communication processes in racial pain care.

when a patient and provider share the same racial background) tend to report greater perceived similarity and shared understanding with their providers, regardless of patient race, compared with those in racially discordant interactions (ie, when a patient and provider have different racial backgrounds).^{45–49} Thus, we predict that racially discordant medical interactions are associated with less behavioural coordination than racially concordant ones at all behavioural levels.⁵⁰ Recent work supports this prediction in simulated medical interactions, where researchers experimentally created concordant versus discordant dyads by assigning participants to either a ‘yellow group’ or a ‘green group’. To make group membership personally meaningful, participants were told that their group assignment was based on their core beliefs and values. Researchers found that movement synchrony (one aspect of non-verbal coordination) between individuals playing the roles of ‘patient’ and ‘provider’ was higher in ideologically concordant dyads (ie, both the ‘patient’ and ‘provider’ belonged to the same colour group). Moreover, this coordination predicted reductions in pain reported by ‘patients’.⁵¹

However, no work has yet tested how coordination varies as a function of racial concordance and how it impacts pain care and experience in real-world medical contexts. Critically, due to the scarcity of black providers, black patients are more likely to see non-black providers even when they prefer to see black providers, whereas white patients are generally able to see white providers if they prefer.^{52–54} In other words, black patients are far less likely than white patients to experience racially concordant medical interactions. Therefore, we posit that behavioural coordination, in turn, serves as the primary mechanism through which patient/provider factors contribute to racial disparities in pain care and pain outcomes (Path D).

METHODS

The overarching goal of this research is to establish the dyadic and dynamic processes underlying patient-provider communication as the key mechanism through which patient and provider factors contribute to racial disparities in both patient-centred and clinical pain outcomes. We will test our hypotheses by addressing three specific aims: (Aim 1) to compare the levels, duration, patterns and context of behavioural coordination in preoperative consultations (both overall and during pain discussion specifically) between black and white patients; (Aim 2) to elucidate links between patient/provider factors and coordination in preoperative consultations; and (Aim 3) to identify specific aspects of behavioural coordination in preoperative consultations that contribute to racial disparities in post-surgical patient-centred outcomes (eg, pain management self-efficacy, quality of life) and clinical outcomes (eg, pain level, prescriptions). These aims will be achieved through an observational study with a convergent mixed methods research design. Table 1 provides an overview of the study design. The study is funded by the National Institutes of Health/National Institute of Nursing Research from 23 September 2022 through 20 June 2027 (R01NR020030).

Our conceptual model of racial pain care disparities is applicable to any medical context that requires pain care (eg, acute pain in the emergency context, chronic pain, cancer pain) given that disparities in pain treatment,⁵⁵ management^{12 14} and outcomes^{56–60} are mirrored in non-surgical contexts. However, the proposed research will focus on surgical pain care for theoretical and methodological reasons. Abundant evidence demonstrates that black patients experience worse surgical care and outcomes than any other patients.⁵⁹ Black patients receive comparatively

Table 1 An overview of the study designs

Aims	Data sources	Analyses	Goals
1	Recorded encounters	Quantitative analyses of behavioural coordination between patients and surgeons during preoperative consultations Qualitative analyses of patients’ and surgeons’ affect (‘this slices’ ratings) and themes of discussion topics during preoperative consultations	Compare the levels, amount, patterns of verbal, paraverbal and non-verbal coordination during preoperative consultations between black and white patients Compare the valence and context of behavioural coordination during preoperative consultations between black and white patients
2	Surgeon baseline survey Patient baseline survey Coded preoperative consultations	Quantitative analyses of associations between the patient and surgeon factors and behavioural coordination during preoperative consultations	Identify specific patient and provider factors that contribute to racial disparities in behavioural coordination
3	Surgeon baseline survey Patient baseline survey Coded preoperative consultations Patient post-recording survey Patient postoperative survey Medical chart review	Quantitative analyses of associations between behavioural coordination during preoperative consultations and post-surgical patient-centred and clinical pain outcomes	Identify specific aspects of behavioural coordination during preoperative consultations that contribute to racial disparities in patient-centred or clinical pain outcomes

less treatment for perioperative pain and/or report comparatively higher postoperative pain intensity across a wide variety of surgical procedures,⁶¹ including mastectomy,⁶² inguinal hernia repair,⁶³ oral surgery,⁶⁴ orthopaedics,^{65–68} appendectomy⁶⁹ and lumbar spine surgery.⁷⁰ Because surgeries are common (about 48 million patients are estimated to undergo surgeries annually⁷¹), a large number of black patients may receive suboptimal perioperative pain treatment and experience worse postoperative pain compared with white patients. Critically, effective patient-provider communication can reduce patient pain experience,^{72–73} and discussing pain care during preoperative consultations represents a key predictor of post-surgical pain care and outcomes.^{74–78} Therefore, effective communication during preoperative consultations represents a primary opportunity for eliminating disparities in pain management. However, patients and providers in racially discordant medical interactions are more likely to have reduced trust and poorer communication, leading to less dyadic decision-making, which may fuel downstream post-surgical pain disparities.⁷⁹

Participants

We will recruit at least 15 surgeons and 150 patients (approximately equal number of black and white patients for each surgeon) from a large university health system in the US South that serves diverse patient populations. A Monte Carlo Simulation with 1000 simulated data sets was conducted based on our smallest predicted effect (ie, a small to moderate effect of surgeon implicit racial prejudice on surgeon communication behaviours in Aim 2, which we expected to require the largest sample size). The power analysis (with 8 surgeon factors and 11 patient factors in a model) revealed that we can achieve adequate power (0.80) with a total of 15 surgeons and 150 patients completing the full suite of surveys.

The eligibility criteria for surgeons are that they: (1) are practicing as attending surgeons at one of the three participating surgery clinics; and (2) have preoperative consultations with patients to discuss surgeries for one of the 18 procedures listed in [table 2](#). We focus on those 18 procedures to minimise variability in pain intensity and duration, as well as procedural workflows and scheduling predictability. Note that the list of eligible procedures is a working list; new procedures will be added as we expand our partnerships with additional surgical divisions. The list of procedures was created and is being updated by surgeon consensus about expected post-surgical pain and validated by published guidelines⁸⁰ regarding opioid prescribing as a proxy for pain outcomes. That said, we will still control for procedure type as a covariate in our analyses. The eligibility criteria for patients are: (1) self-identify as either black/African American or white American; (2) be ≥18 years old; (3)

Table 2 Surgical procedures included in the present study (in alphabetical order)

Procedure	
Adrenalectomy	J-pouch
Amputation	Inguinal hernia repair
Anal condyloma excision	Liver resection
Colon resection	Mastectomy
Colostomy takedown (intra-abdominal)	Nephrectomy
Cystectomy	Parathyroidectomy
Fulguration	Prostatectomy
Gastric bypass	Sleeve gastrectomy
Haemorrhoidectomy	Thyroidectomy

have an appointment for any of the surgical procedures listed in [table 2](#) with one of the participating surgeons; and (4) be able to comprehend documents in English, written at a sixth-grade reading level. The participant recruitment began on 26 October 2023 and is expected to last through 31 May 2027.

Procedures

Data collection

Surgeons who meet the eligibility criteria and provide written consent are asked to complete a 25 min baseline survey online before their first recorded preoperative consultation with a participating patient. The project coordinator monitors patient appointments and identifies those who meet the eligibility criteria. Not every patient who attends preoperative consultations proceeds to have operations due to many different reasons (eg, their surgeon recommends non-surgical treatment after the consultation, the patient decides not to have surgery for their own reasons or the patient is transferred to a provider who is not participating in the research). To maximise the number of patients enrolled in our study with complete data, the project coordinator consults a surgeon on the team weekly to review imaging reports and notes in the charts to determine whether their consultation is likely to result in surgery. If we are uncertain about the likelihood of surgery, we consult with the participating surgeon for further guidance. Patients asked to complete a 45 min baseline survey over the phone following the consent and Health Insurance Portability and Accountability Act authorisation and before the scheduled appointment for preoperative consultation with their participating surgeon. (If necessary, surgeons are given the opportunity to complete the baseline survey in person via a laptop. Similarly, if necessary, patients receive a link to complete the baseline survey over the internet.)

On the day of the preoperative consultation appointment, the patient and surgeon are video-recorded and audio-recorded. We use two GoPro10 cameras

attached to the wall with Velcro: one focusing on the patient, another focusing on the surgeon. We attach the two cameras to the walls of the consultation room prior to the patient entering. The surgeon and patient meet as scheduled while we control the recording from outside of the room using The Remote, a GoPro accessory. Specifically, we stop the recording while the patient undresses for the physical examination and resume it once the patient has dressed again. We manage the camera (de)activation by monitoring the live audio stream. Moreover, a digital voice recorder captures the audio of the interaction to serve as a backup in the rare event of video failure. After the consultation, the patient is escorted to the waiting room (or stays in the examination room if the space is available) to complete a 5 min post-encounter survey.

Patients complete two additional surveys after the video-recorded preoperative consultation. Approximately 7 days before their scheduled surgery, a 5 min preoperative survey is conducted over the phone. Later, approximately 2 weeks after the surgery, a 30 min postoperative survey is completed over the phone after the patient's postoperative appointment. Surgeons receive a US\$50 e-gift card after completing the baseline survey. Patients receive a US\$50 e-gift card for completing the first three parts (ie, completing the baseline survey, participating in a video-recorded preoperative consultation and completing the post-encounter survey), and a US\$20 gift card for completing the preoperative and postoperative surveys. **Figure 2** summarises the survey data collection timeline. Finally, we conduct medical chart reviews to code: (1) comorbidities at the time of preoperative consultations; and (2) clinical outcomes between the time of surgery and the postoperative survey (about a 2-week period).




Figure 2 Overview of the study flow.

Measures

Table 4 provides an overview of the variables assessed in each survey.

Surgeon baseline survey (between consent and the first recorded preoperative consultation)

Demographic information

We assess self-reported age, ethnicity, race, gender identity and US citizenship.

Professional information

Surgeons report the year in which they received their medical degree and also when they started working at the current institution.

Prior experience with cultural competency training

Surgeons are asked to indicate if and when they last took part in cultural competency training. Providers who have participated in such training are further asked four additional questions: (1) What the main goal of the most recent training they attended was (free response box); (2) Whether the training was required (Yes/No); (3) Whether the most recent training was the first training they attended in cultural competency and related topics (Yes/No); and (4) How many hours of cultural competency-related training they have completed over the last 2 years (an approximate number). Lastly, all surgeons are asked to rate their ability to deliver culturally competent care on a scale of 1 (Poor) to 5 (Outstanding).

Prior experience with communication skills training

Similar to the assessment of prior experience with cultural competency training, surgeons are first asked to indicate if and when they last attended training in communication skills. Those who had participated in communication skills training are further asked to respond to four additional questions: (1) What the main goal of the most recent training they attended was (free response box); (2) Whether the training was required (Yes/No); (3) Whether the most recent training was the first training they attended in communication skills and related topics (Yes/No); and (4) How many hours of communication skills-related training they have completed over the last 2 years (an approximate number). All surgeons also rate their communication skills on a scale of 1 (Poor) to 5 (Outstanding).

Prior experiences with black patients

We assess surgeons' past experiences with treating black patients using two items: (1) the frequency of their interactions with black patients in their practice or training (rated on a scale from 1 (Never) to 6 (All the time)); and (2) their self-assessment of communication skills when interacting with black patients (rated on a scale from 1 (Poor) to 5 (Outstanding)).

Perceptual bias in pain recognition

Surgeons' perceptual bias in pain recognition is assessed with a computerised task designed to isolate thresholds

Table 3 An overview of the variables assessed in the surveys

	Surgeons	Patients
Baseline	<ul style="list-style-type: none"> ▶ Demographic information ▶ Professional information ▶ Prior experience with cultural competency training ▶ Prior experience with communication skills training ▶ Prior experiences with black patients ▶ Perceptual bias in pain recognition ▶ Implicit racial prejudice ▶ Explicit racial prejudice ▶ Implicit racial stereotyping ▶ Explicit racial stereotyping ▶ Stereotype threat 	<ul style="list-style-type: none"> ▶ Demographic information ▶ Socioeconomic information ▶ Medical history ▶ Perceived racial discrimination ▶ General trust in surgeons ▶ General satisfaction with healthcare ▶ Pain sensitivity and fear of pain ▶ Self-efficacy for managing pain ▶ State pain level ▶ Quality of life ▶ Depressive symptomatology ▶ Anxiety
Post-encounter		<ul style="list-style-type: none"> ▶ Surgeon-specific trust ▶ Surgeon-specific satisfaction ▶ Expected pain intensity and duration
Preoperative		<ul style="list-style-type: none"> ▶ The number of days until surgery ▶ Resources given for pain and/or pain management ▶ Expected pain intensity and duration
Postoperative		<ul style="list-style-type: none"> ▶ State pain level ▶ Quality of life ▶ Depressive symptomatology ▶ Anxiety ▶ Future healthcare utilisation intentions

for seeing pain on black versus white faces. This task has been extensively used in non-clinical populations. For example, a recent meta-analysis comprising 40 studies and over 6000 participants demonstrated that racial bias in pain perception generalises across perceiver race and robustly predicts a corresponding tendency to recommend prescribing relatively less pain reliever to black (vs white) targets.⁸¹ Moreover, across this work, the influence of this perceptual bias on hypothetical treatment recommendations persists even when accounting for explicit racial biases and stereotypes.^{25 82} While this work was primarily conducted on non-clinical populations, forthcoming work in progress demonstrates that each of these findings generalises to perceivers with training or experience working in clinical contexts.

Surgeons see equal numbers of digitally rendered black and white faces making pain expressions (six morphs per target, ranging from neutral to 100% pain in 20% increments, with expression intensity and facial structure equated across race). For each morph, presented one at a time, surgeons make a continuous rating of perceived pain in response to the question ‘Is this face in pain?’ on a 1–7 scale (1=‘Definitely not in pain’, 7=‘Definitely in pain’). Following our procedures in previous work, these ratings are used to compute points-of-subjective-equality (PSEs) representing surgeons’ perceptual thresholds for recognising pain—in other words, the point in the

morph continuum where someone would be equally likely to see a face as being in pain versus not in pain. Specifically, these ratings are rescored to a 0–1 scale and, using the curve-fitting function fit in MATLAB, fit to an S-shaped psychometric curve approximated by a cumulative normal distribution function. Higher PSE values represent higher pain perception thresholds. This task reliably predicts racial bias in pain treatment recommendations^{25 82} over and above explicit racial bias or racial stereotypes regarding status, strength, threat or pain sensitivity.

Implicit racial prejudice

A well-validated Race Implicit Association Test (Race IAT⁸³) is used to evaluate surgeons’ implicit racial prejudice. Surgeons categorise photographs and words that appear on the computer screen into four groups: two racial groups (white vs black) and two evaluations (good vs bad), which are presented in pairs. The premise is that individuals respond more quickly when the racial category and evaluation mapped onto the same response are strongly associated compared with when they are weakly associated. In the Race IAT, photographs of faces represent the two racial groups, while 16 words (8 positive: marvellous, superb, pleasure, beautiful, joyful, glorious, lovely, wonderful; and 8 negative: tragic, horrible, agony, painful, terrible, awful, humiliate, nasty) represent the

evaluative categories. We used the standard algorithm to compute a D score, which ranges from -2.0 to 2.0, to measure implicit associations (average inter-item consistency $\alpha=0.78^{84}$); this approach has been well validated.^{85–88} Higher scores indicate greater pro-white and anti-black preference.

Explicit racial prejudice

Surgeons' explicit racial prejudice is assessed with two measures: the feeling thermometer and Symbolic Racism Scale 2000 (SR2K). The feeling thermometer is a single scale measure in which surgeons rate their feelings toward each racial social group on a scale from 0 (very cold/unfavourable) to 100 (very warm/favourable^{89 90}). Similar to D scores in the IAT, composite explicit pro-white/anti-black prejudice scores are calculated by subtracting the score for black Americans from a score for white Americans. Positive values indicate explicit preference for white Americans over black Americans, whereas negative values indicate explicit preference for black Americans over white Americans. A value of zero indicates no explicit preference between black and white Americans. The SR2K is the most up-to-date scale of symbolic racism that is designed to measure people's belief system based on the ideas that racial discrimination is no longer an issue in the USA and that black Americans' demands for fairness are unjustified.^{91–93} The SR2K has been found to be reliable (ie, an average α around 0.75) and well-validated in more than two dozen empirical studies.⁹⁴

Implicit racial stereotyping

Implicit racial stereotyping is assessed using the medical cooperativeness IAT, an implicit measure of racial stereotyping commonly used in research on racial health disparities.^{95–97} This task mirrors the Race IAT but replaces the two evaluative categories (good vs bad) with two stereotyping categories (medically cooperative vs medically uncooperative). Specifically, the 16 words categorised into evaluations (good vs bad) in the Race IAT are substituted with 12 words categorised into stereotypes (6 words associated with cooperativeness: willing, cooperative, compliant, reliable, adherent, helpful; and 6 words associated with uncooperativeness: reluctant, doubting, hesitant, apathetic, resistant and lax). D scores are calculated using the same methodology as for the Race IAT. Higher positive values indicate strong implicit associations of white (vs black) Americans with medical cooperativeness, whereas higher negative values indicate stronger implicit associations of black (vs white) Americans with medical cooperativeness.

Explicit racial stereotyping

Surgeons respond to two measures of stereotypes: a 32-item assessment aimed at evaluating general racial stereotypes (16 items for black Americans and 16 items for white Americans, such as unintelligent, lazy, low socioeconomic status),⁹⁸ and a 22-item evaluation designed to gauge stereotypes specifically within the medical context

(11 items for black Americans and 11 items for white Americans, such as medical cooperativeness, health literacy, motivation).⁹⁸ These measures were adapted from previous studies of racial stereotyping.^{28 99} Surgeons are instructed to review statements and indicate the degree to which each statement describes characteristics of black versus white Americans using a Likert scale ranging from 1 (very untrue) to 6 (very true). Consistent with the procedure used in prior research, composite scores are calculated for general and medicine-specific stereotypes separately. More specifically, first, negatively worded items will be reverse-coded, and the average score for each racial category will be computed. Subsequently, the average score for black Americans will be subtracted from the average score for white Americans. Higher positive values indicate a stronger explicit association of white Americans (vs black Americans) with more positive stereotypes, while higher negative values indicate a stronger explicit association of black Americans (vs white Americans) with more positive stereotypes.

Stereotype threat

We assess the degree to which surgeons are concerned about being perceived as prejudiced by black patients by using a modified version of the Explicit Stereotype Threat Scale.¹⁰⁰ The measure consists of four items (eg, 'I worry that my black patients might stereotype me as racist because of my race'), and the scale ranges from 1 (Strongly disagree) to 4 (Strongly agree). Higher scores indicate higher levels of stereotype threat.

Patient baseline survey (between consent and recorded preoperative consultation)

Demographic information

We assess self-reported age, race, ethnicity, gender identity, pronouns they use, sexual orientation, US citizenship, height and weight (to compute body mass index), physical activity, marital status, education, income and health insurance.

Medical history

Patients also indicate if, for what and how many surgeries they have had previously. Patients also report if they have ever been prescribed opioids for severe pain and whether it was helpful.

Perceived racial discrimination

Patients are asked to complete two well-validated measures of perceived racial discrimination that have been used in health research and social psychology research previously. The first is the Brief Perceived Ethnic Discrimination Questionnaire-Community Version ($\alpha=0.87$) that assesses both daily and lifetime experience of multiple forms of discrimination (eg, exclusion, stigmatisation, threat) in multiple domains (eg, work, public places).¹⁰¹ Participants will be asked to use a scale ranging from 1 (Never) to 5 (Very often). The second measure assesses the perceptions of racial discrimination at both personal and group level.^{102 103}

More specifically, five items are designed to assess the degree to which patients have experienced discrimination personally ($\alpha=0.84$), and three items are designed to assess the degree to which patients perceive other members of their racial group in general experience racial discrimination ($\alpha=0.84$). Both subscales are assessed by using response items that range from 1 (Strongly disagree) to 5 (Strongly agree).

General trust in surgeons

We assess patients' expectations about their trust in surgeons in general at the participating health system with a modified version of the Wake Forest Physician Trust Scale, which has been found to have better internal consistency ($\alpha=0.93$, test-retest reliability=0.75), validity, discriminability and scale distribution as compared with other trust scales.^{104 105} To account for patients having never met with a surgeon, they are instructed to indicate how they expect to feel regarding trust in their surgeons. The response items range from 1 (Strongly disagree) to 5 (Strongly agree).

General satisfaction with care

We use a modified version of the Patient Satisfaction Questionnaire Form III¹⁰⁶ (PSQ-III) to assess patients' satisfaction with the specific healthcare system in general, as opposed to satisfaction with surgeons within the specific healthcare system, because the vast majority of patients have had no prior experience with surgeons. In particular, we use three subscales: general satisfaction ($\alpha=0.88$), interpersonal aspects ($\alpha=0.82$) and communication ($\alpha=0.82$). The response items range from 1 (Strongly disagree) to 5 (Strongly agree).

Pain sensitivity and fear of pain

We assess pain sensitivity and fear of pain because they are predictive of post-surgical pain outcomes.^{107–117} Pain sensitivity is assessed with the Pain Sensitivity Questionnaire (PSQ). The PSQ correlates with pain induced experimentally in healthy participants^{118 119} and chronic pain patients.^{120 121} Fear of pain is assessed with the short form of the Pain Anxiety Symptoms Scale,¹²² which has shown strong internal consistency, reliability and good predictive and construct validity.^{123–126}

Self-efficacy for managing pain

Self-efficacy for managing pain is defined as the confidence individuals have in engaging in activities while in pain. It has been shown to predict not only experiences of and reactions to pain but also overall clinical outcomes.^{127–132} Pain self-efficacy is assessed with the well-validated Pain Self-Efficacy Questionnaire,¹³³ which has been used in multiple clinical settings.^{134–137}

State pain level

We assess state pain level with the Numerical Rating Scale (NRS-11), the most widely used, well-validated^{138–141} scale in the clinical setting. The scale ranges from 0 (no pain) to 10 (very severe).

Quality of life

Quality of life is assessed with the 36-item Short Form (SF-36) Health Survey,¹⁴² which has been validated in diverse populations,¹⁴³ including pain patients.^{144–146} We use the SF-36 Health Survey (rather than pain-specific quality of life) because patients in our sample have a variety of medical conditions although they all require surgeries.

Depressive symptomatology

We use the Personal Health Questionnaire Depression Scale (PHQ-8),¹⁴⁷ as opposed to the more widely used PHQ-9,^{148 149} to avoid directly asking patients to report their suicidal thoughts (ie, the last item in the PHQ-9). This 8-item scale is designed to capture both overall depression severity and specific symptoms, has been well-validated,¹⁵⁰ is strongly correlated with the PHQ-9 ($r=0.99$) and can be self-administered.

Anxiety

Anxiety is assessed using the State-Trait Anxiety Inventory,¹⁵¹ a 40-item scale measuring acute and chronic anxiety symptoms. The scale has been used extensively, including in black Americans.^{152 153}

Patient post-encounter survey (immediately after recorded preoperative consultation)

Surgeon-specific trust

We assess patient trust in the specific surgeon they have just met with a modified version of the Wake Forest Physician Trust Scale.¹⁰⁴ Items are modified to reflect the specific surgeon a patient saw during their preoperative consultation.

Surgeon-specific satisfaction

Patient satisfaction with the specific surgeon they have just met with is assessed with a modified version of the PSQ-III.¹⁰⁶ Again, items are modified to reflect the specific surgeon a patient saw during their preoperative consultation.

Expected pain intensity and duration

Patients report their expected post-surgical pain, once again using the NRS-11,^{138–141} ranging from 0 (no pain) to 10 (worst possible pain). Specifically, patients report these expectations for five separate points in time—immediately after their procedure, 1 week after their procedure, 2 weeks after their procedure, 1 month after their procedure and 3 months after their procedure. For each time point, patients are given the opportunity to select a 'Not sure' option.

Patient preoperative survey (approximately 7 days before surgery)

First, patients report the number of days until their surgery. Next, patients are asked about the resources they have been directed to regarding pain and/or pain management. Patients answer separate questions regarding resources (1) their surgeon recommended, (2) other medical professionals recommended, (3) family, friends or coworkers recommended or (4) that

they themselves sought out. In each case, patients first give a 'Yes', 'No' or 'I'm not sure' response. In instances where patients supply a 'Yes' response, a free response box appears, asking them to describe the resources they were directed to or sought out.

Patients then complete the same adapted NRS-11^{138–141} measure of their expected post-surgical pain using the same approach (and at the same time points) as in the patient baseline survey.

Patient postoperative survey (approximately 2 weeks after surgery)

State pain level, quality of life, depressive symptomology, state and trait anxiety and future healthcare service utilisation intentions are once again assessed using the same methods as in the patient baseline survey. For each of these measures, we compute a difference score (via subtraction) representing the change in each outcome from baseline to postoperative follow-up.

Medical chart reviews

We conduct medical chart reviews and code comorbidities at the time of preoperative consultations. Comorbidities will be used to control for potential systematic variability in both post-surgical patient-centred and clinical outcomes. We will also code clinical outcomes between the time of surgery and the postoperative survey (about a 2-week period). Specifically, we will code (1) type of anaesthetic used in procedure¹⁵⁴; (2) pain prescriptions (ie, type, dosage and duration); (3) the number of healthcare visits for post-surgical pain management (greater visits indicating poorer pain management); (4) documented pain score (if any)¹⁵⁵; (5) surgical complications; and (6) pain-related comorbidities that are independent of pain directly related to the surgical procedure. Finally, surgeons on the research team will independently code the appropriateness of anaesthetic/pain prescriptions (−3 (pain is grossly undertreated) to +3 (pain is grossly overtreated), with 0 being the right pain treatment) for a given surgical procedure using the published guidelines⁸⁰ while taking into account comorbidities. They will be kept blind to patient race or surgeons. Their scores will be averaged to compute a composite score.

Analysis

Analysis of recorded preoperative consultations

We are going to use recorded preoperative consultations to assess the levels, duration, patterns and context of behavioural coordination. Specifically, we will (1) obtain numerical quantifications of verbal, paraverbal and non-verbal coordination between patients and surgeons (eg, the level and duration of coordination, the frequency of disruptions to coordination, patterns of coordination); (2) qualitatively describe the valence and context of quantified verbal, non-verbal and paraverbal coordination; (3) test whether the level of behavioural coordination during preoperative consultations is higher for white (vs black) patients; and (4) identify which aspects of behavioural

coordination differ between black and white patients and understand why these differences occur.

Quantitative analysis of behavioural coordination

We will assess coordination between patients and surgeons in preoperative consultations, both overall and during pain discussion specifically, using cross recurrence quantification analysis (CRQA). CRQA is a cutting-edge, fully automated statistical analysis that enables us to quantify the level and duration of behavioural coordination at verbal, paraverbal and non-verbal levels during dyadic social interactions. It also provides information about who initiates changes in the levels of coordination and whether these changes are followed by the other member of the dyad (ie, identification of a leader and a follower). CRQA takes two (ie, patient and provider) numbers (or 'signals') in chronological order as input and outputs quantifications of the coordination dynamics of the dyad (recurrence rate, % determinism, diagonal length, entropy, laminarity, trapping-time). CRQA analyses a recurrence plot—a graph that crosses the two signals by putting one on the x-axis and the other on the y-axis and plots a recurrence point when the two signals have overlapping values¹⁵⁶ (see figure 3 for examples). All CRQA metrics are derived from calculations on this graph.¹⁵⁷

We will quantify the coordination dynamics that characterise patient–surgeon interactions in terms of verbal (eg, using unique number-word assignments as input), paraverbal (eg, extracting waveform and specific frequencies/amplitudes from audio signal) and non-verbal behaviours (eg, overall body movement magnitude derived from changes in pixel colour and intensity).^{158 159} Specifically, we will include recurrence rate (coordination over time), per cent determinism (magnitude of coordination over time), diagonal lengths (time-location and length of coordination), entropy (a measure of the unpredictability of the interaction), laminarity (stability of coordination) and trapping time (independence). Technically, we will use the statistical computing environment R/RStudio¹⁵⁹ to calculate CRQA measures for patient–surgeon dyads across each communication modality. Using the R package 'crqa',¹⁶⁰ the coordination dynamics for each dyad will be modelled to optimise hyperparameters (ie, embedding dimension, delay and radius) by finding the lag at which minimal mutual information is observed, fixing the relative information percentage gain in the false nearest neighbours test to 5% relative to the first embedding dimension, and testing a range of radius values that result in such recurrence rates 2% and 10% ($0.02 > \text{Recurrence Rate } \% > 0.10$). CRQA calculations have been described in detail elsewhere^{159–161} and these summary metrics will be used to operationalise communication dynamics across all three modalities.^{162–164} These values can be used as dependent or independent variables depending on the question (eg, does the patient or surgeon lead the interaction (recurrence rate); are racially concordant interactions more/less coordinated (% determinism) or unpredictable (entropy); at what




Figure 3 Example recurrence plots that depict different system types.

time points are communication most synchronised (diagonal lengths)?).

Qualitative analysis of the valence of behavioural coordination

We will use the ‘thin slices’ methodology to obtain ratings of patients’ and surgeons’ positive and negative affect. Research on person perception has shown that people can quickly and accurately judge affect with thin slices (between 2 s and 5 min) of non-verbal behaviours.^{165–167}

Our work has also shown that ratings of physician affect obtained from naive observers using ‘thin slices’ are associated with physicians’ implicit and explicit prejudice.²⁰ Critically, other work has shown the utility of the thin slices methodology in capturing patient emotion specifically during pain discussions.^{168 169} We will create ‘thin slices’ in two ways (there can be overlaps between the two). First, we will edit the video recordings to create the pain versus general (eg, non-pain) discussion sections. For each, we will create a 2 min excerpt, which will begin the moment either a patient or a surgeon initiates a discussion about pain or a topic unrelated to pain (eg, explanation of a surgical procedure; but excluding greeting at the beginning) and will end after 2 min elapse, regardless of whether the discussion on the topic is still going. Second, we will create two 2 min excerpts for each point of disruption to behavioural coordination identified by the CRQA: one capturing patient–surgeon communication up to the disruption, another capturing communication immediately after the disruption.

Each excerpt will be rated by 20 racially diverse research assistants (RAs). Overthinking and reasoning can often disrupt judgement accuracy, and people usually make better judgements when they do not try to explain their judgements.^{164 165} Thus, we will instruct RAs to report their first impressions. RAs will view excerpts one by one and provide their judgements of: (1) patient affect

(eg, positive: cheerful, friendly, warm; negative: cold, nervous, stressed); and (2) surgeon affect.²⁰ The order of these categories and of excerpts within each category will be randomised across RAs. Ratings will be averaged across RAs because aggregated judgements better predict outcomes than judgements made by single observers.¹⁶⁵

Qualitative analysis of the context of behavioural coordination

The recorded preoperative consultations will be transcribed verbatim to identify discussion themes in five steps. Step 1 involves unitisation of thought units. Unitisation is a step in qualitative research where researchers break down a large amount of qualitative information (eg, transcripts, written responses or observations) into smaller pieces called units that make sense on their own (eg, utterances, paragraphs, discrete thoughts).^{170–172} In the current project, we will use discrete thoughts as units. Two RAs will be trained to unitise a few transcripts until they achieve consensus. After the training, the RAs will unitise the remaining transcripts. They will meet with one of the senior research team members after unitising every 10 transcripts to discuss discrepancies and achieve consensus. Step 2 involves categorisation of thought units into three groups: (1) thought units related to pain; (2) thought units not related to pain; and (3) thought units that are being discussed at the moment of the disruptions to behavioural coordination identified by the CRQA. Thought units categorised into the third group are not mutually exclusive from those categorised in the other two categories. Step 3 involves the development of a codebook for each group of thought units. Several senior research team members will review approximately 50% of the thought units within each group inductively to identify a comprehensive list of themes (eg, emotional expressions, affirmation of concerns, questions about post-surgical pain and explanation of pain management

Table 4 Integration of qualitative and quantitative data

Qualitative questions	Behavioural coordination quantified by the CRQA
<p>1. What are the average patient affect, the average surgeon affect and the main discussion themes</p> <p>a. During discussions unrelated to pain in patient–surgeon dyads</p> <p>b. During pain discussions in patient–surgeon dyads</p>	<p>▶ With higher (+1 SD) vs lower (–1SD) levels of coordination overall as well as during pain discussions specifically?</p> <p>▶ With longer (+1 SD) vs shorter (–1SD) duration of coordination overall as well as during pain discussions specifically?</p> <p>▶ Within each type of coordination (eg, synchrony, turn-taking, leader-follower) overall as well as during pain discussions specifically?</p> <p>▶ With different sequential patterns of coordination (ie, increasing, decreasing, stable, fluctuating) overall as well as during pain discussions specifically?</p>
<p>2. What affect a patient and a surgeon each were displaying and what were they discussing:</p>	<p>▶ Immediately before vs after disruptions to coordination?</p>

CRQA, cross recurrence quantification analysis.

plans). Identified themes will then be refined, collapsed and organised into categories. For example, ‘emotion expressions’ and ‘affirmation of concerns’ may be organised under a higher order theme of ‘discussions of emotions’, and ‘emotion expressions’ may be further refined and divided into ‘positive emotion (eg, relief, excitement)’ and ‘negative emotions (eg, fear, anxiety)’. The codebooks will provide coding procedures, rules for coding, descriptions and examples of the codes. Step 4 involves the coding of the remaining thought units. Two RAs will be trained on the three codebooks using ~10% of these thought units (or until they reach consensus) within each group. The codebooks will be fine-tuned during this period. After the training, the RAs will code the remaining thought units while meeting with a senior research team member each time after they code an additional 10% of thought units to discuss discrepancies and reach consensus. We will use NVivo to categorise, sort and link data across transcripts. We will assess the frequency of each theme within each preoperative consultation. Step 5 involves integration of qualitative and quantitative data (table 4). We will use qualitative results to characterise: (1) quantitative variation in the levels (eg, higher vs lower), duration (eg, longer vs shorter), type (ie, synchrony, turn-taking, leader-follower) and sequential patterns (ie, increasing, decreasing, stable, fluctuating) of behavioural coordination across patient–surgeon dyads; and (2) the occurrences of disruptions to behavioural coordination.

Aim 1: to compare the levels, duration, patterns and context of behavioural coordination in preoperative consultations between black and white patients

To test our hypothesis that behavioural coordination during preoperative consultations is higher for white patients than for black patients, we will conduct a regression using a Generalised Estimating Equations (GEE) framework. GEE treats surgeons as a nuisance parameter and corrects biased estimates due to non-independence of a surgeon seeing multiple patients. This analytical approach is common in the existing

patient-provider communication literature^{173–177} and was used successfully in our prior work.^{178 179} The regression will include patient factors (assessed in the patient baseline survey, such as education) and surgeon factors (assessed with the surgeon baseline survey, such as cultural competency training), and clinical factors (eg, procedure type; as well as those obtained through medical chart reviews, such as comorbidities) that are significantly correlated with coordination level as covariates. We will also test if the results are moderated by patient–surgeon racial concordance/discordance. Approximately 80% of the eligible surgeons are white; thus, the majority of black patients are likely to engage in racially discordant consultations, whereas the majority of white patients are likely to engage in racially concordant consultations. However, because the study will be open to surgeons of all racial and ethnic identities, we will explicitly examine the role of racial concordance as a potential moderator. If the statistical power is too weak to detect the expected moderating effect of patient-provider racial concordance/discordance due to a small number of surgeons who identify as black/African American, we will treat patient-provider racial concordance/discordance as a covariate in our analyses instead. Additionally, we will conduct a descriptive examination of data based on patient–surgeon racial concordance/discordance to identify patterns and trends, which will inform research questions and hypotheses for future studies.

To further identify which additional specific aspects of behavioural coordination differ between black and white patients, we will conduct the same set of regressions with GEE (with patient race as a predictor, patient and surgeon factors as covariates and racial concordance as a moderator) for each aspect of quantified behavioural coordination. We will correct the family-wise error rate due to multiple testing with the Bonferroni correction procedure.^{180 181} Finally,

we will explore why aspects of behavioural coordination identified by the previous analyses differ across patient race and patient–surgeon racial concordance by reviewing the integrated data (table 3) and exploring the patterns of valence and themes which characterise coordination in dyads with black versus white patients.

Aim 2: to elucidate links between patient/provider factors and coordination in preoperative consultations

We will assess relationships between patient factors recorded in the patient baseline survey and specific aspects of behavioural coordination (eg, levels, duration, patterns) assessed in Aims 1. Patient factors can be distilled into race-related factors and health-related factors (online supplemental table 1). In Step 1, we use confirmatory factor analysis (CFA) to examine the latent structure of the patient-level variables and determine whether any patient factors can be aggregated in subsequent analyses. In Step 2, we identify covariates that may impact behavioural coordination by computing correlations among all variables assessed in the patient baseline survey and medical chart review. In Step 3, we compute partial correlations between each patient factor and each measure of behavioural coordination, while controlling for covariates identified in Step 2. In Step 4, we will conduct regressions with GEE using the patient factors remaining significant in Step 3, with the same set of covariates included. This approach enables us to examine which patient factors are most strongly associated with coordination. Again, we will correct the family-wise error rate due to conducting multiple regression tests with the Bonferroni correction procedure. Finally, in Step 5, we will explore whether the associations between patient factors and specific aspects of behavioural coordination identified in Step 4 vary by patient race (black vs white) or patient–surgeon racial concordance.

We will also examine relationships between provider factors recorded in the surgeon baseline survey and specific aspects of behavioural coordination (eg, levels, duration, patterns) assessed in Aim 1. Provider factors can be collapsed into health-specific and general racial bias factors. We will first use CFA to determine if any provider factors can be aggregated in subsequent analyses (Step 1). We then identify covariates that may impact coordination by computing correlations among all variables assessed in the surgeon baseline survey (Step 2). After assessing partial correlations (controlling for covariates identified in Step 2) between each provider factor and each measure of behavioural coordination (Step 3), we will use GEE to examine which provider factors are most strongly associated with behavioural coordination while controlling for the same set of covariates (Step 4). Finally, we will

explore whether these relationships vary by patient race (black vs white) or patient–surgeon racial concordance (Step 5).

Aim 3: to identify specific aspects of behavioural coordination in preoperative consultations that contribute to racial disparities in post-surgical patient-centred and clinical outcomes

In Step 1, we identify outcomes of interest demonstrating racial disparities. We will conduct regressions with GEE to assess whether any of the outcomes listed in online supplemental table 2 vary between black and white patients. In Step 2, we identify covariates. We will conduct correlations to assess whether any potentially confounding factors assessed at baseline (eg, demographic factors, individual differences in pain sensitivity and pain anxiety) covary with change from baseline to the preoperative consultations or 2-week post-surgery follow-up in terms of the patient-centred or clinical outcomes identified in Step 1. In Step 3, we calculate partial correlations assessing the extent to which behavioural coordination correlated with racial disparities in the outcomes (identified in Step 1), while controlling for covariates identified in Step 2. Finally, in Step 4, we conduct a series of regressions using GEE for each outcome that remains significant in Step 3. We will include the same covariates as in Step 3 and correct for multiple comparisons using the Bonferroni procedure.

Patient and public involvement

Patients or the public are not involved in the design, or conduct, or reporting, or dissemination plans of our research.

ETHICS AND DISSEMINATION

The study is conducted in accordance with the Declaration of Helsinki, and ethical approval has been obtained from the Virginia Commonwealth University (VCU) Institutional Review Board (HM20023574). Both surgeons and patients are encouraged to ask questions during the consenting process and before they sign the consent form. It is made clear to the surgeons and patients that they have the right to withdraw from the study at any time even after they have signed the consent form. This study involves no more than minimal risk. However, in the event of a serious adverse event, we will report this to the VCU Institutional Review Board within 48 hours. Findings will be disseminated through presentations at scientific conferences, publications in peer-reviewed journals and speaking engagements with clinician stakeholders. We will also share the main findings from this project with patients via a newsletter on completion of the entire project.

DISCUSSION

This proposed research seeks to shift current research paradigms in racial pain care disparities by both taking a novel theoretical and methodological approach. Our theoretical innovations are reflected in our conceptual model,³¹ which places unprecedented importance on dyadic and dynamic communication processes (eg, behavioural coordination) as the key mechanism through which patient/provider factors contribute to racial disparities in pain care and outcomes. Studying the entire chronology of racial pain care disparities—from patients' and providers' attitudes and perceptions, through dyadic interaction, and finally, to outcomes—in natural clinical settings also enables us to identify novel, ecologically valid intervention points. In turn, it is our hope that interventions informed by this approach will be more feasible, scalable and effective, compared with interventions based on laboratory-based experimental work alone.

Identifying and addressing patient and provider factors that fuel racial pain care disparities is an important and ideal goal. However, this goal is unlikely to be effective on its own—or even attainable in the first place. This is because patient and provider factors (eg, patient mistrust, provider bias) reflect individuals' perceptions, attitudes and reactions that exist within and are shaped by broader societal systems that are carefully designed to maintain the status quo that marginalises racial and ethnic minorities.^{182–184} Thus, rather than reducing patient mistrust or provider bias, a viable alternative approach is to train providers to better manage their communication with patients. What makes our approach unique and different from the existing training and interventions focusing on patient-provider communication is, again, our emphasis on behavioural coordination. Current patient-provider communication training programmes are designed to improve providers' communication behaviours in isolation and fail to consider patients' contributions to these dyadic interactions.^{185–191} To reduce racial pain care disparities effectively, it is essential to identify exactly what triggers disruptions in coordination, at which behavioural level, by whom and in what sequence. Above all, it is critical to answer these questions within a naturalistic context. Once we identify specific aspects of behavioural coordination associated with racial disparities in pain care and outcomes, we can design interventions to help providers learn how to detect potential triggers of disruptions in coordination during their communication with patients and proactively avoid disruptions in coordination. Such interventions can also help providers learn how to increase or restore behavioural coordination with patients with diverse communication styles and practice those strategies. In sum, this work will triangulate on factors that provide the biggest return on investment for future interventions.

Limitations

First, one might anticipate that surgeons may try to modify how they interact with patients when being

recorded. However, our goal is to assess the strength of the associations between patient/provider factors and behavioural coordination, not the absolute levels of behavioural coordination. Therefore, as long as there is variability in providers' communication behaviours, this goal is still feasible. Indeed, in previous work using similar video recording procedures, we were able to capture variability in physicians' communication behaviours, which were systematically associated with implicit racial prejudice.^{20 178 179 192–194} There is no theoretical basis to expect that surgeons would differ in social desirability concerns from the physicians in our prior work. Our previous findings are also consistent with a large body of literature showing that bias in medical contexts manifests via automatic rather than deliberative processes.⁹⁵ Consequently, surgeons may be able to modify their verbal behaviours, but they are likely to have difficulty regulating their non-verbal and paraverbal behaviours.

Second, the proposed research focuses only on the planned surgical settings. From a methodological standpoint, focusing on preoperative consultations minimises inherent variability across care contexts (eg, surgical vs non-surgical care, acute vs chronic pain care, planned vs emergency care) and provides greater precision in testing our conceptual model. However, findings from the present study may be generalisable to patient–surgeon dyads in acute pain management in emergency settings, as well as a variety of patient-provider dyads (eg, primary care providers, specialists, integrative medicine providers, pain specialists, physical therapists) in chronic pain management settings. That said, these other contexts may be subject to the influence of other factors that are not present in the current work. After validating our conceptual model in the context of preoperative consultations, our future work will systematically examine whether this model is generalisable to other clinical contexts involving pain. Such empirical evaluation of the model's generalisability will further contribute to theory development. Relatedly, once we test and refine our conceptual model, we will assess its generalisability beyond the current institutional and geographical contexts.

Lastly, despite strong evidence of gender-based pain disparities, we are unable to examine the potential intersectional effects of race and gender in pain disparities due to the small surgeon sample size. Because the proposed study focuses on racial pain disparities, we will treat gender as a covariate rather than a second moderator in our hypothesis testing. This analytical approach aligns with prior research on racial pain disparities.^{12 14 16}

Author affiliations

¹Department of Public Health Sciences, University of Virginia, Charlottesville, Virginia, USA

²Department of Surgery, Virginia Commonwealth University, Richmond, Virginia, USA

³Department of Psychology, Davidson College, Davidson, North Carolina, USA

⁴Virginia Commonwealth University, Richmond, Virginia, USA

⁵Department of Pediatrics, Virginia Commonwealth University, Richmond, Virginia, USA

⁶Department of Psychology, Virginia Commonwealth University, Richmond, Virginia, USA

⁷Department of Psychology, University of Delaware, Newark, Delaware, USA

Acknowledgements We thank Gavin Mullin for his assistance with formatting in-text citations and references.

Contributors NHag and PM-S are the multiple principal investigators and ER, BAE, SCTJ and ACG are the co-investigators of the project. All authors have contributed to the development of this protocol. CE is the former project coordinator, and NHag is the current project coordinator. NHag wrote the first and final drafts of the protocol as well as revised the manuscript. PM-S, CE, ER, BAE and NHag contributed to the drafting and editing process. NHag is the guarantor. All authors have read, edited and approved the manuscript.

Funding This study has been funded by the National Institute of Health/National Institute of Nursing Research (R01 NR020030).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Nao Hagiwara <http://orcid.org/0000-0003-3933-8917>

REFERENCES

- Hunfeld JAM, Perquin CW, Duivenvoorden HJ. Chronic Pain and Its Impact on Quality of Life in Adolescents and Their Families. *J Pediatr Psychol* 2001;26:145–53.
- Katz N. The impact of pain management on quality of life. *J Pain Symptom Manage* 2002;24:S38–47.
- Niv D, Kreitler S. Pain and quality of life. *Pain Pract* 2001;1:150–61.
- Kennedy J, Roll JM, Schraudner T, et al. Prevalence of persistent pain in the U.S. adult population: new data from the 2010 national health interview survey. *J Pain* 2014;15:979–84.
- Gureje O. Comorbidity of pain and anxiety disorders. *Curr Psychiatry Rep* 2008;10:318–22.
- Gureje O. Treating chronic pain in the context of comorbid depression. *Pain* 2008;134:3–4.
- Lépine J-P, Briley M. The epidemiology of pain in depression. *Hum Psychopharmacol Clin Exp* 2004;19:S3–7.
- Gureje O, Von Korff M, Simon GE, et al. Persistent pain and well-being: a World Health Organization Study in Primary Care. *JAMA* 1998;280:147–51.
- Pizzi LT, Carter CT, Howell JB, et al. Work Loss, Healthcare Utilization, and Costs among US Employees with Chronic Pain. *Disease Management & Health Outcomes* 2005;13:201–8.
- Andrew R, Derry S, Taylor RS, et al. The costs and consequences of adequately managed chronic non-cancer pain and chronic neuropathic pain. *Pain Pract* 2014;14:79–94.
- Phillips CJ. Economic burden of chronic pain. *Expert Rev Pharmacoecon Outcomes Res* 2006;6:591–601.
- Anderson KO, Green CR, Payne R. Racial and ethnic disparities in pain: causes and consequences of unequal care. *J Pain* 2009;10:1187–204.
- Shavers VL, Bakos A, Sheppard VB. Race, ethnicity, and pain among the U.S. adult population. *J Health Care Poor Underserved* 2010;21:177–220.
- Lee P, Le Saux M, Siegel R, et al. Racial and ethnic disparities in the management of acute pain in US emergency departments: Meta-analysis and systematic review. *Am J Emerg Med* 2019;37:1770–7.
- Green CR, Anderson KO, Baker TA, et al. The unequal burden of pain: confronting racial and ethnic disparities in pain. *Pain Med* 2003;4:277–94.
- Mossey JM. Defining racial and ethnic disparities in pain management. *Clin Orthop Relat Res* 2011;469:1859–70.
- Penner LA, Dovidio JF, Hagiwara N, et al. An Analysis of Race-related Attitudes and Beliefs in Black Cancer Patients: Implications for Health Care Disparities. *J Health Care Poor Underserved* 2016;27:1503–20.
- Arnett MJ, Thorpe RJ, Gaskin DJ, et al. Race, Medical Mistrust, and Segregation in Primary Care as Usual Source of Care: Findings from the Exploring Health Disparities in Integrated Communities Study. *J Urban Health* 2016;93:456–67.
- Chen FM, Fryer GE Jr, Phillips RL Jr, et al. Patients' beliefs about racism, preferences for physician race, and satisfaction with care. *Ann Fam Med* 2005;3:138–43.
- Hagiwara N, Dovidio JF, Eggly S, et al. The effects of racial attitudes on affect and engagement in racially discordant medical interactions between non-Black physicians and Black patients. *Group Process Intergroup Relat* 2016;19:509–27.
- LaVeist TA, Isaac LA, Williams KP. Mistrust of health care organizations is associated with underutilization of health services. *Health Serv Res* 2009;44:2093–105.
- Thomson R, Murtagh M, Khaw FM. Tensions in public health policy: patient engagement, evidence-based public health and health inequalities. *Qual Saf Health Care* 2005;14:398–400.
- Williamson LD, Bigman CA. A systematic review of medical mistrust measures. *Patient Educ Couns* 2018;101:1786–94.
- Pascoe EA, Smart Richman L. Perceived discrimination and health: a meta-analytic review. *Psychol Bull* 2009;135:531–54.
- Mende-Siedlecki P, Qu-Lee J, Backer R, et al. Perceptual contributions to racial bias in pain recognition. *J Exp Psychol Gen* 2019;148:863–89.
- Green AR, Carney DR, Pallin DJ, et al. Implicit bias among physicians and its prediction of thrombolysis decisions for black and white patients. *J Gen Intern Med* 2007;22:1231–8.
- Hoffman KM, Trawalter S. Assumptions about life hardship and pain perception. *Group Process Intergroup Relat* 2016;19:493–508.
- Hoffman KM, Trawalter S, Axt JR, et al. Racial bias in pain assessment and treatment recommendations, and false beliefs about biological differences between blacks and whites. *Proc Natl Acad Sci USA* 2016;113:4296–301.
- Trawalter S, Hoffman KM, Waytz A. Racial bias in perceptions of others' pain. *PLoS ONE* 2012;7:e48546.
- Sabin JA, Greenwald AG. The influence of implicit bias on treatment recommendations for 4 common pediatric conditions: pain, urinary tract infection, attention deficit hyperactivity disorder, and asthma. *Am J Public Health* 2012;102:988–95.
- Mende-Siedlecki P, Rivet EB, Grover AC, et al. Making Meaningful Impacts: Centering Breakdowns in Dyadic Communication Processes in Racial Disparities in Surgical Pain Care. *Ann Surg* 2022;276:e646–8.
- Craig KD. The social communication model of pain. *Canadian Psy* 2009;50:22–32.
- Craig KD. Social communication model of pain. *Pain* 2015;156:1198–9.
- Duran N, Dale R, Richardson D, et al. The self-organization of human interaction. *Psychol Learn Motiv* 2013;59:43–95.
- Lakin JL. *Behavioral mimicry and interpersonal synchrony*. Berlin, Boston: DE GRUYTER, 2013:539–76.
- Richardson DC, Dale R, Kirkham NZ. The Art of Conversation Is Coordination. *Psychol Sci* 2007;18:407–13.
- Abney D, Paxton A, Dale R, et al. Cooperation in sound and motion: complexity matching in collaborative interaction. *Open Science Framework* [Preprint] 2021.
- Abney DH, Paxton A, Dale R, et al. Complexity matching in dyadic conversation. *J Exp Psychol Gen* 2014;143:2304–15.
- Schmidt RC, Fitzpatrick P. Embodied synchronization and complexity in a verbal interaction. *Nonlinear Dyn Psychol Life Sci* 2019;23:199–228.
- Vicaria IM, Dickens L. Meta-Analyses of the Intra- and Interpersonal Outcomes of Interpersonal Coordination. *J Nonverbal Behav* 2016;40:335–61.

- 41 Louwerse MM, Dale R, Bard EG, *et al.* Behavior matching in multimodal communication is synchronized. *Cogn Sci* 2012;36:1404–26.
- 42 Chemero A. Sensorimotor empathy. *J Conscious Stud* 2016;23:152.
- 43 Keller PE, Novembre G, Hove MJ. Rhythm in joint action: psychological and neurophysiological mechanisms for real-time interpersonal coordination. *Philos Trans R Soc Lond B Biol Sci* 2014;369:20130394.
- 44 Koehne S, Hatri A, Cacioppo JT, *et al.* Perceived interpersonal synchrony increases empathy: Insights from autism spectrum disorder. *Cognition* 2016;146:8–15.
- 45 Anderson SR, Gianola M, Perry JM, *et al.* Clinician-Patient Racial/Ethnic Concordance Influences Racial/Ethnic Minority Pain: Evidence from Simulated Clinical Interactions. *Pain Med* 2020;21:3109–25.
- 46 Moore C, Coates E, Watson A, *et al.* 'It's Important to Work with People that Look Like Me': Black Patients' Preferences for Patient-Provider Race Concordance. *J Racial and Ethnic Health Disparities* 2023;10:2552–64.
- 47 Nazione S, Perrault EK, Keating DM. Finding Common Ground: Can Provider-Patient Race Concordance and Self-disclosure Bolster Patient Trust, Perceptions, and Intentions? *J Racial Ethn Health Disparities* 2019;6:962–72.
- 48 Schoenthaler A, Allegante JP, Chaplin W, *et al.* The effect of patient-provider communication on medication adherence in hypertensive black patients: does race concordance matter? *Ann Behav Med* 2012;43:372–82.
- 49 Street RL Jr, O'Malley KJ, Cooper LA, *et al.* Understanding concordance in patient-physician relationships: personal and ethnic dimensions of shared identity. *Ann Fam Med* 2008;6:198–205.
- 50 Tunçgenç B, Cohen E. Movement Synchrony Forges Social Bonds across Group Divides. *Front Psychol* 2016;7:782.
- 51 Goldstein P, Losin EAR, Anderson SR, *et al.* Clinician-Patient Movement Synchrony Mediates Social Group Effects on Interpersonal Trust and Perceived Pain. *J Pain* 2020;21:1160–74.
- 52 Brown J. The battle to become a Black physician in the USA: past and present. *The Lancet* 2024;403:426–8.
- 53 Ly DP. Historical Trends in the Representativeness and Incomes of Black Physicians, 1900–2018. *J Gen Intern Med* 2022;37:1310–2.
- 54 Mora H, Obayemi A, Holcomb K, *et al.* The National Deficit of Black and Hispanic Physicians in the US and Projected Estimates of Time to Correction. *JAMA Netw Open* 2022;5:e2215485.
- 55 Fiscella K, Sanders MR. Racial and Ethnic Disparities in the Quality of Health Care. *Annu Rev Public Health* 2016;37:375–94.
- 56 Akinbami LJ, Moorman JE, Simon AE, *et al.* Trends in racial disparities for asthma outcomes among children 0 to 17 years, 2001–2010. *J Allergy Clin Immunol* 2014;134:547–53.
- 57 Arthur M, Hedges JR, Newgard CD, *et al.* Racial disparities in mortality among adults hospitalized after injury. *Med Care* 2008;46:192–9.
- 58 Duru OK, Harawa NT, Kermah D, *et al.* Allostatic load burden and racial disparities in mortality. *J Natl Med Assoc* 2012;104:89–95.
- 59 Haider AH, Scott VK, Rehman KA, *et al.* Racial disparities in surgical care and outcomes in the United States: a comprehensive review of patient, provider, and systemic factors. *J Am Coll Surg* 2013;216:482–92.
- 60 O'Keefe EB, Meltzer JP, Bethea TN. Health disparities and cancer: racial disparities in cancer mortality in the United States, 2000–2010. *Front Public Health* 2015;3:51.
- 61 Ng B, Dimsdale JE, Rollnik JD, *et al.* The effect of ethnicity on prescriptions for patient-controlled analgesia for post-operative pain. *Pain* 1996;66:9–12.
- 62 Beletsky A, Burton BN, Finneran IV JJ, *et al.* Association of race and ethnicity in the receipt of regional anesthesia following mastectomy. *Reg Anesth Pain Med* 2021;46:118–23.
- 63 Memtsoudis SG, Besculides MC, Swamidoss CP. Do race, gender, and source of payment impact on anesthetic technique for inguinal hernia repair? *J Clin Anesth* 2006;18:328–33.
- 64 Faucett J, Gordon N, Levine J. Differences in postoperative pain severity among four ethnic groups. *J Pain Symptom Manage* 1994;9:383–9.
- 65 Memtsoudis SG, Poeran J, Zubizarreta N, *et al.* Anesthetic Care for Orthopedic Patients: Is There a Potential for Differences in Care? *Anesthesiology* 2016;124:608–23.
- 66 Goodman SM, Mandl LA, Parks ML, *et al.* Disparities in TKA Outcomes: Census Tract Data Show Interactions Between Race and Poverty. *Clin Orthop Relat Res* 2016;474:1986–95.
- 67 Goodman SM, Mehta B, Zhang M, *et al.* Disparities in Total Hip Arthroplasty Outcomes: Census Tract Data Show Interactions Between Race and Community Deprivation. *J Am Acad Orthop Surg* 2018;26:e457–64.
- 68 Lavernia CJ, Alcerro JC, Contreras JS, *et al.* Ethnic and racial factors influencing well-being, perceived pain, and physical function after primary total joint arthroplasty. *Clin Orthop Relat Res* 2011;469:1838–45.
- 69 McDonald DD. Gender and ethnic stereotyping and narcotic analgesic administration. *Res Nurs Health* 1994;17:45–9.
- 70 Khor S, Lavalley D, Cizik AM, *et al.* Development and Validation of a Prediction Model for Pain and Functional Outcomes After Lumbar Spine Surgery. *JAMA Surg* 2018;153:634.
- 71 Hall MJ, Schwartzman A, Zhang J, *et al.* Ambulatory Surgery Data From Hospitals and Ambulatory Surgery Centers: United States, 2010. *Natl Health Stat Report* 2017;1:1–15.
- 72 Frantsve LME, Kerns RD. Patient-provider interactions in the management of chronic pain: current findings within the context of shared medical decision making. *Pain Med* 2007;8:25–35.
- 73 Jimenez N, Garrouette E, Kundu A, *et al.* A review of the experience, epidemiology, and management of pain among American Indian, Alaska Native, and Aboriginal Canadian peoples. *J Pain* 2011;12:511–22.
- 74 Israel JS, Poore SO. The Clinical Conundrum of Perioperative Pain Management in Patients with Opioid Dependence. *Plast Reconstr Surg (1946)* 2013;131:657e–8e.
- 75 Ljungqvist O, Scott M, Fearon KC. Enhanced Recovery After Surgery: A Review. *JAMA Surg* 2017;152:292–8.
- 76 Theunissen M, Peters ML, Bruce J, *et al.* Preoperative anxiety and catastrophizing: a systematic review and meta-analysis of the association with chronic postsurgical pain. *Clin J Pain* 2012;28:819–41.
- 77 Mazer LS, Lovett PJ, Miller JM, *et al.* Improving satisfaction with postoperative pain management for patients with persistent pain: a preoperative pain medicine consultation intervention. *J Perioper Pract* 2020;30:345–51.
- 78 Smith RJ, Rhodes K, Paciotti B, *et al.* Patient Perspectives of Acute Pain Management in the Era of the Opioid Epidemic. *Ann Emerg Med* 2015;66:246–52.
- 79 Smedley BD, Stith AY, Nelson AR, eds. Institute of medicine committee on understanding and eliminating racial and ethnic disparities in health care. In: *Unequal treatment: confronting racial and ethnic disparities in health care*. Washington DC: National Academies Press, 2003.
- 80 Mayo clinic surgical outcomes program recommendations for adult discharge opioid prescriptions. 2018. Available: <http://links.lww.com/SLA/B477> [Accessed 27 Aug 2020].
- 81 Lin J, Drain A, Goharad A, *et al.* What factors predict anti-Black bias in pain perception? An internal meta-analysis across 40 experimental studies. *Social & Personality Psych* 2024;18:e12901.
- 82 Mende-Siedlecki P, Lin J, Ferron S, *et al.* Seeing no pain: Assessing the generalizability of racial bias in pain perception. *Emotion* 2021;21:932–50.
- 83 Greenwald AG, McGhee DE, Schwartz JKL. Measuring individual differences in implicit cognition: the implicit association test. *J Pers Soc Psychol* 1998;74:1464–80.
- 84 Greenwald AG, Nosek BA, Banaji MR. Understanding and using the implicit association test: I. An improved scoring algorithm. *J Pers Soc Psychol* 2003;85:197–216.
- 85 Greenwald AG, Poehlman TA, Uhlmann EL, *et al.* Understanding and using the Implicit Association Test: III. Meta-analysis of predictive validity. *J Pers Soc Psychol* 2009;97:17–41.
- 86 Cunningham WA, Preacher KJ, Banaji MR. Implicit attitude measures: consistency, stability, and convergent validity. *Psychol Sci* 2001;12:163–70.
- 87 Lane KA, Banaji MR, Nosek BA, *et al.* Understanding and using the implicit association test: iv. what we know (so far). In: Wittenbrink B, Schwarz NS, eds. *Implicit measures of attitudes: processes and controversies*. New York, NY: Guilford Press, 2007.
- 88 Nosek BA, Greenwald AG, Banaji MR. Understanding and using the Implicit Association Test: II. Method variables and construct validity. *Pers Soc Psychol Bull* 2005;31:166–80.
- 89 Campbell DT. *White attitudes toward black people*. Ann Arbor, MI: Institute for Social Research, 1971.
- 90 Axt JR. The Best Way to Measure Explicit Racial Attitudes Is to Ask About Them. *Soc Psychol Personal Sci* 2018;9:896–906.
- 91 Henry PJ, Sears DO. The Symbolic Racism 2000 Scale. *Polit Psychol* 2002;23:253–83.
- 92 Tarman C, Sears DO. The Conceptualization and Measurement of Symbolic Racism. *J Polit* 2005;67:731–61.
- 93 Sears DO, Henry PJ. The origins of symbolic racism. *J Pers Soc Psychol* 2003;85:259–75.
- 94 Sears DO, Henry PJ. Over thirty years later: A contemporary look at symbolic racism. *Adv Exp Soc Psychol* 2005;37:95–125.

- 95 Hagiwara N, Dovidio JF, Stone J, *et al.* Applied Racial/Ethnic Healthcare Disparities Research Using Implicit Measures. *Soc Cogn* 2020;38:s68–97.
- 96 Sabin J, Nosek BA, Greenwald A, *et al.* Physicians' implicit and explicit attitudes about race by MD race, ethnicity, and gender. *J Health Care Poor Underserved* 2009;20:896–913.
- 97 Oliver MN, Wells KM, Joy-Gaba JA, *et al.* Do physicians' implicit views of African Americans affect clinical decision making? *J Am Board Fam Med* 2014;27:177–88.
- 98 Hagiwara N, Duffy C, Quillin J. Implicit and explicit racial prejudice and stereotyping toward Black (vs. White) Americans: The prevalence and variation among genetic counselors in North America. *J Genet Couns* 2023;32:397–410.
- 99 Wittenbrink B, Judd CM, Park B. Evidence for racial prejudice at the implicit level and its relationship with questionnaire measures. *J Pers Soc Psychol* 1997;72:262–74.
- 100 Goff PA, Steele CM, Davies PG. The space between us: stereotype threat and distance in interracial contexts. *J Pers Soc Psychol* 2008;94:91–107.
- 101 Brondolo E, Kelly KP, Coakley V, *et al.* The Perceived Ethnic Discrimination Questionnaire: Development and Preliminary Validation of a Community Version1. *J Appl Social Psychol* 2005;35:335–65.
- 102 Hagiwara N, Alderson CJ, McCauley JM. 'We get what we deserve': the belief in a just world and its health consequences for Blacks. *J Behav Med* 2015;38:912–21.
- 103 Hagiwara N, Alderson CJ, Mezek B. Differential Effects of Personal-Level vs Group-Level Racial Discrimination on Health among Black Americans. *Ethn Dis* 2016;26:453–60.
- 104 Hall MA, Camacho F, Dugan E, *et al.* Trust in the medical profession: conceptual and measurement issues. *Health Serv Res* 2002;37:1419–39.
- 105 Hall MA, Zheng B, Dugan E, *et al.* Measuring patients' trust in their primary care providers. *Med Care Res Rev* 2002;59:293–318.
- 106 Marshall GN, Hays RD. The patient satisfaction questionnaire short-form (psq-18). Santa Monica, CA Rand; 1994.
- 107 Azimi P, Benzel EC. Cut-Off Value for Pain Sensitivity Questionnaire in Predicting Surgical Success in Patients with Lumbar Disc Herniation. *PLoS ONE* 2016;11:e0160541.
- 108 Bjørnnes AK, Lie I, Parry M, *et al.* Association between self-perceived pain sensitivity and pain intensity after cardiac surgery. *J Pain Res* 2018;11:1425–32.
- 109 Valeberg BT, Høvik LH, Gjelle KH. Relationship between self-reported pain sensitivity and pain after total knee arthroplasty: a prospective study of 71 patients 8 weeks after a standardized fast-track program. *J Pain Res* 2016;9:625–9.
- 110 Rehberg B, Mathivon S, Combescure C, *et al.* Prediction of Acute Postoperative Pain Following Breast Cancer Surgery Using the Pain Sensitivity Questionnaire. *Clin J Pain* 2017;33:57–66.
- 111 Kim H-J, Lee J-I, Kang K-T, *et al.* Influence of Pain Sensitivity on Surgical Outcomes After Lumbar Spine Surgery in Patients With Lumbar Spinal Stenosis. *Spine (Phila Pa 1986)* 2015;40:193–200.
- 112 Kim H-J, Park J-H, Kim J-W, *et al.* Prediction of Postoperative Pain Intensity after Lumbar Spinal Surgery Using Pain Sensitivity and Preoperative Back Pain Severity. *Pain Med* 2014;15:2037–45.
- 113 Tuna T, Boz S, Van Obbergh L, *et al.* Comparison of the Pain Sensitivity Questionnaire and the Pain Catastrophizing Scale in Predicting Postoperative Pain and Pain Chronicization After Spine Surgery. *Clin Spine Surg* 2018;31:E432–40.
- 114 Duchow J, Schlörcke E, Hüppe M. Self-rated pain sensitivity and postoperative pain. *Schmerz* 2013;27:371–9.
- 115 Sommer M, de Rijke JM, van Kleef M, *et al.* Predictors of acute postoperative pain after elective surgery. *Clin J Pain* 2010;26:87–94.
- 116 Turk DC, Wilson HD. Fear of pain as a prognostic factor in chronic pain: conceptual models, assessment, and treatment implications. *Curr Pain Headache Rep* 2010;14:88–95.
- 117 Granot M, Ferber SG. The Roles of Pain Catastrophizing and Anxiety in the Prediction of Postoperative Pain Intensity. *Clin J Pain* 2005;21:439–45.
- 118 McIntyre MH, Kless A, Hein P, *et al.* Validity of the cold pressor test and pain sensitivity questionnaire via online self-administration. *PLoS ONE* 2020;15:e0231697.
- 119 Ruscheweyh R, Marziniak M, Stumpfenhorst F, *et al.* Pain sensitivity can be assessed by self-rating: Development and validation of the Pain Sensitivity Questionnaire. *Pain* 2009;146:65–74.
- 120 Ruscheweyh R, Verneuer B, Dany K, *et al.* Validation of the Pain Sensitivity Questionnaire in chronic pain patients. *PAIN* 2012;153:1210–8.
- 121 Sellers AB, Ruscheweyh R, Kelley BJ, *et al.* Validation of the English language pain sensitivity questionnaire. *Reg Anesth Pain Med* 2013;38:508–14.
- 122 McCracken LM, Dhinra L. A short version of the Pain Anxiety Symptoms Scale (PASS-20): preliminary development and validity. *Pain Res Manag* 2002;7:45–50.
- 123 Roelofs J, McCracken L, Peters ML, *et al.* Psychometric evaluation of the Pain Anxiety Symptoms Scale (PASS) in chronic pain patients. *J Behav Med* 2004;27:167–83.
- 124 Abrams MP, Carleton RN, Asmundson GJG. An exploration of the psychometric properties of the PASS-20 with a nonclinical sample. *J Pain* 2007;8:879–86.
- 125 Carleton RN, Asmundson GJG. The multidimensionality of fear of pain: construct independence for the fear of Pain Questionnaire-Short Form and the Pain Anxiety Symptoms Scale-20. *J Pain* 2009;10:29–37.
- 126 Coons MJ, Hadjistavropoulos HD, Asmundson GJG. Factor structure and psychometric properties of the Pain Anxiety Symptoms Scale-20 in a community physiotherapy clinic sample. *Eur J Pain* 2004;8:511–6.
- 127 Meredith P, Strong J, Feeney JA. Adult attachment, anxiety, and pain self-efficacy as predictors of pain intensity and disability. *Pain* 2006;123:146–54.
- 128 Asghari A, Nicholas MK. Pain self-efficacy beliefs and pain behaviour. A prospective study. *Pain* 2001;94:85–100.
- 129 Turner JA, Ersek M, Kemp C. Self-efficacy for managing pain is associated with disability, depression, and pain coping among retirement community residents with chronic pain. *J Pain* 2005;6:471–9.
- 130 Porter LS, Keefe FJ, Garst J, *et al.* Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: associations with symptoms and distress. *Pain* 2008;137:306–15.
- 131 Somers TJ, Wren AA, Shelby RA. The context of pain in arthritis: self-efficacy for managing pain and other symptoms. *Curr Pain Headache Rep* 2012;16:502–8.
- 132 Brister H, Turner JA, Aaron LA, *et al.* Self-efficacy is associated with pain, functioning, and coping in patients with chronic temporomandibular disorder pain. *J Orofac Pain* 2006;20:115–24.
- 133 Nicholas MK. The pain self-efficacy questionnaire: Taking pain into account. *Eur J Pain* 2007;11:153–63.
- 134 Costa DSJ, Asghari A, Nicholas MK. Item response theory analysis of the Pain Self-Efficacy Questionnaire. *Scand J Pain* 2017;14:113–7.
- 135 Di Pietro F, Catley MJ, McAuley JH, *et al.* Rasch analysis supports the use of the Pain Self-Efficacy Questionnaire. *Phys Ther* 2014;94:91–100.
- 136 McWilliams LA, Kowal J, Wilson KG. Development and evaluation of short forms of the Pain Catastrophizing Scale and the Pain Self-efficacy Questionnaire. *Eur J Pain* 2015;19:1342–9.
- 137 Chiarotto A, Falla D, Polli A, *et al.* Validity and Responsiveness of the Pain Self-Efficacy Questionnaire in Patients With Neck Pain Disorders. *J Orthop Sports Phys Ther* 2018;48:204–16.
- 138 Bijur PE, Latimer CT, Gallagher EJ. Validation of a Verbally Administered Numerical Rating Scale of Acute Pain for Use in the Emergency Department. *Acad Emerg Med* 2003;10:390–2.
- 139 Hollen PJ, Gralla RJ, Kris MG, *et al.* A comparison of visual analogue and numerical rating scale formats for the Lung Cancer Symptom Scale (LCSS): Does format affect patient ratings of symptoms and quality of life? *Qual Life Res* 2005;14:837–47.
- 140 Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain* 1986;27:117–26.
- 141 Williamson A, Hoggart B. Pain: a review of three commonly used pain rating scales. *J Clin Nurs* 2005;14:798–804.
- 142 Ware JE, Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) Project. *J Clin Epidemiol* 1998;51:903–12.
- 143 Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE Open Med* 2016;4:2050312116671725.
- 144 Elliott TE, Renier CM, Palcher JA. Chronic pain, depression, and quality of life: correlations and predictive value of the SF-36. *Pain Med* 2003;4:331–9.
- 145 Meyer-Rosberg K, Burckhardt CS, Huizar K, *et al.* A comparison of the SF-36 and Nottingham Health Profile in patients with chronic neuropathic pain. *Eur J Pain* 2001;5:391–403.
- 146 Bronfort G, Bouter LM. Responsiveness of general health status in chronic low back pain: a comparison of the COOP charts and the SF-36. *Pain* 1999;83:201–9.
- 147 Razykov I, Ziegelstein RC, Whooley MA, *et al.* The PHQ-9 versus the PHQ-8--is item 9 useful for assessing suicide risk in coronary artery disease patients? Data from the Heart and Soul Study. *J Psychosom Res* 2012;73:163–8.

- 148 Kroenke K, Spitzer RL. The PHQ-9: A New Depression Diagnostic and Severity Measure. *Psychiatr Ann* 2002;32:509–15.
- 149 Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–13.
- 150 Wu Y, Levis B, Riehm KE, et al. Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. *Psychol Med* 2020;50:1368–80.
- 151 Spielberger CD. Manual for the state-trait anxiety inventory (self-evaluation questionnaire). 1970.
- 152 Vorauer JD, Hunter AJ, Main KJ, et al. Meta-stereotype activation: evidence from indirect measures for specific evaluative concerns experienced by members of dominant groups in intergroup interaction. *J Pers Soc Psychol* 2000;78:690–707.
- 153 Williams MT, Chapman LK, Wong J, et al. The role of ethnic identity in symptoms of anxiety and depression in African Americans. *Psychiatry Res* 2012;199:31–6.
- 154 Jotwani R, Turnbull ZA, White RS. The economic cost of racial disparities in perioperative care. *J Comp Eff Res* 2020;9:317–20.
- 155 Cintron A, Morrison RS. Pain and Ethnicity in the United States: A Systematic Review. *J Palliat Med* 2006;9:1454–73.
- 156 Webber CL Jr, Zbilut JP. Dynamical assessment of physiological systems and states using recurrence plot strategies. *J Appl Physiol* 1994;76:965–73.
- 157 Marwan N, Carmenromano M, Thiel M, et al. Recurrence plots for the analysis of complex systems. *Phys Rep* 2007;438:237–329.
- 158 Paxton A, Dale R. Frame-differencing methods for measuring bodily synchrony in conversation. *Behav Res* 2013;45:329–43.
- 159 Coco MI, Dale R. Cross-recurrence quantification analysis of categorical and continuous time series: an R package. *Front Psychol* 2014;5:510.
- 160 Coco M, Mønster D, Leonardi G, et al. n.d. Unidimensional and Multidimensional Methods for Recurrence Quantification Analysis with crqa. *R J* 13:145.
- 161 Wallot S. Multidimensional Cross-Recurrence Quantification Analysis (MdCrQA) – A Method for Quantifying Correlation between Multivariate Time-Series. *Multivariate Behav Res* 2019;54:173–91.
- 162 Angus D. Recurrence Methods for Communication Data, Reflecting on 20 Years of Progress. *Front Appl Math Stat* 2019;5:54.
- 163 Miles K, Weisser A, Kallen RW, et al. Behavioral dynamics of conversation, (mis)communication and coordination in noisy environments. *Sci Rep* 2023;13:20271.
- 164 Tolston M, Ariyabuddhipongs K, Riley MA, et al. Cross-recurrence quantification analysis of the influence of coupling constraints on interpersonal coordination and communication. In: *Translational recurrences: from mathematical theory to real-world applications* 2014. Springer International Publishing, 2014: 157–71.
- 165 Ambady N, Bernieri FJ, Richeson JA, et al, eds. Toward a histology of social behavior: judgmental accuracy from thin slices of the behavioral stream. In: *Advances in experimental social psychology*. New York: Academic Press, 2000.:32.
- 166 Ambady N, Rosenthal R. Thin slices of expressive behavior as predictors of interpersonal consequences: A meta-analysis. *Psychol Bull* 1992;111:256–74.
- 167 Ambady N. The Perils of Pondering: Intuition and Thin Slice Judgments. *Psychol Inq* 2010;21:271–8.
- 168 Henry SG, Eggly S. The effect of discussing pain on patient-physician communication in a low-income, black, primary care patient population. *J Pain* 2013;14:759–66.
- 169 Henry SG, Penner LA, Eggly S. Associations between thin slice ratings of affect and rapport and perceived patient-centeredness in primary care: Comparison of audio and video recordings. *Patient Educ Couns* 2017;100:1128–35.
- 170 Auld F, White AM. Rules for Dividing Interviews Into Sentences. *J Psychol* 1956;42:273–81.
- 171 Hatfield JD, Weider-Hatfield D. The comparative utility of three types of behavioral units for interaction analysis. *Commun Monogr* 1978;45:44–50.
- 172 Murray EJ. A content-analysis method for studying psychotherapy. *Psychological Monographs: General and Applied* 1956;70:1–31.
- 173 Rao JK, Anderson LA, Inui TS, et al. Communication Interventions Make A Difference in Conversations Between Physicians and Patients. *Med Care* 2007;45:340–9.
- 174 Roter DL, Hall JA, Katz NR. Relations Between Physicians?? Behaviors and Analogue Patients?? Satisfaction, Recall, and Impressions. *Med Care* 1987;25:437–51.
- 175 Salmon P, Ring A, Dowrick CF, et al. What do general practice patients want when they present medically unexplained symptoms, and why do their doctors feel pressurized? *J Psychosom Res* 2005;59:255–60.
- 176 Schirmer JM, Mauksch L, Lang F, et al. Assessing communication competence: a review of current tools. *Fam Med* 2005;37:184–92.
- 177 Zoppi K, Epstein RM. Is communication a skill? Communication behaviors and being in relation. *Fam Med* 2002;34:319–24.
- 178 Hagiwara N, Penner LA, Gonzalez R, et al. Racial attitudes, physician-patient talk time ratio, and adherence in racially discordant medical interactions. *Soc Sci Med* 2013;87:123–31.
- 179 Hagiwara N, Slatcher RB, Eggly S, et al. Physician Racial Bias and Word Use during Racially Discordant Medical Interactions. *Health Commun* 2017;32:401–8.
- 180 Estimation of the Medians for Dependent Variables. *Ann Math Statist* 1959;30:192–7.
- 181 Dunn OJ. Multiple Comparisons among Means. *J Am Stat Assoc* 1961;56:52–64.
- 182 Kraus MW, Callaghan B. Noblesse oblige? Social status and economic inequality maintenance among politicians. *PLoS ONE* 2014;9:e85293.
- 183 Maxwell A, Shields T. The Fate of Obamacare: Racial Resentment, Ethnocentrism and Attitudes about Healthcare Reform. *Race Soc Probl* 2014;6:293–304.
- 184 Wilson WJ. Toward a Framework for Understanding Forces that Contribute to or Reinforce Racial Inequality. *Race Soc Probl* 2009;1:3–11.
- 185 Makoul G. MSJAMA. Communication skills education in medical school and beyond. *JAMA* 2003;289:93.
- 186 How to improve the medical students' communication skills. *Med Teach* 2008;30:639–41.
- 187 Morgado P, Lemos AR, Almeida S, et al. A structured remediation program for communication skills. *Int J Med Educ* 2019;10:161–2.
- 188 Drazen JM, Shields HM, Loscalzo J. A Division of Medical Communications in an Academic Medical Center's Department of Medicine. *Acad Med* 2014;89:1623–9.
- 189 Street RL Jr, De Haes H. Designing a curriculum for communication skills training from a theory and evidence-based perspective. *Patient Educ Couns* 2013;93:27–33.
- 190 Al Odhayani A, Ratnapalan S. Teaching communication skills. *Can Fam Physician* 2011;57:1216–8.
- 191 Feraco AM, Brand SR, Mack JW, et al. Communication Skills Training in Pediatric Oncology: Moving Beyond Role Modeling. *Pediatric Blood & Cancer* 2016;63:966–72.
- 192 Penner LA, Dovidio JF, Gonzalez R, et al. The Effects of Oncologist Implicit Racial Bias in Racially Discordant Oncology Interactions. *J Clin Oncol* 2016;34:2874–80.
- 193 Hagiwara N, Mezuk B, Elston Lafata J, et al. Study protocol for investigating physician communication behaviours that link physician implicit racial bias and patient outcomes in Black patients with type 2 diabetes using an exploratory sequential mixed methods design. *BMJ Open* 2018;8:e022623.
- 194 Hagiwara N, Harika N, Carmany EP, et al. Racial disparities in cancer genetic counseling encounters: study protocol for investigating patient-genetic counselor communication in the naturalistic clinical setting using a convergent mixed methods design. *BMC Cancer* 2023;23:983.