

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

Effect of Pasta in the Context of Low Glycemic Index Dietary Patterns on Body Weight and Markers of Global Adiposity: A Systematic Review and Meta-analysis of Randomized Controlled Trials in Adults

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019438
Article Type:	Research
Date Submitted by the Author:	31-Aug-2017
Complete List of Authors:	Chiavaroli, Laura; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit; University of Toronto, Nutritional Sciences Kendall, Cyril; University of Toronto, Nutritional Sciences Braunstein, Catherine; University of Toronto, Nutritional Sciences; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit Blanco Mejia, Sonia; Clinical Nutrition and Risk Factor Modification Centre, St. Michael's Hospital; Toronto 3D Knowledge Synthesis and Clinical Trials Unit, St. Michael's Hospital Leiter, Lawrence; St. Michael's Hospital, Endocrinology; University of Toronto, Nutritional Sciences Jenkins, David; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Center; University of Toronto, Nutritional Sciences Sievenpiper, John; McMaster University, Department of Pathology and Molecular Medicine; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Centre
Primary Subject Heading:	Nutrition and metabolism
Secondary Subject Heading:	Cardiovascular medicine
Keywords:	body weight, pasta, glycemic index, glycaemic index, systematic review and meta-analysis, weight loss

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Effect of Pasta in the Context of Low Glycemic Index Dietary Patterns on Body Weight
2 and Markers of Global Adiposity: A Systematic Review and Meta-analysis of Randomized
3 Controlled Trials in Adults

4 Laura Chiavaroli^{1,2}, Cyril WC Kendall^{1,2,3}, Catherine R Braunstein^{1,2}, Sonia Blanco Mejia^{1,2},
5 Lawrence A Leiter^{1,2,4-6}, David JA Jenkins^{1,2,4-6}, John L Sievenpiper^{1,2,4-6}

6 1 Department of Nutritional Sciences, University of Toronto, Toronto, Canada

7 2 Toronto 3D Knowledge Synthesis and Clinical Trials Unit, St. Michael’s Hospital, Toronto,
8 Canada

9 3 College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, Canada

10 4 Department of Medicine, University of Toronto, Toronto, Canada

11 5 Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Toronto, Canada

12 6 Division of Endocrinology and Metabolism, St. Michael’s Hospital, Toronto, Canada

14 Corresponding Author:

15 Dr. John L Sievenpiper, MD, PhD, FRCPC, St. Michael’s Hospital, #6137-61 Queen Street East,
16 Toronto, ON, M5C 2T2, CANADA

17 Telephone: +1 416 867 7475; Fax: +1 416 867 7495; Email: john.sievenpiper@utoronto.ca

19 Word Count: 4,816

20 Tables: 1

21 Figures: 3

22 Online only supplementary material included

24 **Keywords:** body weight, pasta, glycemic index, glycaemic index, systematic review and meta-
25 analysis, weight loss

ABSTRACT

Objective: Carbohydrates have been implicated in the epidemic of obesity. To assess the effect of pasta alone or pasta in the context of low glycemic index (GI) dietary patterns on adiposity, we conducted a systematic review and meta-analysis.

Design: Systematic review and meta-analysis of randomized controlled trials (RCTs) with GRADE assessment.

Eligibility criteria for selecting studies: MEDLINE, Embase, CINAHL, and the Cochrane Library were searched through 07 February 2017. We included RCTs of ≥ 3 -weeks assessing the effect of pasta alone or pasta in the context of low-GI dietary patterns on measures of global (body weight, BMI, body fat) and regional (waist circumference [WC], waist-to-hip ratio [WHR], sagittal abdominal diameter [SAD]) adiposity in adults. Two independent reviewers extracted data and assessed risk of bias. Data were pooled using the generic inverse-variance method and expressed as mean differences (MDs) with 95% confidence intervals (95% CIs). Heterogeneity was assessed (Cochran Q statistic) and quantified (I²-statistic). GRADE assessed the overall quality of the evidence.

Results: We identified no trials of the effect of pasta alone and 32 trial comparisons (n=2,448 participants) of the effect of pasta in the context of low-GI dietary patterns. Pasta in the context of low-GI dietary patterns significantly reduced body weight (MD=-0.63kg [95% CIs, -0.84, -0.42kg]) and BMI (MD=-0.26kg/m² [95% CIs, -0.36, -0.16kg/m²]) compared with higher-GI dietary patterns. There was no effect on other measures of adiposity. The evidence was graded as

1
2
3 52 moderate for body weight, BMI, WHR, and SAD owing to a downgrade for indirectness and low
4
5 53 for WC and body fat owing to downgrades for indirectness and inconsistency.
6
7
8 54

9
10 55 **Conclusions:** The available RCTs demonstrate that pasta in the context of low-GI dietary
11
12 56 patterns does not adversely affect adiposity. Future trials should assess the effect of pasta in the
13
14 57 context of other healthy dietary patterns.
15
16
17 58

18
19 59 **Protocol registration:** ClinicalTrials.gov Identifier, NCT02961088
20
21
22 60
23
24 61

25
26 62 **Strengths and limitations of this study**

- 27
28 63 - The present systematic review and meta-analysis was based on a comprehensive search
29
30 64 and include a large number of randomized controlled trials which provide the best
31
32 65 protection against bias
33
34 66 - We used the Grading of Recommendations Assessment, Development, and Evaluation
35
36 67 (GRADE) system to evaluate the strength and quality of the evidence.
37
38 68 - There was evidence of unexplained inconsistency in the treatment estimates across trials
39
40 69 for waist circumference and body fat.
41
42 70 - The generalizability of our results is questionable with evidence of indirectness in the
43
44 71 pooled estimates for all body and adiposity outcomes, as all of the trials assessed pasta in
45
46 72 the context of low-GI dietary patterns (no trials assessed the effect of pasta alone or in the
47
48 73 context of other dietary patterns) and most of the available trials did not quantify the
49
50 74 amount of pasta consumed.
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

As the role of saturated fat in chronic disease has been called into question, carbohydrates have come under attack in the media^{1,2}, popular books³⁻⁹, statements of health advocacy groups¹⁰ and commentaries in leading medical journals^{11,12}. Much of the attention has focused on sugars but traditional carbohydrate staples like pasta, rice, and breads are increasingly being implicated in the epidemics of overweight and obesity^{1,7}. Although advantages for weight related outcomes have been shown for dietary patterns that are high in these foods but low in glycemic index (GI)^{13,14}, high in whole grains^{15,16}, and/or high in dietary fibre^{17,18}, there has been a general lack of recognition of the importance of carbohydrate quality.

Pasta is an important example of a food which is considered a refined carbohydrate but has a low-GI, a property that has been exploited extensively in studies of low-GI dietary patterns. It remains unclear whether pasta alone or in the context of a low-GI dietary pattern contributes to weight gain. We undertook a systematic review and meta-analysis of randomized controlled trials (RCTs) with a full GRADE assessment to quantify the effect of pasta alone or in the context of low-GI dietary patterns on body weight and measures of adiposity relevant to the prevention and management of overweight and obesity.

METHODS

Design

Our protocol followed the guideline of the Cochrane Handbook for Systematic Reviews of Interventions¹⁹, and findings are reported according to the guidelines of the Preferred Reporting

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Items for Systematic Reviews and Meta-Analyses²⁰. The protocol is registered at
clinicaltrials.gov (identifier, NCT02961088).

Data sources and searches

We searched MEDLINE (<http://www.nlm.nih.gov/bsd/pmresources.html>), Embase
(<https://www.embase.com>), CINAHL (<https://health.ebsco.com/products/the-cinahl-database>),
and the Cochrane Library (<http://www.cochranelibrary.com/>) through 07 February 2017. The full
search terms used in this study are presented in **Supplemental Tables S1-S2**. Briefly, we
searched using variations of the terms pasta and glycemic index and glycemic load and body
weight and BMI. The search was limited to human studies and had no language restrictions.
Reference lists of selected studies and reviews were also searched to identify additional articles.

Study selection

We include RCTs that investigated the effect of pasta consumed alone or in the context of low-
GI dietary patterns that emphasized pasta in comparison with higher GI diets that did not include
pasta on body weight or other measures of global (BMI, body fat) or abdominal (waist
circumference, waist-to-hip ratio, sagittal abdominal diameter, or visceral adipose tissue assessed
by imaging modalities) adiposity in participants of all health backgrounds. Trials were excluded
if they had diet duration of <3 weeks, did not intend to use a calorie- and macronutrient-matched
comparator arm that was higher in GI; included pregnant or breast-feeding women or children; or
did not provide suitable end-point data. When multiple publications existed for the same study,
the article with the most information was included (n=6). Published abstracts were not included.

120 Data extraction

121 Two reviewers (LC, CB) assessed the titles and abstracts of all identified studies and
122 independently reviewed and extracted relevant data from each report, including study design,
123 blinding, sample size, participant characteristics, follow-up duration, identification of pasta in the
124 low-GI diet only, comparator diet, macronutrient profile, funding source, and outcome data. The
125 primary outcome was body weight, and secondary outcomes included markers of global (BMI,
126 body fat) and abdominal (waist circumference, waist-to-hip ratio, sagittal abdominal diameter, or
127 visceral adipose tissue assessed by imaging modalities) adiposity. Mean±SD differences between
128 test and control arms were extracted for each outcome.

129 In those trials where the data were included in figures and not provided numerically, we used the
130 software program Plot Digitizer (<http://plotdigitizer.sourceforge.net/>) to extract the data.

131 Additional information was requested from the authors of all included trials. Disagreement were
132 resolved by consensus or where necessary by a third author (SBM).

133

134 Risk of bias assessment

135 Risk of bias for each individual study was assessed using the Cochrane Risk of Bias tool¹⁹. The
136 level of bias was evaluated for sequence generation, allocation concealment, blinding,
137 incomplete outcome data, and selective reporting and determined overall as either low (proper
138 methods taken to reduce bias), high (improper methods creating bias), or unclear (insufficient
139 information provided to determine the bias level).

140

141 Statistical analysis

Baseline, end-of-study, and between-treatment differences in body weight, BMI, waist circumference, body fat, waist-to-hip ratio, sagittal abdominal diameter or visceral adipose tissue assessed by imaging modalities were recorded as means $\pm$ SDs. If not provided, between-treatment differences in change-from-baseline or end differences were calculated by subtracting means and variance measures such as SEs were imputed with the use of published formulas¹⁹. Missing SDs were imputed with the use of the pooled SD from other studies included in the analysis¹⁹.

Data analyses were conducted using Review Manager version 5.3 (RevMan) (Copenhagen, Denmark: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014) for primary analyses and Stata version 13 (College Station, TX: StataCorp LP) for subgroup analyses. A generic inverse-variance method with random-effects models was used to calculate pooled mean differences and 95% confidence intervals (CIs). Random-effects models were used even in the absence of statistically significant inter-study heterogeneity, as they yield more conservative summary effect estimates in the presence of residual heterogeneity. Change-from-baseline differences were preferred over end differences and paired analyses were applied to all crossover trials with the use of a within-individual correlation coefficient between treatments of 0.5 as described by Elbourne et al.²¹

Inter-study heterogeneity was assessed by the Cochran Q statistic, where $P < 0.10$ was considered statistically significant, and quantified by the I^2 statistic, where $I^2 \geq 50\%$ indicates substantial heterogeneity¹⁹. Sensitivity analysis were performed which included the removal of each single study from the meta-analyses one at a time and recalculation of the summary effect. An influential outlier was considered a study whose removal changed the magnitude of the pooled effect by $>10\%$. Sensitivity analysis were also conducted using different correlation coefficient

values for crossover trials (0.25 and 0.75) to test for the robustness of the effect size, conducting analyses using fixed effects models and restricting analyses to those trials for which pasta intake could be quantified.

If ≥ 10 trial comparisons were available, then sources of heterogeneity were explored by subgroup analyses. A priori categorical subgroups analyses were assessed by meta-regression analyses. These included patient type (normal body weight, overweight or obese [average baseline BMI $>27\text{kg/m}^2$]), diabetes, coronary heart disease), follow-up (<24 -weeks, ≥ 24 weeks), baseline BMI (BMI ≤ 30 , $>30\text{kg/m}^2$), design (parallel, crossover), energy balance (negative on both arms [weight loss diets], neutral on both arms [weight maintaining diets]) and dose of pasta (based on the median). A priori categorical subgroup analyses also included the following dietary factors: GI (absolute level [≤ 55 , >55 ; glucose scale], within-treatment change, between-treatment change), fat intake (absolute level [$<30\%$, $\geq 30\%$ energy], within-treatment change, between-treatment change), carbohydrate intake (absolute level [$<50\%$, $\geq 50\%$ energy], within-treatment change, between-treatment change), protein intake (absolute level [$<20\%$, $\geq 20\%$ energy], within-treatment change, between-treatment change), dietary fibre intake (absolute level [$<28\text{g/day}$, $\geq 28\text{g/day}$], within-treatment change, between-treatment change), and risk of bias. A priori continuous meta-regression analyses were conducted on the absolute levels and within- and between-treatment changes of these same dietary factors in the intervention arms of pasta in the context of low-GI dietary patterns. Linear and non-linear pasta intake dose-response analyses were assessed by using continuous meta-regression analyses and spline curve modeling (MKSPLINE procedure), respectively. Publication bias was assessed by visual inspection of funnel plots and the Egger²² and Begg²³ tests, when ≥ 10 trial comparisons were available. If

publication bias was suspected, then adjustment for funnel plot asymmetry was done by imputing missing study data using the Duval and Tweedie trim and fill method ²⁴.

Grading the evidence

The grading of recommendations assessment, development, and evaluation (GRADE) approach was used to assess the confidence in the pooled effect estimates by assessing the overall quality and strength of the evidence ²⁵. Evidence was graded as high, moderate, low or very low quality. The included RCTs were graded as high quality evidence by default and downgraded based on pre-specified criteria. Criteria to downgrade evidence included risk of bias (weight of studies show risk of bias assessed by the Cochrane Risk of Bias tool), inconsistency (substantial unexplained heterogeneity, $I^2>50\%$, $P<0.01$), indirectness (presence of factors that limited the generalizability of the results), imprecision (the 95% CI for effect estimates were wide or crossed pre-specified minimally important differences [MIDs] for harm), and publication bias (significant evidence of small-study effects).

Patient involvement

No patients were directly involved in the development of the research question, selection of the outcome measures, design and implementation of the study, or interpretation of the results.

RESULTS

Search results

Figure 1 shows the flow of the literature. We identified 4876 reports of which 29 met eligibility criteria. No reports were identified that assessed the effect of pasta alone, while 29 reports (including 32 trial comparisons involving 2448 participants) were identified that assessed the effect of pasta in the context of low-GI dietary patterns on any adiposity outcome in adults²⁶⁻⁵⁴. Of the 32 trial comparisons that assessed the effect of pasta in the context of a low-GI dietary pattern, there were 32 trial comparisons for body weight, 18 trial comparisons for BMI^{27,28,31-33,35,36,39-41,43-46,48,49,52,53}, 18 trial comparisons for waist circumference^{27,28,31,33,34,36,38-40,42,44-47,52,53}, 10 trial comparisons for body fat^{27,28,31-33,36,38,41,43,53}, 6 trial comparisons for waist-to-hip ratio^{31,39,40,44,45,52}, and 3 trial comparisons for sagittal abdominal diameter^{34,39}. There was only one trial comparison identified³⁵ for visceral adipose tissue assessed by imaging modalities, thus a meta-analysis could not be undertaken for this outcome.

Trial characteristics

Table 1 and **Supplemental Table S3** show the characteristics of all included trials of the effect of pasta in the context of low-GI dietary patterns. The majority of trials had a parallel design (26/32) with a median follow-up of 12 weeks (IQR: 9–21 weeks) and a median number of participants per trial of 43 (IQR: 21–112). Most participants were middle aged (median age: 50 y; IQR: 40–58 y) men and women (median ratio of men to women: 0.4:1 in available trials). The median baseline BMI across studies was 30.4kg/m² (IQR: 28.2–32.0). Regarding metabolic phenotype, 21 (66%) trials included participants who were overweight or obese (had a baseline BMI $\geq$ 27kg/m²), 10 (31%) had diabetes and one trial (3%) with coronary heart disease (CHD). We did not retrieve any trials where participants had a normal BMI at baseline ($\leq$ 25kg/m²),

although 6 trials did not include BMI >25 kg/m² as part of criteria, the average baseline BMI was ≥27 kg/m², therefore categorized as overweight.

Risk of bias

Supplemental Figures S1 and S2 show the individual Cochrane Risk of Bias tool assessments for each of the included trials of the effect of pasta in the context of low-GI dietary patterns. No serious risk of bias was detected.

Effect of Pasta in the Context of Low-GI dietary patterns and Body Weight

Figure 2 shows the effect of pasta in the context of low-GI dietary patterns on the primary outcome body weight. Pooled analyses showed pasta in the context of low-GI dietary patterns had the effect of reducing body weight by -0.63 kg (95% CI:-0.84, -0.42 kg; P<0.001) compared with higher GI control diets with no evidence of heterogeneity (I²=0%, P-heterogeneity=0.51).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Global Adiposity

Figure 3 and Supplemental Figure S3-S4 show the effect of pasta in the context of low-GI dietary patterns on markers of global adiposity. Pooled analyses showed that pasta in the context of low-GI dietary patterns had the effect of reducing BMI by -0.26kg/m² (n=18 trials; MD=-0.26kg/m²; 95% CI:-0.36, -0.16 kg/m²; P<0.001) compared with higher GI control diets with no evidence of heterogeneity (I²=0%, P-heterogeneity=0.90). There was no effect on body fat (n=10 trials; MD=-0.01%, 95% CI:-0.58, 0.56%; P=0.98) with evidence of substantial heterogeneity (I²=65%, P<0.01).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Abdominal Adiposity

Figure 3 and **Supplemental Figures S5-S7** show the pooled estimates for the markers of abdominal adiposity. Pooled analyses did not show a significant effect of pasta in the context of low-GI dietary patterns compared with higher-GI control diets on waist circumference (n=18 trials; MD=-0.46cm, 95% CI:-1.05, 0.14cm; P=0.13), waist-to-hip ratio (n=6 trials; MD=-0.00, 95% CI:-0.01, 0.00; P=0.27), or sagittal abdominal diameter (n=3 trials, MD=-0.09cm, 95% CI:-0.34, 0.16cm; P=0.48) There was only evidence of substantial heterogeneity in the analyses for waist circumference ($I^2=62\%$, P-heterogeneity<0.01).

Sensitivity analyses

We conducted four sets of sensitivity analyses (**Supplemental Tables S4-5, Supplemental Figures S8-9**). The systematic removal of each trial did not modify the direction or significance of the effect estimates or the evidence of heterogeneity for any of the outcomes with the exception of waist circumference (**Supplemental Table S4**). In the sensitivity analysis for waist circumference, two studies were influential outliers in that their removal altered the magnitude of the pooled effect in the remaining studies by >10%, where the removal of the studies of McMillan-Price et al. (high protein comparison)⁴² and Jenkins et al.⁴⁴ rendered the results for waist circumference statistically significant (MD= -0.62cm; 95% CI: -1.19, -0.05; P=0.03) and (MD= -0.61cm; 95% CI: -1.18, -0.04; P=0.04), respectively; forest plots not shown). Heterogeneity remained significant in both cases ($I^2= 55\%$, P-heterogeneity <0.01 and $I^2=50\%$, P-heterogeneity= 0.01, respectively). Sensitivity analyses using correlation coefficients of 0.25 and 0.75 for paired analyses of crossover trials also did not modify the results (**Supplemental**

Table S5). In the sensitivity analyses where fixed effects models were applied (**Supplemental Figure S8**), the direction, magnitude and significance of the pooled estimates were very similar to those produced by the random effects models with the exception of the sensitivity analysis for waist circumference which was significant (MD= -0.62; 95% CI: -0.93, -0.32; P<0.001). Finally, restricting analyses to the 11 trials for which pasta intake could be quantified (median pasta intake, 3.33 servings/week [range, 1.75 – 7 servings/week]) showed a similar reduction in body weight (MD= -0.70 kg; 95% CI:-1.10, -0.29 kg; P<0.001) when pasta was consumed in the context of low-GI dietary patterns compared with the higher GI control arms without evidence of heterogeneity ($I^2=0\%$, P-heterogeneity=0.68) (**Supplemental Figure S9**).

Subgroup analyses

We were only able to conduct a priori categorical and continuous subgroup analyses for body weight, BMI, body fat and waist circumference. Subgroup analyses for waist-to-hip ratio and sagittal abdominal diameter could not be assessed, owing to <10 trial comparisons in each case. **Supplemental Figures S10-S12** show the categorical a priori subgroup analyses for body weight. There was no evidence of significant effect modification in any of the subgroup analyses for body weight, including no effect modification of follow-up when comparing studies less than 24 weeks duration to those greater than or equal to 24 weeks (-0.63kg vs -0.57kg, respectively) (**Supplemental Figure S10**). Neither was there evidence of significant effect modification in any of the subgroup analyses for BMI, body fat or waist circumference (**Supplemental Figures S13-20**).

Supplemental Table S6 and **Supplemental Figures S21-22** show the continuous subgroup analyses for body weight. There was evidence of significant effect modification by carbohydrate and protein intake, where an increase in carbohydrate intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight loss ($\beta = -0.07$, 95% CI: -0.12, -0.01, $I^2 = 0.00\%$, $P = 0.02$), and an increase in protein intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight gain ($\beta = 0.15$, 95% CI: 0.03, 0.27, $I^2 = 0.00\%$, $P = 0.02$). None of the other continuous subgroup analyses were significant. There was no evidence of significant effect modification in any of the continuous subgroup analyses for BMI (**Supplemental Table S7**). For body fat, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of difference in GI between intervention and control groups, where greater difference in GI between the groups was associated with greater reduction in body fat in the intervention group ($\beta = -0.09$, 95% CI: -0.15, -0.03, $I^2 = 19.39\%$, $P = 0.01$) (**Supplemental Table S8**). None of the other continuous subgroup analyses were significant. For waist circumference, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of absolute carbohydrate level and absolute protein level, where greater carbohydrate level in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with greater loss in waist circumference ($\beta = -0.11$, 95% CI: -0.19, -0.04, $I^2 = 27.06\%$, $P < 0.01$) and a lower protein level in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with an increase in waist circumference ($\beta = 0.20$, 95% CI: 0.01, 0.38, $I^2 = 43.92\%$, $P = 0.04$) (**Supplemental Table S9**). None of the other continuous subgroup analyses were significant.

1
2
3 322 **Dose-response analyses**

4
5 323 **Supplemental Tables S6 and S10** and **Supplemental Figure S23** show the dose-response
6
7
8 324 analysis for the 11 trials for which pasta intake could be quantified. No evidence of a linear dose
9
10 325 response was seen for pasta intake by meta-regression analyses (**Supplemental Table S6**).
11
12 326 There was also no evidence of a non-linear dose response by MKSPLINE (p=0.85)
13
14 327 (**Supplemental Figure S23**) or piecewise linear meta-regression analyses (**Supplemental Table**
15
16 328 **S10**).
17
18
19 329

20
21 330 **Publication Bias**

22
23 331 **Supplemental Figures S24-S27** shows the funnel plots for body weight, BMI, body fat and
24
25 332 waist circumference. There was no evidence of funnel-plot asymmetry. Formal testing with the
26
27 333 Egger and Begg tests did not show evidence of small-study effects (P>0.05 for both). Publication
28
29 334 bias was not assessed for waist-to-hip ratio and sagittal abdominal diameter, owing to <10 trial
30
31 335 comparisons.
32
33
34 336

35
36
37 337 **GRADE Assessment**

38
39 338 **Supplemental Table S11** shows a summary of the GRADE assessments for the effect of pasta in
40
41 339 the context of low-GI dietary patterns on body weight and measures of adiposity. The evidence
42
43 340 was graded as moderate for body weight, BMI, waist-to-hip ratio and sagittal abdominal
44
45 341 diameter owing to downgrades for indirectness and low for waist circumference and body fat,
46
47 342 owing to downgrades for indirectness and inconsistency (I²=59%, P-heterogeneity<0.001;
48
49 343 I²=66%, P-heterogeneity<0.01, respectively).
50
51
52
53

DISCUSSION

The present systematic review and meta-analysis quantified the effect of pasta alone and pasta in the context of low-GI dietary patterns on body weight and other markers of adiposity. We identified no trial comparisons for the effect of pasta alone and 32 trial comparisons involving 2448 participants who were predominantly middle-aged and overweight or obese for the effect of pasta in the context of low-GI dietary patterns. The primary pooled analysis demonstrated that pasta in the context of low-GI dietary patterns did not contribute to weight gain, resulting in a significant weight loss of -0.63 kg when compared to diets higher in GI over a median follow-up of 12 weeks. The lack of harm was reflected in the established clinical secondary outcome measures of global (BMI and body fat) and abdominal adiposity (waist circumference, waist-to-hip ratio and sagittal abdominal diameter). These findings were robust across subgroups. The findings did not differ by metabolic phenotype in those who were overweight or obese or had diabetes, which is noteworthy since these are populations who would benefit from weight management strategies. There was also no effect modification by the energy balance of the design such that the weight loss was seen even under conditions of neutral energy balance (in which participants were instructed to consume dietary advice ad libitum), suggesting that encouragement of the consumption of pasta in the context of a low-GI dietary patterns does not cause harm and may even lead to spontaneous weight loss. There was also no effect modification by follow-up either in continuous meta-regression or categorical, where the 24 trials with <24 weeks follow-up had a weight reduction similar to those 8 trials with ≥24 weeks follow-up (-0.63 kg vs -0.57 kg, respectively). This is of relevance since many dietary studies are successful in demonstrating weight loss in the short term but not over the long term.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Findings in the context of existing studies

We are not aware of any RCTs directly assessing the effect of pasta intake on health parameters including body weight. Our findings, however, agree with earlier systematic reviews and meta-analyses of RCTs of the effect of low-GI dietary patterns irrespective of pasta intake on body weight and adiposity. The systematic review and meta-analysis by Thomas et al. in 2007 found a significant -1.1kg weight loss and -1.3kg/m² reduction in BMI favouring low-GI or glycemic load (GL) diets compared to control diets in 6 RCTs of 5 weeks to 6 months in duration in overweight or obese individuals¹³. Another systematic review and meta-analysis by Schwingshackl et al. in 2013 found a -0.62kg weight loss favouring low-GI/GL diets compared to higher GI/GL diets in 14 RCTs with greater than 6 months duration in overweight individuals (BMI>25kg/m²)¹⁴.

Our findings also agree with trials in which pasta was emphasized in the context of other healthy dietary patterns. One trial done in children in Spain given Mediterranean dietary advice which included increasing the intake of pasta found that approximately 11.3% of the participants in the Mediterranean diet group who were classified as overweight and obese changed their weight status to normal weight compared to only approximately 2.6% of the participants in the control group⁵⁵.

Other lines of evidence from observational studies have demonstrated benefits of pasta consumption on body weight and adiposity. Pasta intake was recently assessed in the Moli-sani study and the Italian Nutrition & HEalth Survey (INHES), a cross-sectional study of over 20,000 Italians from all over Italy⁵⁶. The study demonstrated that higher pasta consumption was

391 associated with lower BMI, waist circumference and waist-to-hip ratio and with a lower
392 prevalence of overweight and obesity⁵⁶. Furthermore, greater pasta intake was associated with
393 better adherence to the Mediterranean diet, a dietary pattern which has a demonstrated
394 cardiovascular benefit⁵⁷. Similar associations between greater pasta intake and lower body
395 weight have also been observed in prospective U.S. cohorts⁵⁸. A pooled analysis of the 3
396 Harvard cohorts also showed that higher GI diets (which would preclude pasta) were associated
397 with weight gain⁵⁹.

399 Although the product form of pasta can vary widely, including in shape (e.g. macaroni, spaghetti,
400 linguine), ingredients (e.g. type of wheat, egg content), and processing technique (e.g. drying
401 temperature), studies have demonstrated that when comparing pastas varying in these
402 parameters, despite slight variations in glycemic response among pastas, glycemic responses are
403 still lower compared to a control, e.g. white bread^{60,61}. One concern of the choice of pasta as a
404 carbohydrate food is that it is a refined food low in fibre. Although there are whole grain pasta
405 options available, studies have demonstrated that fiber added to pasta, does not significantly
406 affect the glucose or insulin response, the secretion of gut hormones or satiety^{62,63}. Furthermore,
407 pasta has a similar glycemic response compared to many fiber-rich carbohydrates, including
408 barley, legumes and steel cut oats, and still a lower glycemic response compared to other fiber-
409 rich foods including whole wheat bread, breakfast cereals like bran flakes and potatoes with
410 skin⁶⁴.

411
412 The mechanism by which pasta in the context of low-GI dietary patterns lead to weight loss even
413 under conditions of ad libitum dietary advice is unclear. Lower GI diets may result in greater

1
2
3 414 body weight reduction compared to higher GI diets because lower GI foods may be more
4
5 415 satiating⁶⁵ and have been shown to delay hunger and decrease subsequent energy intake¹³. Low-
6
7
8 416 GI dietary patterns are also characterized by high fiber content^{64,66} which may also contribute to
9
10 417 improvements in satiety and hunger¹⁷. Furthermore, studies which have compared ad libitum
11
12 418 low-GI dietary patterns to standard energy-restricted low-fat diets, demonstrated equivalency or
13
14 419 better weight loss when following the low-GI diet, despite the fact that they could eat as much as
15
16 420 they desired^{13,67}. Thus, voluntary energy intake may be lower after low-GI meals, as has been
17
18 421 previously demonstrated⁶⁸.

22 422

24 423 **Strengths and limitations**

26 424 The strengths of the present systematic review and meta-analysis include that it is
27
28 425 comprehensive, includes RCTs which protects against bias and uses GRADE to evaluate the
29
30 426 quality of evidence. Additionally, a large number of trials were identified (32 trials) for the
31
32 427 primary outcome of body weight, the median follow-up period was 12 weeks which allows for
33
34 428 the assessment of a moderate duration of intervention, none of the trials were rated as having a
35
36 429 serious risk of bias, and there was no evidence of publication bias.

40 430

42 431 There are several limitations. First, the present systematic review and meta-analysis showed
43
44 432 evidence of inconsistency in the treatment estimates across trials for some of the outcomes
45
46 433 assessed. There was evidence of unexplained heterogeneity in waist circumference ($I^2=62\%$) and
47
48 434 in body fat ($I^2=65\%$). Although the inconsistency in these outcomes may have related to
49
50 435 measurement error⁶⁹ in the different techniques for measuring waist circumference and body fat,
51
52 436 we were unable to conduct sensitivity or subgroup analyses to explore this source of

56 437

58 438

60 439

heterogeneity. Second, there was evidence of indirectness. Most of the available trials did not quantify the amount of pasta consumed in the context of the low-GI dietary patterns. Although sensitivity analyses in which analyses were restricted to the 11 trials that did quantify (providing a median 3.33 servings/week) pasta intake did not meaningfully alter our estimates (-0.70kg versus -0.63kg), it is difficult to quantify the effect of pasta in these diets. There is also the question of translation to other background diets. Whether the observed effect of pasta in the context of low-GI dietary patterns will hold in the context of other healthy dietary patterns, such as Mediterranean and Vegetarian dietary patterns, is unclear. Although there is no biological reason to doubt that the findings would hold, there is a lack of direct evidence to support this conclusion. If the question had been asked from the perspective of benefit as opposed to that of harm, then the relatively short duration of the included trials may be another reason to downgrade for indirectness. In the absence of long-term trials (>1 year diet duration), it is difficult to conclude with certainty that the observed lack of harm implies an actual sustainable benefit. Finally, there was some evidence of imprecision for benefit but not harm. Whereas the 95% CI of the pooled estimates did not overlap with our pre-specified MID for harm (that is, they did not contain evidence for harm) and so were not downgraded for imprecision, the upper bound of the 95% CI would overlap with the lower bound of the same MID to assess the precision of the evidence for benefit for some outcomes.

Balancing these strengths and limitations, GRADE assessed the overall quality and strength of the available evidence of the effect of pasta in the context of low-GI dietary patterns as moderate for the primary outcome of body weight and the secondary outcomes of BMI, waist-to-hip ratio and sagittal abdominal diameter owing to downgrades for indirectness. The evidence was

assessed as low for the other secondary outcomes of body fat and waist circumference owing to downgrades for indirectness and inconsistency.

Implications

These results are important considering the negative messages directed at the public regarding carbohydrates, which is influencing their food choices, as is evident in recent reductions in carbohydrate intake⁷⁰⁻⁷², and in particular reductions in pasta consumption^{70,73-76}. Contrary to these concerns, the available evidence shows that when pasta is consumed in the context of low-GI dietary patterns that there is not weight gain but rather marginally clinically significant weight loss (>0.5kg)⁷⁷.

Although we were able to approximate the amount of pasta consumed in one third of included trials, it is unclear what the effect of pasta would be in the context of other dietary patterns. Low-GI dietary patterns, however, shares many similarities with a Mediterranean diet, which emphasizes many low-GI foods and has demonstrated a CVD benefit⁵⁷.

Current clinical practice guidelines already suggest the replacement of high GI foods with low-GI foods for improvement of glycemic control and cardiovascular risk factors^{78,79}. The present evidence means that pasta may be highlighted as an important example of a low-GI food which can contribute to a low-GI dietary pattern, a pattern which in turn may potentially improve cardio-metabolic risk without an adverse effect on weight control.

CONCLUSIONS

In conclusion, pasta consumed in the context of low-GI dietary patterns does not have an adverse effect on body weight and adiposity outcomes of importance in the prevention and management of overweight and obesity. The results are generalizable in the context of a high carbohydrate diet composed of low-GI foods with or without the intention of weight loss in middle-aged individuals who are overweight or obese or have diabetes. Although the clinical significance of the observed weight loss is debatable, this finding increases our confidence that pasta in the context of low-GI dietary patterns does not result in weight gain. Further research may change our confidence in the estimates for our primary outcome body weight and several key secondary outcomes including BMI and two measures of abdominal adiposity, waist-to-hip ratio and sagittal abdominal diameter. More research, however, is needed, to improve our estimates for the secondary outcomes, body fat and waist circumference and assess whether our findings extend to related cardio-metabolic outcomes. There is also a need for more randomized trials of >1 year diet duration to clarify whether the lack of harm for pasta in the context of low-GI dietary patterns will translate into meaningful long-term benefits. Other randomized trials should focus on whether pasta will have similar effects in the context of other healthy dietary patterns such as a Mediterranean diet.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

Figure 1: Literature Search

Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). n = 2448. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.
CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.
BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

527 Acknowledgements

528 We wish to thank Teruko Kishibe, an information specialist in the Scotiabank Health Sciences
529 Library, St Michael's Hospital, for her help in the development of search terms used. Aspects of
530 this work were presented at the Società Italiana di Nutrizione Umana (SINU) Congress, in
531 Parma, Italy, on April 12, 2017 and at the 35th International Symposium on Diabetes and
532 Nutrition, of the Diabetes and Nutrition Study Group (DNSG) of the European Association for
533 the Study of Diabetes (EASD) in Skagen, Denmark, on June 20 and 21, 2017.

534 Data Sharing

535 No additional data available.

537 Exclusive License

538 The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf
539 of all authors, an exclusive license (or non-exclusive for government employees) on a worldwide
540 basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in BMJ
541 editions and any other BMJ PGL products and sublicenses such use and exploit all subsidiary
542 rights, as set out in our license.

543 Ethics Approval

544 Not required.

545 Authorship:

546 All authors had full access to all of the data (including statistical reports and tables) in this study
547 and take full responsibility for the integrity of the data and the accuracy of the data analysis.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conception and design: C.W.C. Kendall, J.L. Sievenpiper.

Analysis and interpretation of the data: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Drafting of the article: L. Chiavaroli, J.L. Sievenpiper.

Critical revision of the article for important intellectual content: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Final approval of the article: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Obtaining of funding: C.W.C. Kendall, D.J.A. Jenkins, J.L. Sievenpiper.

Administrative, technical, or logistic support: S. Blanco Mejia.

Collection and assembly of data: L. Chiavaroli, C.R. Braunstein, S. Blanco Mejia.

Guarantor: J.L. Sievenpiper.

Transparency declaration: As guarantor, J.L. Sievenpiper affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Funding:

 This work was funded by the Canadian Institutes of Health Research (funding reference number, 129920) through the Canada-wide Human Nutrition Trialists' Network (NTN). The Diet, Digestive tract, and Disease (3-D) Centre, funded through the Canada Foundation for Innovation (CFI) and the Ministry of Research and Innovation's Ontario Research Fund (ORF),

provided the infrastructure for the conduct of this project. **L Chiavaroli** was funded by a Toronto 3D Postdoctoral Fellowship Award. **DJA Jenkins** was funded by the Government of Canada through the Canada Research Chair Endowment. **JL Sievenpiper** was funded by a PSI Graham Farquharson Knowledge Translation Fellowship, Canadian Diabetes Association (CDA) Clinician Scientist award, CIHR INMD/CNS New Investigator Partnership Prize, and Banting & Best Diabetes Centre Sun Life Financial New Investigator Award. None of the sponsors had a role in any aspect of the present study, including design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, approval of the manuscript or decision to publish.

Conflicts of Interest:

All authors have completed the Unified Competing Interest form (available on request from the corresponding author) and declare: **CWC Kendall** has received research support from the Advanced Foods and Material Network, Agrifoods and Agriculture Canada, the Almond Board of California, the American Pistachio Growers, Barilla, the California Strawberry Commission, the Calorie Control Council, CIHR, the Canola Council of Canada, the Coca-Cola Company (investigator initiated, unrestricted grant), Hain Celestial, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Kraft, Loblaw Companies Ltd., Orafiti, Pulse Canada, Saskatchewan Pulse Growers, Solae and Unilever. He has received travel funding, consultant fees and/or honoraria from Abbott Laboratories, the Almond Board of California, the American Peanut Council, the American Pistachio Growers, Barilla, Bayer, the Canola Council of Canada, the Coca-Cola Company, Danone, General Mills, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Loblaw Companies Ltd., the Nutrition Foundation of Italy, Oldways Preservation Trust, Orafiti, Paramount Farms, the Peanut Institute,

1
2
3 593 PepsiCo, Pulse Canada, Sabra Dipping Co., Saskatchewan Pulse Growers, Solae, Sun-Maid, Tate
4
5 594 and Lyle, and Unilever. He is on the Clinical Practice Guidelines Expert Committee for Nutrition
6
7 595 Therapy of the European Association for the Study of Diabetes (EASD) and has served on the
8
9
10 596 scientific advisory board for the Almond Board of California, the International Tree Nut Council,
11
12 597 Oldways Preservation Trust, Paramount Farms and Pulse Canada. **DJAJ** has received research
13
14 598 grants from Saskatchewan Pulse Growers, the Agricultural Bioproducts Innovation Program
15
16 599 through the Pulse Research Network, the Advanced Foods and Material Network, Loblaw
17
18 600 Companies Ltd., Unilever, Barilla, the Almond Board of California, Agriculture and Agri-food
19
20 601 Canada, Pulse Canada, Kellogg's Company, Canada, Quaker Oats, Canada, Procter & Gamble
21
22 602 Technical Centre Ltd., Bayer Consumer Care, Springfield, NJ, Pepsi/Quaker, International Nut
23
24 603 & Dried Fruit (INC), Soy Foods Association of North America, the Coca-Cola Company
25
26 604 (investigator initiated, unrestricted grant), Solae, Haine Celestial, the Sanitarium Company,
27
28 605 Orafit, the International Tree Nut Council Nutrition Research and Education Foundation, the
29
30 606 Peanut Institute, the Canola and Flax Councils of Canada, the Calorie Control Council (CCC),
31
32 607 the CIHR, the Canada Foundation for Innovation and the Ontario Research Fund. He has
33
34 608 received in-kind supplies for trial as a research support from the Almond board of California,
35
36 609 Walnut Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo,
37
38 610 Loblaw Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada,
39
40 611 WhiteWave Foods. He has been on the speaker's panel, served on the scientific advisory board
41
42 612 and/or received travel support and/or honoraria from the Almond Board of California, Canadian
43
44 613 Agriculture Policy Institute, Loblaw Companies Ltd, the Griffin Hospital (for the development of
45
46 614 the NuVal scoring system, the Coca-Cola Company, EPICURE, Danone, Diet Quality Photo
47
48 615 Navigation (DQPN), FareWell, Verywell, True Health Initiative, Saskatchewan Pulse Growers,
49
50
51
52
53
54
55
56
57
58
59
60

Sanitarium Company, Orafit, the Almond Board of California, the American Peanut Council, the International Tree Nut Council Nutrition Research and Education Foundation, the Peanut Institute, Herbalife International, Pacific Health Laboratories, Nutritional Fundamental for Health, Barilla, Metagenics, Bayer Consumer Care, Unilever Canada and Netherlands, Solae, Kellogg, Quaker Oats, Procter & Gamble, the Coca-Cola Company, the Griffin Hospital, Abbott Laboratories, the Canola Council of Canada, Dean Foods, the California Strawberry Commission, Haine Celestial, PepsiCo, the Alpro Foundation, Pioneer Hi-Bred International, DuPont Nutrition and Health, Spherix Consulting and WhiteWave Foods, the Advanced Foods and Material Network, the Canola and Flax Councils of Canada, the Nutritional Fundamentals for Health, Agri-Culture and Agri-Food Canada, the Canadian Agri-Food Policy Institute, Pulse Canada, the Saskatchewan Pulse Growers, the Soy Foods Association of North America, the Nutrition Foundation of Italy (NFI), Nutra-Source Diagnostics, the McDougall Program, the Toronto Knowledge Translation Group (St. Michael's Hospital), the Canadian College of Naturopathic Medicine, The Hospital for Sick Children, the Canadian Nutrition Society (CNS), the American Society of Nutrition (ASN), Arizona State University, Paolo Sorbini Foundation and the Institute of Nutrition, Metabolism and Diabetes. He received an honorarium from the United States Department of Agriculture to present the 2013 W.O. Atwater Memorial Lecture. He received the 2013 Award for Excellence in Research from the International Nut and Dried Fruit Council. He received funding and travel support from the Canadian Society of Endocrinology and Metabolism to produce mini cases for the Canadian Diabetes Association (CDA). He is a member of the International Carbohydrate Quality Consortium (ICQC). His wife, ALJ, is a director and partner of Glycemic Index Laboratories, Inc., and his sister received funding through a grant from the St. Michael's Hospital Foundation to develop a cookbook for

one of his studies. **JL Sievenpiper** has received research support from the Canadian Institutes of health Research (CIHR), Canadian Diabetes Association (CDA), PSI Foundation, Banting and Best Diabetes Centre (BBDC), Canadian Nutrition Society (CNS), American Society for Nutrition (ASN), Calorie Control Council, INC International Nut and Dried Fruit Council Foundation, National Dried Fruit Trade Association, The Coca-Cola Company (investigator initiated, unrestricted), Dr. Pepper Snapple Group (investigator initiated, unrestricted), The Tate and Lyle Nutritional Research Fund at the University of Toronto, and The Alberta Pulse Growers Glycemic Control and Cardiovascular Disease in Type 2 Diabetes Fund at the University of Toronto. He has received in-kind research support from the Almond board of California, Walnut Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo, Loblaw Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada, WhiteWave Foods. He has received reimbursement of travel expenses, speaker fees, and/or honoraria from the Canadian Diabetes Association (CDA), Canadian Nutrition Society (CNS), University of Alabama at Birmingham, Oldways Preservation Trust, Nutrition Foundation of Italy (NFI), Diabetes and Nutrition Study Group (DNSG) of the European Association for the Study of Diabetes (EASD), International Life Sciences Institute (ILSI) North America, Dr. Pepper Snapple Group, Corn Refiners Association, World Sugar Research Organization, Dairy Farmers of Canada, Società Italiana di Nutrizione Umana (SINU), III World Congress of Public Health Nutrition, C3 Collaborating for Health, Sprim Brasil, WhiteWave Foods, Rippe Lifestyle, mdBriefcase, Federation of European Nutrition Societies (FENS), New York Academy of Sciences, International Diabetes Federation, American Heart Association (AHA), American Society for Nutrition (ASN), FoodMinds LLC, Memac Ogilvy & Mather LLC, Pulse Canada, Pepsico, BCFN Foundation, The Ginger Network LLC, and Dietitians of Canada. He has ad hoc

consulting arrangements with Winston & Strawn LLP, Perkins Coie LLP, and Tate & Lyle. He is a member of the European Fruit Juice Association Scientific Expert Panel. He is on Clinical Practice Guidelines Expert Committees of the Canadian Diabetes Association (CDA) European Association for the study of Diabetes (EASD), and Canadian Cardiovascular Society (CCS), as well as an expert writing panel of the American Society for Nutrition (ASN). He serves as an unpaid scientific advisor for the Food, Nutrition, and Safety Program (FNSP) and the Technical Committee on Carbohydrates of the International Life Science Institute (ILSI) North America. He is a member of the International Carbohydrate Quality Consortium (ICQC), Executive Board Member of the Diabetes and Nutrition Study Group (DNSG) of the EASD, and Director of the Toronto 3D Knowledge Synthesis and Clinical Trials foundation. His wife is an employee of Unilever Canada. No competing interests were declared by **CR Braunstein, S Blanco Mejia, and LA Leiter.**

References

1. Malhotra A. Eating MORE fat while cutting carbs and quitting sugar can help you lose weight and be happier, says top cardiologist. Mail Online 2016.

2. Taubes G. Diet Advice That Ignores Hunger. The New York Times. Sunday Review ed 2015.

3. Ludwig D. Always hungry? Conquer cravings, retrain your fat cells and lose weight permanently. New York: Grand Central Life & Style; 2016.

4. Taubes G. Why We Get Fat And What To Do About It. . New York: Alfred A Knopf; 2011.

5. Atkins R. Dr Atkins' Diet Revolution: The High Calorie Way to Stay Thin Forever. : Bantam Books; 1973.

6. Perlmutter D. Grain Brain: The surprising truth about wheat, carbs, and sugar -- Your brain's silent killers: Little, Brown And Company; 2013.

7. David W. Wheat Belly: HarperCollins; 2012.

8. Cordain L. The Paleo Diet Revised: Lose weight and get healthy by eating the foods you were designed to eat: Houghton Mifflin Harcourt; 2010.

9. Teicholz N. The Big Fat Surprise: Why Butter, Meat and Cheese Belong in a Healthy Diet: Simon and Schuster; 2014.

10. National Obesity Forum, The Public Health Collaboration. Eat Fat, Cut The Carbs and Avoid Snacking To Reverse Obesity and Type 2 Diabetes.

11. Malhotra A. Saturated fat is not the major issue. BMJ 2013;347:f6340.

12. Malhotra A. Author's reply to Lichtenstein, Tedstone and Pyne, Mann and colleagues, Lim, and Clifton. BMJ 2013;347:f6855.

13. Thomas DE, Elliott EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. Cochrane Database Syst Rev 2007;CD005105.

14. Schwingshackl L, Hoffmann G. Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. Nutr Metab Cardiovasc Dis 2013;23:699-706.

15. Pol K, Christensen R, Bartels EM, Raben A, Tetens I, Kristensen M. Whole grain and body weight changes in apparently healthy adults: a systematic review and meta-analysis of randomized controlled studies. Am J Clin Nutr 2013;98:872-84.

16. Harland JI, Garton LE. Whole-grain intake as a marker of healthy body weight and adiposity. Public Health Nutr 2008;11:554-63.

17. Slavin JL. Dietary fiber and body weight. Nutrition 2005;21:411-8.

18. Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: Health Implications of Dietary Fiber. J Acad Nutr Diet 2015;115:1861-70.

19. Higgins JP, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions [Internet]. Version 5.1.0. The Cochrane Collaboration. Version current March 2011 [cited 2015 Sep 23]. Available from: www.cochrane-handbook.org.

20. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Ann Intern Med 2009;151:264-9, W64.

21. Elbourne DR, Altman DG, Higgins JP, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. Int J Epidemiol 2002;31:140-9.

22. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ 1997;315:629-34.

23. Begg CB. A measure to aid in the interpretation of published clinical trials. Stat Med 1985;4:1-9.

24. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 2000;56:455-63.

25. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383-94.
26. Jebb SA, Lovegrove JA, Griffin BA, et al. Effect of changing the amount and type of fat and carbohydrate on insulin sensitivity and cardiovascular risk: the RISCK (Reading, Imperial, Surrey, Cambridge, and Kings) trial. *Am J Clin Nutr* 2010;92:748-58.
27. Philippou E, Bovill-Taylor C, Rajkumar C, et al. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 2009;58:1703-8.
28. Philippou E, McGowan BM, Brynes AE, Dornhorst A, Leeds AR, Frost GS. The effect of a 12-week low glycaemic index diet on heart disease risk factors and 24 h glycaemic response in healthy middle-aged volunteers at risk of heart disease: a pilot study. *Eur J Clin Nutr* 2008;62:145-9.
29. Sichieri R, Moura AS, Genelhu V, Hu F, Willett WC. An 18-mo randomized trial of a low-glycemic-index diet and weight change in Brazilian women. *Am J Clin Nutr* 2007;86:707-13.
30. Wolever TM, Mehling C. High-carbohydrate-low-glycaemic index dietary advice improves glucose disposition index in subjects with impaired glucose tolerance. *Br J Nutr* 2002;87:477-87.
31. Pereira EV, Costa Jde A, Alfenas Rde C. Effect of glycemic index on obesity control. *Arch Endocrinol Metab* 2015;59:245-51.
32. Karl JP, Roberts SB, Schaefer EJ, et al. Effects of carbohydrate quantity and glycemic index on resting metabolic rate and body composition during weight loss. *Obesity (Silver Spring)* 2015;23:2190-8.
33. Buscemi S, Cosentino L, Rosafio G, et al. Effects of hypocaloric diets with different glycemic indexes on endothelial function and glycemic variability in overweight and in obese adult patients at increased cardiovascular risk. *Clin Nutr* 2013;32:346-52.
34. Larsen TM, Dalskov SM, van Baak M, et al. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med* 2010;363:2102-13.
35. Solomon TP, Haus JM, Kelly KR, et al. A low-glycemic index diet combined with exercise reduces insulin resistance, postprandial hyperinsulinemia, and glucose-dependent insulinotropic polypeptide responses in obese, prediabetic humans. *Am J Clin Nutr* 2010;92:1359-68.
36. Philippou E, Neary NM, Chaudhri O, et al. The effect of dietary glycemic index on weight maintenance in overweight subjects: a pilot study. *Obesity (Silver Spring)* 2009;17:396-401.
37. Abete I, Parra D, Martinez JA. Energy-restricted diets based on a distinct food selection affecting the glycemic index induce different weight loss and oxidative response. *Clin Nutr* 2008;27:545-51.
38. Aston LM, Stokes CS, Jebb SA. No effect of a diet with a reduced glycaemic index on satiety, energy intake and body weight in overweight and obese women. *Int J Obes (Lond)* 2008;32:160-5.
39. Jensen L, Sloth B, Krog-Mikkelsen I, et al. A low-glycemic-index diet reduces plasma plasminogen activator inhibitor-1 activity, but not tissue inhibitor of proteinases-1 or plasminogen activator inhibitor-1 protein, in overweight women. *Am J Clin Nutr* 2008;87:97-105.
40. Bellisle F, Dalix AM, De Assis MA, et al. Motivational effects of 12-week moderately restrictive diets with or without special attention to the Glycaemic Index of foods. *Br J Nutr* 2007;97:790-8.
41. de Rougemont A, Normand S, Nazare JA, et al. Beneficial effects of a 5-week low-glycaemic index regimen on weight control and cardiovascular risk factors in overweight non-diabetic subjects. *Br J Nutr* 2007;98:1288-98.
42. McMillan-Price J, Petocz P, Atkinson F, et al. Comparison of 4 diets of varying glycemic load on weight loss and cardiovascular risk reduction in overweight and obese young adults: a randomized controlled trial. *Arch Intern Med* 2006;166:1466-75.
43. Visek J, Lacigova S, Cechurova D, Rusavy Z. Comparison of a low-glycemic index vs standard diabetic diet. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:112-6.

1
2
3 779 44. Jenkins DJ, Kendall CW, Vuksan V, et al. Effect of lowering the glycemic load with canola oil on
4 780 glycemic control and cardiovascular risk factors: a randomized controlled trial. *Diabetes Care*
5 781 2014;37:1806-14.
6 782 45. Jenkins DJ, Kendall CW, Augustin LS, et al. Effect of legumes as part of a low glycemic index diet
7 783 on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled
8 784 trial. *Arch Intern Med* 2012;172:1653-60.
9 785 46. Yusof BN, Talib RA, Kamaruddin NA, Karim NA, Chinna K, Gilbertson H. A low-GI diet is
10 786 associated with a short-term improvement of glycaemic control in Asian patients with type 2 diabetes.
11 787 *Diabetes Obes Metab* 2009;11:387-96.
12 788 47. Wolever TM, Gibbs AL, Mehling C, et al. The Canadian Trial of Carbohydrates in Diabetes (CCD),
13 789 a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on
14 790 glycated hemoglobin but reduction in C-reactive protein. *Am J Clin Nutr* 2008;87:114-25.
15 791 48. Jenkins DJ, Kendall CW, McKeown-Eyssen G, et al. Effect of a low-glycemic index or a high-cereal
16 792 fiber diet on type 2 diabetes: a randomized trial. *JAMA* 2008;300:2742-53.
17 793 49. Jimenez-Cruz A, Bacardi-Gascon M, Turnbull WH, Rosales-Garay P, Severino-Lugo I. A flexible,
18 794 low-glycemic index mexican-style diet in overweight and obese subjects with type 2 diabetes improves
19 795 metabolic parameters during a 6-week treatment period. *Diabetes Care* 2003;26:1967-70.
20 796 50. Heilbronn LK, Noakes M, Clifton PM. The effect of high- and low-glycemic index energy
21 797 restricted diets on plasma lipid and glucose profiles in type 2 diabetic subjects with varying glycemic
22 798 control. *J Am Coll Nutr* 2002;21:120-7.
23 799 51. Fontvieille AM, Acosta M, Rizkalla SW, et al. A moderate switch from high to low glycemic-index
24 800 foods for 3 weeks improves metabolic control of type I (IDDM) diabetic subjects. . *Diabetes Nutr Metab*
25 801 1988;1:139-43.
26 802 52. Frost GS, Brynes AE, Bovill-Taylor C, Dornhorst A. A prospective randomised trial to determine
27 803 the efficacy of a low glycaemic index diet given in addition to healthy eating and weight loss advice in
28 804 patients with coronary heart disease. *Eur J Clin Nutr* 2004;58:121-7.
29 805 53. de Assis Costa J, de Cassia Goncalves Alfensa R. The consumption of low glycemic meals reduces
30 806 abdominal obesity in subjects with excess body weight. *Nutricion Hospitalaria* 2012;27:1178-83.
31 807 54. Fontvieille AM, Rizkalla SW, Penfornis A, Acosta M, Bornet FR, Slama G. The use of low
32 808 glycaemic index foods improves metabolic control of diabetic patients over five weeks. *Diabet Med*
33 809 1992;9:444-50.
34 810 55. Bibiloni MDM, Fernandez-Blanco J, Pujol-Plana N, et al. [Improving diet quality in children
35 811 through a new nutritional education programme: INFADIMED]. *Gac Sanit* 2017.
36 812 56. Pounis G, Castelnuovo AD, Costanzo S, et al. Association of pasta consumption with body mass
37 813 index and waist-to-hip ratio: results from Moli-sani and INHES studies. *Nutr Diabetes* 2016;6:e218.
38 814 57. Estruch R, Ros E, Salas-Salvado J, et al. Primary prevention of cardiovascular disease with a
39 815 Mediterranean diet. *N Engl J Med* 2013;368:1279-90.
40 816 58. Shay CM, Van Horn L, Stamler J, et al. Food and nutrient intakes and their associations with
41 817 lower BMI in middle-aged US adults: the International Study of Macro-/Micronutrients and Blood
42 818 Pressure (INTERMAP). *Am J Clin Nutr* 2012;96:483-91.
43 819 59. Smith JD, Hou T, Ludwig DS, et al. Changes in intake of protein foods, carbohydrate amount and
44 820 quality, and long-term weight change: results from 3 prospective cohorts. *Am J Clin Nutr* 2015;101:1216-
45 821 24.
46 822 60. Granfeldt Y, Bjorck I. Glycemic response to starch in pasta: a study of mechanisms of limited
47 823 enzyme availability. . *J Cereal Sci* 1991;14:47-61.
48 824 61. Granfeldt Y, Bjorck I, Hagander B. On the importance of processing conditions, product thickness
49 825 and egg addition for the glycaemic and hormonal responses to pasta: a comparison with bread made
50 826 from 'pasta ingredients'. *Eur J Clin Nutr* 1991;45:489-99.

62. Frost GS, Brynes AE, Dhillon WS, Bloom SR, McBurney MI. The effects of fiber enrichment of pasta and fat content on gastric emptying, GLP-1, glucose, and insulin responses to a meal. *Eur J Clin Nutr* 2003;57:293-8.
63. Korczak R, Timm D, Ahnen R, Thomas W, Slavin JL. High Protein Pasta is Not More Satiating than High Fiber Pasta at a Lunch Meal, Nor Does it Decrease Mid-Afternoon Snacking in Healthy Men and Women. *J Food Sci* 2016;81:S2240-5.
64. Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care* 2008;31:2281-3.
65. Ludwig DS. Dietary glycemic index and obesity. *J Nutr* 2000;130:280S-3S.
66. Riccardi G, Rivellese AA, Giacco R. Role of glycemic index and glycemic load in the healthy state, in prediabetes, and in diabetes. *Am J Clin Nutr* 2008;87:269S-74S.
67. Saris WH, Astrup A, Prentice AM, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the CARMEN study. The Carbohydrate Ratio Management in European National diets. *Int J Obes Relat Metab Disord* 2000;24:1310-8.
68. Ludwig DS, Majzoub JA, Al-Zahrani A, Dallal GE, Blanco I, Roberts SB. High glycemic index foods, overeating, and obesity. *Pediatrics* 1999;103:E26.
69. Wang J, Thornton JC, Bari S, et al. Comparisons of waist circumferences measured at 4 sites. *Am J Clin Nutr* 2003;77:379-84.
70. Makarem N, Scott M, Quatromoni P, Jacques P, Parekh N. Trends in dietary carbohydrate consumption from 1991 to 2008 in the Framingham Heart Study Offspring Cohort. *Br J Nutr* 2014;111:2010-23.
71. Rehm CD, Penalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults, 1999-2012. *JAMA* 2016;315:2542-53.
72. Golzarand M, Mirmiran P, Jessri M, Toolabi K, Mojarrad M, Azizi F. Dietary trends in the Middle East and North Africa: an ecological study (1961 to 2007). *Public Health Nutr* 2012;15:1835-44.
73. Leclercq C, Arcella D, Piccinelli R, et al. The Italian National Food Consumption Survey INRAN-SCAI 2005-06: main results in terms of food consumption. *Public Health Nutr* 2009;12:2504-32.
74. Bonaccio M, Di Castelnuovo A, Bonanni A, et al. Decline of the Mediterranean diet at a time of economic crisis. Results from the Moli-sani study. *Nutr Metab Cardiovasc Dis* 2014;24:853-60.
75. Sofi F, Innocenti G, Dini C, et al. Low adherence of a clinically healthy Italian population to nutritional recommendations for primary prevention of chronic diseases. *Nutr Metab Cardiovasc Dis* 2006;16:436-44.
76. Ferdman RA. This is what happens when everyone is terrified of carbs. *The Washington Post* 2015 September 2, 2015.
77. Johnston BC, Kanters S, Bandayrel K, et al. Comparison of weight loss among named diet programs in overweight and obese adults: a meta-analysis. *JAMA* 2014;312:923-33.
78. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;129:S1-45.
79. Anderson TJ, Gregoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. *Can J Cardiol* 2013;29:151-67.

Table 1: Summary of trial characteristics

Trial Characteristics*	ALL	Neutral Energy Balance	Negative Energy Balance
Trial Number (n)	32	23	9
Trial Size (total, range)	2448 (8 - 250)	1989 (8 - 250)	459 (13 - 123)
Male: Female ^a (%)	40:60	47:53	27:73
Age (years)	50 (40 - 58)	52.0 (42.1 – 59.5)	49.5 (34.2 – 53.0)
Metabolic Phenotype (OW/OB:DM:CHD) (%)	66:31:3	57:39:4	89:11:0
Setting (IP:OP) (%)	3:97	4:96	0:100
Baseline Body Weight (kg) ^b	85.5 (80.0 - 91.9)	84.1 (79.5 – 87.5)	92.5 (86.1 – 93.9)
Baseline BMI (kg/m ²) ^c	30.4 (28.2 - 32.0)	29.5 (27.4 – 31.4)	31.7 (30.1 – 32.9)
Study Design (C:P) (%)	19:81	26:74	0:100
Dose Pasta (servings/week) ^d	3.3 (2.3 - 3.5)	3.4 (2.9 – 4.1)	2.3 (2.3-3.5)
GI in Pasta/LGI group	49.0 (44.0 - 55.1)	46.5 (49.9 – 55.5)	44.0 (42.3 - 49.4)
GI in Higher GI group	62.5 (61.6 - 63.2)	63.3 (60.1 – 64.4)	61.0 (59.2 – 66.6)
Calorie reduction in Pasta/LGI group (kcal) ^e	-179 (-90 - -448)	-165 (-74 - -313)	-447 (-134 - -594)
Calorie reduction in Higher GI group (kcal) ^e	-181 (-93 - -401)	-160 (-40 - -248)	-470 (-172 - -561)
Feeding Control (Met:Supp:DA) (%)	6:44:50	4:48:48	11:33:56
Follow-Up Duration (Weeks)	12 (9-21)	12 (6-24)	12 (10 - 21)
Funding Sources (A:I:AI:NR) (%)	47:9:25:19	44:13:26:17	56:0:22:22

* median (inter quartile range), unless otherwise indicated

^a24/32 trials provided data on sex

^b 30/32 trials reported baseline body weight

^c 28/32 trials reported baseline BMI

^d 11/32 trials provided data from which dose could be approximated

^e 20/32 trials provided data from which to approximate changes in caloric intake

A, agency; AI, agency and industry; BMI, body mass index; C, crossover design; CHD, coronary heart disease; DA, dietary advice; DM, diabetes; GI, glycemic index; I, industry; IP, inpatient; LGI, low glycemic index; Met, metabolic; NR, not reported; OB, obese; OP, outpatient; OW, overweight; P, parallel design; Suppl, supplemented/provision of certain food




Figure 1: Literature Search
40x54mm (300 x 300 DPI)




Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). $n = 2448$. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

40x54mm (300 x 300 DPI)

For peer review only




Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I² statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

40x54mm (300 x 300 DPI)

Online Supplemental Information

Table of Contents

Supplemental Tables..... 4

Supplemental Table S1: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials 4

Supplemental Table S2: PICO framework of the search strategy 6

Supplement Table S3a: Trial characteristics 7

Supplement Table S3b: Trial characteristics continued..... 8

Supplement Table S3c: Trial characteristics continued 9

Supplement Table S3d: Trial characteristics continued..... 10

Supplement Table S3e: Trial characteristics continued..... 11

Supplemental Table S4: Sensitivity analysis of the systematic removal of each trial* 12

Supplemental Table S5: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 for crossover trials 15

Supplement Table S6. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹ 16

Supplemental Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹ 18

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹ 20

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹ 22

Supplementary Table S10. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg). 24

Supplementary Table S11: GRADE assessment of study quality 25

Supplemental Table S12: PRISMA Checklist 28

Supplemental Figures 31

Supplemental Figure S1: Cochrane risk of bias summary for all included trials..... 31

Supplemental Figure S2: Risk of bias proportion graph for all included trials..... 32

Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038) 33

Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285) 34

Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	35
Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)	36
Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)	37
Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448) ..	38
Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)	39
Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	40
Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)	42
Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	44
Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	46
Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) continued (n = 1038)	48
Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	50
Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	52
Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	54
Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	56
Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)	58
Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380). 60	
Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)	62

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19) 63

Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure) 64

Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg) 65

Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²) 66

Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%) 67

Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) 68

Supplemental Tables

Supplemental Table S1: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials

Database	Search Period	Search Terms
Medline	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw. 17. overweight.tw. 18. weight*.tw. 19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
Embase	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw.

		17. overweight.tw. 18. weight*.tw. 19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
The Cochrane Library	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycemic index/ 8. glycaemic ind*.tw. 9. glycemic ind*.tw. 10. glycemic load*.tw. 11. glycaemic load*.tw. 12. exp body weight/ 13. body weight*.tw. 14. BMI.tw. 15. body mass index/ 16. body mass index.tw. 17. weight*.tw. 18. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 19. 12 or 13 or 14 or 15 or 16 or 17 20. 18 and 19 21. limit 20 to animals 22. 20 not 21

Supplemental Table S2: PICO framework of the search strategy

PICO framework ^a defined in the present systematic review and meta-analysis			
Participants	Interventions	Comparators	Outcomes
Adult men and women excluding pregnant or breastfeeding women	Low glycemic index interventions where pasta is included as part of the intervention	Higher glycemic index diets where pasta is not included as part of the intervention	Body weight Body mass index (BMI) Body Fat (%) Waist circumference Waist-to-hip ratio

a Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA and PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 2015; 4:1.

<https://doi.org/10.1186/2046-4053-4-1>

Supplement Table S3a: Trial characteristics

Overweight/Obese Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)§§	Dietary Prescription	Funding Source
Karl et al. 2015 †††	39 (19M:20F)**	OB, OP					2.33	USA	P	Negative+ Neutral	17		CR to 2/3kcal; Metabolic	Agency
Low GI	20		56 (5)*	92.9 (13.6)*	32.3 (3.4)*	42:133						68:15:16		
Higher GI	19		56 (5)*	94 (9.7)*	33.4 (2.6)*	61:201						70:16:14		
Pereira et al. 2015	19 (4M:15F)**	OW, IP/OP					NR	Brazil	P	Neutral	6.4		Ad libitum	Unknown
Low GI	10		28(5)	80.0(12.6)	29.9 (2.1)	41.2(2.2) ¥ **						48.3:16.1:32.8		
Higher GI	9		26(3)	79.1(12.2)	29.1 (2.0)	74.1(2.9) ¥ **						54.6:12.7:34.4		
Buscemi et al. 2013	40 (19M:21F)**	OW/OB, high CVD risk, OP					NR	Italy	P	Negative	12		CR to 20kcal/kg/d; Ad libitum	Unknown
Low GI	19		51 (8)	93.8 (17.3)	34.3 (6.6)	48.1: 138						56:18:26		
Higher GI	21		49 (8)	93.2 (14.4)	34.5 (5.1)	59.3: 174						57:16:27		
Costa et al. 2012	17 (7M:10F)	OW, IP/OP	25.4 (5.8)	84.1(16.3)	26.3(3.2)		NR	Brazil	C	Neutral	4		Ab libitum,2 meals+3 fruit/d provided	NR
Low GI						47.5(3.8)						58.6:13.9:25.5		
Higher GI						61.6(2.8)						55.4:14.2:30.3		
Jebb et al. 2010 - High MUFA	225**	OH, some CV risk factors, OP	~52		~28.5 ⁺		1.75	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	115			83.7 (69.6-93.1)¶¶¶		~55.2						44.6:16.4:35.7****		
Higher GI	110			80.5 (70.0-92.1)¶¶¶		~63.3						44.9:15.3:35.6****		
Jebb et al. 2010 - Low Fat	250**	OH, some CV risk factors, OP	~52		~28.5 ⁺		3.5	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	117			79.4 (70.1- 91.8)¶¶¶		~56.3						41.5:14.2:26.1****		
Higher GI	111			80.7 (71.4- 91.4)¶¶¶		~64.4						41.1:15.7:27.5****		
Larsen et al. 2010-High Pro	231**	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	124		42.1 (6.5)	88.5 (15.6)		~56.5: 108.9						~43:22:32		
Higher GI	107		42 (5.7)	89.5 (17.1)		~61.4: 113.1						~45:23:31		
Larsen et al. 2010 - Low Pro	203 **	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	106		42.2 (5.7)	88.4 (15.7)		~56.2: 121						~51:18:30		
Higher GI	97		42 (5.9)	86.6 (13.8)		~61.6: 137.9						~51:17:31		

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S3b: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Solomon et al. 2010	22 (8M:14F)**	OB, Pre-T2DM, OP					7	USA	P	Neutral	12		Metabolic plus exercise program	Agency
Low GI	10 (3M:7F)		67 (6)	97.4 (12.0)	34.9 (1.1)	39.8 (0.9)								
Higher GI	12 (5M:7F)		64 (3)	94.7 (15.2)	34.1 (1.1)	80.0 (2.1)								
Philippou et al. 2009- 6 mo	38**	≥1 CHD risk factors, OP	(35-65) ¶				NR	UK	P	Negative (for all but n=2)	24		500kcal CR; Ad libitum	Agency
Low GI	22			91.3(14.8)***	29.1 (3.6)***	50.6 (4.6): 114.4(31.5)								
Higher GI	16			97.5(16.4)***	30.5 (3.5)***	63.2(5.6): 175.0(45.6)								
Philippou et al. 2009- 4 mo	42**	OW, OP	(18-65)¶				NR	UK	P	Neutral	16		Ad libitum	Unknown
Low GI	23			87.2 (15.3)	32.5 (4.8)	49.7(5.7):89.7(27.5)								
Higher GI	19			83.6 (13.4)	31.3 (4.8)	63.7(9.4):136.8(56.3)								
Abete et al. 2008	32 (18M:14F)	OB, OP	36(7)				2.33	Spain	P	Negative	8		30% CR; Ad libitum, 3-day menus	Agency
Low GI	16			94.3(16.1)	32.8 (4.3)	(40-45)¶								
Higher GI	16			94.4(13.1)	32.2 (4.4)	(60-65)¶								
Aston et al. 2008	19 (0M:19F)**	OW/OB, OP	51.9(7.6)	87.5 (15)	33.1 (4.9)		3.33	UK	C	Neutral	12		Ad libitum, key CHO foods provided	Agency
Low GI						55.5(3.8): 133.8(22.9)****								
Higher GI						63.9(3): 138.8(30.5)****								
Jensen et al. 2008	44 (0M:44F)**	OW, OP	(20-40)¶				3	Denmark	P	Neutral	10		Ad libitum, partial provision, menu plans	Agency, Industry
Low GI	22 (0M:22F)			77.9(6.9)	27.4 (1.5)	72¥								
Higher GI	22 (0M:22F)			80.2(1.4)	27.6 (0.3)	95¥								
Philippou et al. 2008	13 (5M:8F)**	≥1 CHD risk factors, OP					NR	UK	P	Negative	12		500kcal CR; Ad libitum	Agency
Low GI	7 (3M:4F)		54 (49-58)¶¶	81.5 (4.7)***	28.7 (2.1)***	51.3(51.0-52.0): 105.6 (76.9-110.1)¶¶								
Higher GI	6 (2M:4F)		45 (39-50)¶¶	89.7 (12.8)***	31.5 (4.4)***	59.3(59.2-64.0): 114.7(98.5-134.9)¶¶								

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department of GEZ-LTA Erasmus Hogeschool. All rights reserved. No reuse allowed without permission.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplement Table S3c: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Bellisle et al. 2007	65 (0M:65F)**	OW/OB, OP					NR	France	P	Neutral	12		Ad libitum	Industry
Low GI	35		46.1 (13.6)	80 (13.2)	30.2 (4.1)	na								
Higher GI	30		45.3 (12.0)	79 (13.1)	30.4 (4.4)	na								
de Rougemont et al. 2007	38 (22M:16F)**	OW, OP					2.8	France	P	Neutral	5		Ad libitum, some foods provided	Agency, Industry
Low GI	19		36.3 (8.7)	77.2 (9.6)	27.5 (1.3)	46.5 (1.3)						6 (6):19.8 (1.7):37.7 (4.4)		
Higher GI	19		40.4 (9.6)	77.3 (9.2)	27.2 (1.3)	66.3 (2.6)						6 (6):19.8 (1.7):38.4 (3.1)		
Sichieri et al. 2007	123 (0M:123F)**	OW, OP					NR	Brazil	P	Negative	72		100-300kcal CR; 6-d menu and exchange lists provided	Agency
Low GI	63		37.2 (5.4)*	67.7 (6.6)*	na	21(38): 74(84)						6 (6.3): 13.3: 27.2(4.6)		
Higher GI	60		37.5 (5.6)*	68.5 (7.5)*	na	51(28): 199(43)						6 (6.2): 12.3: 26.1(4.7)		
McMillian-Price et al. 2006-High Carb	64(16M:48F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	32		30.5 (7.9)	87.1 (15.3)	30.6 (4.5)	45 (6):89 (28)						6 (6):19 (0):22 (6)		
Higher GI	32		31.8 (9.6)	86 (10.7)	30.9 (3.4)	70 (6):129 (45)						6 (6):18 (6):19 (6)		
McMillian-Price et al. 2006-High Protein	65(15M:50F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	33		34.6 (8.6)	88.4 (17.2)	32.1 (5.2)	44 (6):59 (23)						6 (11):26 (6):28 (6)		
Higher GI	32		30.2 (8.5)	87.7 (16.4)	31.3 (4.5)	59 (6):75 (17)						6 (6):28 (6):27 (6)		
Wolever et al. 2002	24 (5M:19F)**	IGT, OP					NR	Canada	P	Neutral	16		Ad libitum, partial provision	Agency
Low GI	13(3M:10F)		55.2 (10.8)	79.7 (13.1)***	29.7 (4.3)	54.4 (2.5):91.8 (9.4)						6 (6.1):19.4 (1.8):24.7 (5.8)		
Higher GI	11(2M:9F)		58.8 (13.3)	76.4 (20.4)***	29.3 (7.3)	59.3 (2.0):96.8 (11.6)						6 (6.6):17.4 (2.3):27.9 (6.3)		

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S3d: Trial characteristics continued

Diabetes Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Jenkins et al. 2014	141(77M:64F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum, bread supplement	Industry Association
Low GI	70 (38M:32F)		59 (10)	85 (20)	30 (5)	~51:53						~38.5:19.8:37.2		
Higher GI	71 (39M:32F)		59 (10)	84 (19)	31 (6)	~62:89						~49.2:19.8:27.4		
Visek et al. 2014	20 (12M:8F)	T2DM, OP	62.7 (5.8)	91.9 (14.1)	32 (4.2)		NR	Czech Republic	C	Neutral	12		Ad libitum	Agency
Low GI						49 (48-51)¶¶						~37.2:18.0:36.0		
Higher GI						68 (61-72)¶¶						~36.2:17.3:40.0		
Jenkins et al. 2012	121 (61M: 60F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum	Agency
Low GI	60		58 (10.1)	85.6 (20.1)	31.4 (7.0)	47: 80						45.4:22.8:30.5		
Higher GI	61		61 (7.8)	82.5 (17.2)	29.9 (5.5)	58: 100						48.3:21.4:28.5		
Yusof et al. 2009	100**	T2DM, OP	NR				NR	Malaysia	P	Neutral	12		Ad libitum, key foods provided to lowGI group	Agency
Low GI	51			69.12 (13.33)	27.05 (4.91)	57(6): 108(32)						52(4):18(3):30(4)		
Higher GI	49			66.83 (11.50)	26.79 (4.65)	64(5): 131(30)						54(4):17(3):28(5)		
Jenkins et al 2008	210 (125M:82F)	T2DM, OP					NR	Canada	P	Neutral	24		Ad libitum	Agency
Low GI	106 (65M:41F)		60 (10)	87.0 (20.0)	30.6 (6.0)	49.4: 91.5						44.0:21.2:33.3		
Higher GI	104 (63M:41F)		61 (9)	87.8 (19.4)	31.2 (5.8)	59.3: 117.9						47.5:20.7:30.5		
Wolever et al. 2008	103	T2DM, OW/OB, OP					NR	Canada	P	Neutral	52		Ad libitum, key foods provided	Agency
Low GI	55		60.6 (7.5)*	81.1 (18.7)*	31.6 (4.5)*	55.1 (3.0): 133 (14.8)						46.7:20.6(3.0):26.5 (5.9)		
Higher GI	48		60.4 (7.9)*	84.4(18.0)*	30.1 (4.3)*	63.2 (2.8): 135 (20.8)						46.5:20.4 (2.8):30.8 (4.8)		

Supplement Table S3e: Trial characteristics continued

Diabetes Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance§	Duration (wks)	Dietary Composition % (SD)§§	Dietary Prescription	Funding Source
Jimenez-Cruz et al. 2003	14 (6M:8F)**	T2DM, OP	59 (34)											
Low GI				91.6 (24.3)	32.4 (6.0)	44(3.4): 86(19.8)		NR	Mexico	C	Neutral	60:21:23	Ad libitum	Industry
Higher GI				92.6 (25.4)	32.3 (6.0)	56(4.9): 139(27.3)						64:18:20		
Heilbronn et al. 2002	45 (23M:22F)**	T2DM, OW, OP			NR			3.5	Australia	P	Negative		CR to 1500kcal/d; Ad libitum, partial provision	Unknown
Low GI	24 (11M:13F)		56.0(9.4)	91.7(16.2)		43						58.9(2.9):22.2 (1.5):17.9 (3.9)		
Higher GI	21 (12M:9F)		57.5(9.6)	93.2 (13.3)		75						60.3(2.3):21.7 (0.9):17.1 (2.3)		
Fontvieille et al. 1992	18 (12M:6F)	T1DM/T2DM, OP	47.2(11.6)	NR	24.8(2.6)			4.7	France	C	Neutral		Ad libitum	Agency, Industry
Low GI						38.1(5.3)						58.2(2.2):18.0(2.5):36.2(6.8)		
Higher GI						64.2(3.1)						67.3(2.3):18.8(1.6):36.3(6.0)		
Fontvieille et al. 1988	8 (4M:4F)	T1DM, OP	43.5 (9.9)	NR	24.1 (6.8)			3.5	France	C	Neutral		Ad libitum, partial provision	Agency, Industry
Low GI						46.5(2.5)						58.1(1.5):17.4 (1.4):35.0 (2.8)		
Higher GI						60.1 (5.1)						62.5(2.5):16.9 (1.7):36.0 (2.8)		
CHD Trial														
Frost et al. 2004	55 (48M:7F)**	CHD, OP						NR	UK	P	Neutral §§§		Ad Libitum	Unknown
Low GI	26 (23M:3F)		63.6 (9.4)	81.2 (12.2)	26.9 (3.3)	50(4):115(39)						59 (5):18 (5):31 (5)		
Higher GI	29 (25M:4F)		61.8 (9)	81.7 (16.7)	28.7 (4.6)	57(4):106(34)						62 (5):18 (5):32 (10)		

Abbreviations: BMI=Body Mass Index; C=Crossover design; Carb=Carbohydrate; CED= carbohydrate exchange diet; CHD= Coronary Heart Disease; CR= calorie restriction; CV=cardiovascular; GI= glycemic index; GL=glycemic load; IGT= impaired glucose tolerance; IP=inpatient; IR= insulin resistant; High Carb= high carbohydrate; HI= hyperinsulinemic; M = male; mo=months; MUFA=monounsaturated fatty acids; na=not available; T1DM=type 1 diabetes mellitus; T2DM = type 2 diabetes mellitus; OB= obese; OH=otherwise healthy; OP = outpatient; OW= overweight; P=Parallel design; SD= standard deviation; F = female; UK= United Kingdom; USA= United States of America; wks=weeks; y= years

§Negative energy balance diets were designed for weight loss, Neutral energy balance diets for weight maintenance, where all diets are isocaloric between test and control groups; §§Energy from carbohydrate: protein: fat for the planned diet or if not planned; §§§ Participants were advised to lose weight if they had a BMI>28kg/m2.

¶ Range of values; ¶¶ median and interquartile range (IQR); ¶¶¶ Actual GI/GL or if not available, planned;

* Calculated before dropout; **Completer Analysis, as used in data analysis; *** completer analysis, data obtained from authors; **** based on less participants than in analysis;

+ approximate based on all study arms; ¥ approximate based on test meals; ‡ approximate based on n=45 from Sloan et al. 2004, the original publication of this study;

‡‡‡ analysis includes weight loss and weight maintenance phases

Supplemental Table S4: Sensitivity analysis of the systematic removal of each trial*

	MD [95% CI], P-value I ² , P-value					
	Body Weight (kg) N=32	BMI (kg/m ²) N=18	Body Fat (%) N=10	Waist Circumference (cm) N=18	Waist-to-hip Ratio N=6	Sagittal Abdominal Diameter (cm) N=3
All studies	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.01 [-0.58, 0.56], P=0.98 65%, P=0.003	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92
Removal of:						
OW/OB						
Pereira et al. 2015	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.54	-0.24 [-0.34, -0.14], P<0.01 0.00%, P=0.93	0.19 [-0.25, 0.63], P=0.39 36%, P=0.13	-0.37 [-1.04, 0.31], P=0.28 62%, P<0.01	0.00 [-0.01, 0.01], P=0.67 0.00%, P=0.58	n/a
Karl et al. 2015	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.50	-0.25 [-0.35, -0.16], P<0.01 0.00%, P=0.91	0.10 [-0.51, 0.71], P=0.74 64%, P<0.01	n/a	n/a	n/a
Buscemi et al. 2013	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.86	-0.08 [-0.63, 0.48], P=0.79 64%, P<0.01	-0.46 [-1.06, 0.15], P=0.14 64%, P<0.01	n/a	n/a
Costa et al. 2012	-0.62 [-0.84, -0.41], P<0.01 0.00%, P=0.46	-0.26 [-0.37, -0.16], P<0.01 0.00%, P=0.86	-0.06 [-0.73, 0.61], P=0.86 66%, P<0.01	-0.33 [-0.94, 0.28], P=0.29 59%, P<0.01	n/a	n/a
Jebb et al. 2010 - LowFat	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - LowPro	-0.60 [-0.81, -0.38], P<0.01 0.00%, P=0.57	n/a	n/a	-0.45 [-1.09, 0.18], P=0.16 64%, P<0.01	n/a	-0.05 [-0.39, 0.29], P=0.79 0.00%, P=0.86
Jebb et al. 2010- HighMUFA	-0.64 [-0.85, -0.43], P<0.01 0.00%, P=0.65	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - HighPro	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.48	n/a	n/a	-0.37 [-1.00, 0.26], P=0.26 62%, P<0.01	n/a	-0.13 [-0.45, 0.19], P=0.42 0.00%, P=0.92
Solomon et al. 2010	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	n/a	n/a	n/a	n/a

	0.00%, P=0.47	0.00%, P=0.88				
Philippou et al. 2009 - 6mnth	-0.65 [-0.86, -0.44], P<0.01 0.00%, P=0.59	-0.28 [-0.37, -0.18], P<0.01 0.00%, P=0.97	-0.12 [-0.71, 0.48], P=0.70 65%, P<0.01	-0.55 [-1.15, 0.04], P=0.07 61%, P<0.01	n/a	n/a
Philippou et al. 2009 - 4mnth	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.89	-0.09 [-0.72, 0.54], P=0.78 66%, P<0.01	-0.44 [-1.08, 0.19], P=0.17 64%, P<0.01	n/a	n/a
Abete et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a	n/a
Philippou et al. 2008	-0.62 [-0.83, -0.42], P<0.01 0.00%, P=0.48	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.89	-0.06 [-0.68, 0.55], P=0.84 68%, P<0.01	-0.41 [-1.02, 0.19], P=0.18 63%, P<0.01	n/a	n/a
Aston et al. 2008	-0.66 [-0.87, -0.44], P<0.01 0.00%, P=0.52	n/a	-0.00 [-0.70, 0.69], P=0.99 68%, P<0.01	-0.54 [-1.14, 0.07], P=0.08 62%, P<0.01	n/a	n/a
Jensen et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.27 [-0.36, -0.17], P<0.01 0.00%, P=0.88	n/a	-0.44 [-1.05, 0.18], P=0.16 64%, P<0.01	0.00 [-0.01, 0.00], P=0.24 0.00%, P=0.41	-0.09 [-0.35, 0.18], P=0.51 0.00%, P=0.69
de Rougemont et al. 2007	-0.57 [-0.80, -0.34], P<0.01 0.00%, P=0.53	-0.25 [-0.36, -0.14], P<0.01 0.00%, P=0.87	0.06 [-0.57, 0.70], P=0.84 67%, P<0.01	n/a	n/a	n/a
Sichieri et al. 2007	-0.66 [-0.88, -0.45], P<0.01 0.00%, P=0.53	n/a	n/a	n/a	n/a	n/a
Bellisle et al. 2007	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.88	n/a	-0.47 [-1.08, 0.13], P=0.13 64%, P<0.01	0.00 [-0.01, 0.00], P=0.14 0.00%, P=0.70	n/a
McMillan-Price et al. 2006 - HighCHO	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.51	n/a	n/a	-0.38 [-1.01, 0.25], P=0.23 63%, P<0.01	n/a	n/a
McMillan-Price et al. 2006 - HighPro	-0.70 [-0.91, -0.49], P<0.01 0.00%, P=0.91	n/a	n/a	-0.62 [-1.19, -0.05], P=0.03 55%, P<0.01	n/a	n/a
Wolever et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Diabetes:						
Vissek et al. 2014	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	0.01 [-0.57, 0.60], P=0.96	n/a	n/a	n/a

	0.00%, P=0.46	0.00%, P=0.86	68%, P<0.01			
Jenkins et al. 2014	-0.66 [-0.88, -0.43], P<0.01 0.00%, P=0.47	-0.29 [-0.39, -0.18], P<0.01 0.00%, P=0.94	n/a	-0.61 [-1.18, -0.04], P=0.04 50%, P=0.01	0.00 [-0.01, 0.00], P=0.21 0.00%, P=0.43	n/a
Jenkins et al. 2012	-0.62 [-0.84, -0.40], P<0.01 0.00%, P=0.46	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.87	n/a	-0.44 [-1.05, 0.17], P=0.16 64%, P<0.01	0.00 [-0.01, 0.01], P=0.64 0.00%, P=0.53	n/a
Yusof et al. 2009	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.87	n/a	-0.33 [-0.95, 0.28], P=0.29 58%, P<0.01	n/a	n/a
Jenkins et al. 2008	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.36, -0.15], P<0.01 0.00%, P=0.87	n/a	n/a	n/a	n/a
Wolever et al. 2008	-0.64 [-0.84, -0.43], P<0.01 0.00%, P=0.57	n/a	n/a	-0.50 [-1.10, 0.09], P=0.10 62%, P<0.01	n/a	n/a
Jimenez-Cruz et al. 2003	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.86	n/a	n/a	n/a	n/a
Heilbronn et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a	n/a
Fontvieille et al. 1992	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Fontvielle et al. 1988	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
CHD						
Frost et al. 2004	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.17], P<0.01 0.00%, P=0.89	n/a	-0.48 [-1.09, 0.12], P=0.12 63%, P<0.01	0.00 [-0.01, 0.00], P=0.25 1%, P=0.40	n/a

*Sensitivity analysis included the removal of each single study from the meta-analyses one at a time and the summary effect was recalculated. An influential outlier was considered a study whose removal changed the magnitude of the pooled effect by >10%.

BMI, body mass index; CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; MD, mean difference; mnt, month; MUFA, monounsaturated fatty acids; n/a, not applicable; OB, obese; OW, overweight; Pro, protein

Supplemental Table S5: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 in crossover trials

Outcome (no. crossover trials/total)	MD (95% CI), P-value I ² , P-value		
	Correlation Coefficient used in the Primary Analysis	Correlation coefficient used in Sensitivity Analyses	
	0.5	0.25	0.75
Body Weight, kg (6*/32)	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.43], P<0.01 0.00%, P=0.51
BMI, kg/m ² (3/18)	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.90	-0.26 [-0.35, -0.17], P<0.01 0.00%, P=0.90
Body Fat, % (3*/10)	-0.01 [-0.58, 0.56], P=0.98 65%, P<0.01	-0.00 [-0.58, 0.58], P=0.99 64%, P<0.01	-0.02 [-0.57, 0.54], P=0.96 66%, P<0.01
Waist Circumference, cm (2*/18)	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	-0.44 [-1.04, 0.15], P=0.14 60%, P<0.01	-0.46 [-1.07, 0.14], P=0.13 65%, P<0.01
Waist-to-hip Ratio (0/6)	-0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	n/a	n/a
Sagittal Abdominal Diameter, cm (0/6)	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92	n/a	n/a

* one of these crossover trials, however, did not require the use of a correlation coefficient as complete data was available

BMI, body mass index; CI, confidence interval; MD, mean difference; no., number

Supplement Table S6. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	24.1 – 37.1 kg/m ²	32	2448	-0.04 (-0.13, 0.06)	0.00%	0.454
Follow-up	3 – 72 wks	32	2448	0.01 (-0.01, 0.02)	0.00%	0.314
Dose Pasta	1.75 – 7.0 serv/wk	11	740	0.29 (-0.91, 1.50)	0.00%	0.595
GI ³	21 - 72	31	2383	-0.01 (-0.04, 0.02)	0.45%	0.482
Difference in GI ²	4.9 - 40.2	31	2383	-0.00 (-0.04, 0.03)	2.75%	0.811
Fiber ³	8.0 - 39.4g/d	27	1851	0.02 (-0.01, 0.06)	0.00%	0.221
Change in Fiber ⁴	-7.5 - +13.8g/d	17	1571	-0.02 (-0.07, 0.04)	23.13%	0.595
Difference in Fiber ²	-8.5 - +15.3g/d	27	1851	-0.01 (-0.06, 0.04)	0.00%	0.701
Saturated Fat ³	5.1 - 12.5%	14	1396	-0.05 (-0.30, 0.20)	34.32%	0.648
Change in Saturated Fat ⁴	-8.2 - -1.2%	12	1309	-0.01 (-0.29, 0.28)	46.27%	0.965
Difference in Saturated Fat ²	-2.0 - +2.3%	14	1396	-0.34 (-1.08, 0.39)	32.80%	0.329
CHO ³	37.2 - 68.0%	30	2345	-0.01 (-0.05, 0.03)	0.00%	0.541
Change in CHO ⁴	-7.2 - +10.1%	19	2046	-0.07 (-0.12, -0.01)	0.00%	0.016
Difference in CHO ²	-11.1 - +5.4%	30	2345	0.02 (-0.05, 0.10)	0.00%	0.547
Protein ³	13.3 - 26.1%	30	2345	0.04 (-0.04, 0.12)	0.00%	0.332

Change in Protein ⁴	-0.5 - +8.6%	19	2046	0.15 (0.03, 0.27)	0.00%	0.017
Difference in Protein ²	-2.5 - +3.4%	30	2345	-0.18 (-0.38, 0.02)	0.00%	0.069
Fat ³	16.0 - 37.7%	30	2345	0.01 (-0.05, 0.06)	0.00%	0.852
Change in Fat ⁴	-12.2 - +5.4%	19	2046	0.05 (-0.02, 0.12)	17.73%	0.122
Difference in Fat ²	-4.4 - +10.6%	30	2345	0.02 (-0.05, 0.09)	0.00%	0.610

- 1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body weight on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body weight. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.
- 2 Difference in diet variable between the intervention and control arms
- 3 Intake at the end of study in the intervention arm
- 4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index

Supplemental Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 37.1 kg/m ²	18	1038	-0.01 (-0.06, 0.03)	0.00%	0.559
Follow-up	4 – 24 wks	18	1038	0.00 (-0.01, 0.02)	0.00%	0.559
Dose Pasta*	2.33 – 7.0 serv/wk	4	143			
GI ³	39.8 - 72	17	973	0.01 (-0.01, 0.03)	0.00%	0.153
Difference in GI ²	7.0 - 40.2	17	973	-0.01 (-0.03, 0.01)	0.00%	0.275
Fiber ³	8.0 - 39.4g/d	16	935	0.01 (-0.01, 0.02)	0.00%	0.366
Change in Fiber ⁴	-1.8 - +12.4g/d	10	758	0.00 (-0.03, 0.03)	0.00%	0.846
Difference in Fiber ²	-4.9 - +13.0g/d	16	935	-0.00 (-0.02, 0.02)	0.00%	0.821
Saturated Fat* ³	7.6 - 12.5%	7				
Change in Saturated Fat* ⁴	-2.4 - -1.2%	6				
Difference in Saturated Fat* ²	-1.0 - +2.3%	7				
CHO ³	37.2 - 68.0%	16	935	-0.00 (-0.02, 0.01)	0.00%	0.594
Change in CHO ⁴	-5.6 - +3.2%	10	758	-0.02 (-0.06, 0.02)	0.00%	0.325
Difference in CHO ²	-11.1 - +2.0%	16	935	-0.01 (-0.03, 0.02)	0.00%	0.531
Protein ³	13.9 – 22.8%	16	935	0.00 (-0.04, 0.04)	0.00%	0.905

Change in Protein ⁴	-0.2 - +3.0%	10	758	0.01 (-0.11, 0.14)	0.00%	0.786
Difference in Protein ²	-2.5 - +3.4%	16	935	-0.03 (-0.12, 0.06)	0.00%	0.441
Fat ³	16.0 - 37.7%	16	935	0.00 (-0.02, 0.02)	0.00%	0.717
Change in Fat ⁴	-4.8 - +5.4%	10	758	0.02 (-0.02, 0.06)	0.00%	0.240
Difference in Fat ²	-4.4 - +10.6%	16	935	0.01 (-0.01, 0.04)	0.00%	0.299

*For Dose, there were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in BMI on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in BMI. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	10	285	-0.02 (-0.26, 0.23)	67.69%	0.890
Follow-up	4 – 24 wks	10	285	0.06 (-0.06, 0.17)	66.00%	0.303
Dose Pasta*	2.33 - 3.33 serv/wk	3	96			
GI ³	39.8 - 72	10	285	0.12 (-0.00, 0.25)	52.21%	0.053
Difference in GI ²	7.0 - 40.2	10	285	-0.09 (-0.15, -0.03)	19.39%	0.008
Fiber* ³	8.0 - 39.4g/d	8	228			
Change in Fiber* ⁴	-1.8 - +12.4g/d	4	87			
Difference in Fiber* ²	-4.9 - +13.0g/d	8	228			
Saturated Fat* ³	7.6 - 12.5%	3	93			
Change in Saturated Fat* ⁴	-2.4 - -1.2%	2	51			
Difference in Saturated Fat* ²	-1.0 - +2.3%	3	93			
CHO* ³	37.2 - 68.0%	9	247			
Change in CHO* ⁴	-5.6 - +3.2%	4	87			
Difference in CHO* ²	-11.1 - +2.0%	9	247			
Protein* ³	13.9 – 22.8%	9	247			

Change in Protein* ⁴	-0.2 - +3.0%	4	87		
Difference in Protein* ²	-2.5 - +3.4%	9	247		
Fat* ³	16.0 - 37.7%	9	247		
Change in Fat* ⁴	-4.8 - +5.4%	4	87		
Difference in Fat* ²	-4.4 - +10.6%	9	247		

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body fat on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body fat. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	18	1380	0.10 (-0.13, 0.32)	59.02%	0.372
Follow-up	4 – 52 wks	18	1380	0.05 (-0.03, 0.13)	60.50%	0.225
Dose Pasta*	3.0 – 3.33 serv/wk	2	63			
GI ³	41.2 – 72.0	17	1315	-0.01 (-0.12, 0.10)	66.00%	0.841
Difference in GI ²	4.9 – 32.9	17	1315	-0.02 (-0.10, 0.06)	64.51%	0.596
Fiber ³	8.0 - 39.4g/d	15	1258	0.05 (-0.05, 0.14)	59.60%	0.342
Change in Fiber ⁴	-1.8 - +13.8g/d	13	1176	0.01 (-0.17, 0.20)	70.59%	0.861
Difference in Fiber ²	-4.9 - +15.3g/d	15	1258	-0.00 (-0.14, 0.13)	64.86%	0.939
Saturated Fat* ³	6.1 – 10.1%	8	604			
Change in Saturated Fat* ⁴	-7.6 - -1.2%	7	562			
Difference in Saturated Fat* ²	-2.0 - +2.3%	8	604			
CHO ³	38.5 - 58.6%	16	1277	-0.11 (-0.19, -0.04)	27.06%	0.007
Change in CHO ⁴	-7.2 - +10.1%	13	1176	-0.09 (-0.23, 0.04)	57.19%	0.148
Difference in CHO ²	-11.1 - +5.4%	16	1277	-0.01 (-0.18, 0.17)	60.81%	0.947
Protein ³	13.9 – 26.1%	16	1277	0.20 (0.01, 0.38)	43.92%	0.038

Change in Protein ⁴	-0.3 - +8.6%	13	1176	0.19 (-0.09, 0.46)	63.38%	0.165
Difference in Protein ²	-2.5 - +3.4%	16	1277	-0.14 (-0.66, 0.38)	62.79%	0.570
Fat ³	22.0 - 37.2%	16	1277	0.10 (-0.05, 0.26)	53.48%	0.185
Change in Fat ⁴	-12.2 - +5.4%	13	1176	0.08 (-0.07, 0.24)	61.15%	0.274
Difference in Fat ²	-4.4 - +10.6%	16	1277	0.09 (-0.06, 0.25)	47.17%	0.220

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in waist circumference on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in waist circumference. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplementary Table S10. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg)

Dose threshold of servings of pasta consumed per week	Dose ranges of servings of pasta consumed per week	β (95% CIs) *	Residual I^2 †	p -value
3.0	≤ 3.0	-0.70 (-3.27, 1.86)	0.00%	0.890
	> 3.0	0.91 (-0.89, 2.70)		
3.33	≤ 3.33	0.05 (-1.80, 1.89)	0.00%	0.518
	> 3.33	0.44 (-1.75, 2.63)		
3.5	≤ 3.5	0.09 (-1.65, 1.82)	0.00%	0.888
	> 3.5	0.46 (-1.89, 2.81)		

* β is the slope derived from the piecewise linear meta-regression analyses and represents the treatment effect on body weight for doses above and below each dose-threshold representing servings of pasta consumption per week. † The residual I^2 value indicates heterogeneity unexplained by each dose-threshold.

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department of Health Technology, Innovation and Society, University of Cambridge. All rights reserved.

Supplementary Table S11: GRADE assessment of study quality

Quality assessment*							№ of patients		Effect	Quality Importance
№ of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)	
Body Weight (follow up: median 12 weeks)										
32	randomised trials	not serious	not serious	serious ^a	not serious	none	1294	1228	MD -0.63 kg (-0.84 to -0.42)	⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
BMI (follow up: median 12 weeks)										
18	randomised trials	not serious	not serious	serious ^a	not serious	none	551	538	MD -0.26 kg/m² (-0.36 to -0.16)	⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Waist Circumference (follow up: median 12 weeks)										
18	randomised trials	not serious	serious ^b	serious ^a	not serious	none	717	624	MD -0.46 cm (-0.05 to 0.14)	⊕⊕○○ LOW ^{a,b} Due to downgrade for inconsistency and indirectness

Quality assessment*							№ of patients		Effect	Quality Importance	
№ of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)		
Body Fat (follow up: median 12 weeks)										12 April 2018. Downloaded from http://bmjopen.bmj.com/ on June 5, 2025 at Department of Clinical Epidemiology, AI training, and similar technologies.	
10	randomised trials	not serious	serious ^c	serious ^a	not serious	none	176	165	MD -0.01 % (-0.58 to 0.56)		⊕⊕○○ LOW ^{a,c} Due to downgrade for inconsistency and indirectness
Waist-to-hip Ratio (follow up: median 12 weeks)											
6	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	223	222	MD -0.00 (-0.01 to 0.00)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Sagittal Abdominal Diameter (follow up: median 26 weeks)										Erasmus University Medical Center Rotterdam	
3	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	237	214	MD -0.09 cm (-0.34 to 0.16)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

CI, Confidence interval; GI, glycemic index; LGI, low glycemic index; MD, Mean difference

*All outcomes started with high quality evidence since all studies were randomized controlled trials. Risk of Bias –We rated down for risk of bias if the majority of studies were considered to be at high risk of bias. Inconsistency – We assessed inconsistency using I^2 estimates where an I^2 of 50% or higher indicates substantial heterogeneity. I^2 is the percentage of variability in the treatment estimates that is attributable to heterogeneity between studies. We rated down for inconsistency if there was substantial heterogeneity that was unexplained by any a priori sensitivity or subgroup analyses. Indirectness – We rated down for indirectness if there were factors present relating to the population, interventions, and outcomes that limited the generalizability of the results. Imprecision – We rated down for imprecision if the 95% confidence interval (95% CI) crossed the minimally important difference (MID) for harm. MIDs used for each outcome are: 0.5kg for body weight (based on Johnston et al. JAMA 2014;312(9):923-33); 0.2m² for BMI; 2.0cm for waist circumference; 2.0% for body fat; 0.02 for waist-to-hip ratio; 2.0cm for sagittal abdominal diameter.

- a. Downgrade for serious indirectness, since most of the available trials did not quantify the amount of food consumed in the context of the low-GI dietary patterns
- b. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2=62\%$, P -heterogeneity<0.001), which could not be explained
- c. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2= 65\%$, P -heterogeneity=0.003), which could not be explained
- d. No downgrade for publication bias, as publication bias could not be assessed due to lack of power for assessing funnel plot asymmetry and small study effects (<10 trials included in the meta-analysis)

-2017-019438 on 2 April 2019. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool. All rights reserved. No reuse allowed without permission. For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Supplemental Table S12: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5-6, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1

Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6, Supplemental Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	7-9

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7-8
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	9-11, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10-11, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	11, Supplemental Figures S1-S2

Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	11-12, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	11-12, Figures 2-3, Supplemental Figures S3-S7
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	11, Supplemental Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	12-15, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policymakers).	16-19
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	19-21
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	21-22
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	25-26


From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Supplemental Figures


	Random Sequence Generation	Allocation Concealment	Blinding of Participants, Personnel and Outcome Assessment	Incomplete Outcome Data	Selective Reporting		Random Sequence Generation	Allocation Concealment	Blinding of Participants, Personnel and Outcome Assessment	Incomplete Outcome Data	Selective Reporting
Abete et al. 2008	?	?	?	+	+	Jimenez-Cruz et al. 2003	?	?	?	?	+
Aston et al. 2008	+	?	+	+	+	Karl et al. 2015	+	+	+	+	+
Bellisle et al. 2007	?	?	?	?	+	Larsen et al. 2010 -LowPro	+	+	+	+	+
Buscemi et al. 2012	+	+	+	?	+	Larsen et al. 2010 -HighPro	+	+	+	+	+
Cost et al. 2012	?	?	+	+	+	McMillan-Price et al. 2006- HighCHO	?	?	+	+	+
de Rougemont et al. 2007	?	+	+	+	+	McMillan-Price et al. 2006- HighPro	?	?	+	+	+
Fontvielle et al. 1992	?	?	+	+	+	Pereira et al. 2015	?	?	+	?	+
Fontvielle et al. 1988	?	?	+	+	+	Philippou et al. 2008	?	?	+	?	+
Frost et al. 2004	?	?	+	+	+	Philippou et al. 2009-4mo	?	?	+	?	+
Heilbronn et al. 2002	?	?	+	?	+	Philippou et al. 2009-6mo	?	?	?	?	+
Jebb et al. 2010 - HighMUFA	+	?	+	?	+	Sicheri et al. 2007	+	?	+	?	+
Jebb et al. 2010 - LowFat	+	?	+	?	+	Solomon et al. 2010	?	?	+	+	+
Jenkins et al. 2014	+	+	+	+	+	Visek et al. 2014	?	?	+	+	+
Jenkins et al. 2012	+	+	+	+	+	Wolever et al. 2008	+	+	?	+	+
Jenkins et al. 2008	+	+	+	+	+	Wolever et al. 2002	+	+	+	+	+
Jensen et al. 2008	?	?	+	+	+	Yusof et al. 2009	+	?	+	+	+

Supplemental Figure S1: Cochrane risk of bias summary for all included trials
Summary of risk of bias ratings for each individual study included in the meta-analysis.



Supplemental Figure S2: Risk of bias proportion graph for all included trials


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P<0.10$ and quantified with the use of the I^2 statistic where $I^2\geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


BMI, body mass index; CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot’s bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran’s Q statistic at a significance level of P<0.10 and quantified with the use of the I² statistic where I²≥50% is considered to be evidence of substantial heterogeneity and ≥75% considerable heterogeneity.


CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance fixed effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.



Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids



Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and it indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.

BMI, body mass index; btw, between; CHD, coronary heart disease; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; serv, serving; trt, treatment; wk, week.


*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.0 kg (-0.08, 0.65) (1 vs. 2) to 1.32kg (-4.28, 6.91) (1 vs. 3) to -1.29kg (-6.91, 4.32) (2 vs. 3).

† data available on 31 studies

‖ data available on 17 studies

‡ data available on 27 studies

-2017-019438 on 24 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for use in text and data mining, AI training, and similar technologies.




Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and it indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.

BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); trt, treatment.
§ data available on 30 studies
¥ data available on 19 studies
¶ data available on 14 studies
Ψ data available on 12 studies

2017-019438 on 23 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For copyright, including for text and data mining, AI training, and similar technologies.




Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p = 0.05$ indicated that the effect size differed between levels of the subgroup.

LRB=Low Risk of Bias, URB= Unclear Risk of Bias, HRB=High Risk of Bias. *Within and between subgroup analysis could not be performed against HRB since no values for HRB subgroup. ** no values for URB subgroup. § Blinding of Participants, Personnel, and Outcome Assessors.

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I² was estimated with the use of the Cochran's Q statistic and indicates the between study

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

heterogeneity unexplained by the variability in response because of the specific between studies factor. 0.05 indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=1 trials), therefore analyses were not performed.

*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.05kg/m² (-0.156, 0.274) (1 vs. 2) to 0.584kg/m² (-1.009, 2.178) (1 vs. 3) to -0.525kg/m² (-2.122, 1.072) (2 vs. 3).


† data available on 17 studies

‡ data available on 16 studies

|| data available on 10 studies

BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) continued (n = 1038)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=7 trials), saturated fat within treatment (n=6 trials) and saturated fat between treatments (n=7 trials), therefore analyses were not performed.

§ data available on 16 studies
¥ data available on 10 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.

2017-019438 on 24 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool. For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml



Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the degree of heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup.


*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.

BMI, body mass index; CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias.

2017-019438 on 24 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml




Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=9), test carbohydrate, protein and fat (n=9), carbohydrate, protein and fat within treatment (n=4), carbohydrate, protein and fat between treatment (n=9), test fibre (n=8), fibre within treatment (n=4), fibre between treatment (n=8), test saturated fat (n=3 trials), saturated fat within treatment (n=2 trials) and saturated fat between treatments (n=3 trials), therefore analyses were not performed.

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For copyright, including for uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup.


*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.

CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=2 trials), therefore analyses were not performed.


*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.15cm (-1.492, 1.818) (1 vs. 2) to 1.608cm (-3.432, 6.648) (1 vs. 3) to -1.445cm (-6.630, 3.740) (2 vs. 3).

† data available on 17 studies

‡ data available on 15 studies

BMI, body mass index; btw, between; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

2017-019438 on 20 April 2015. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For copyright, including for text and data mining, AI training, and similar technologies.



Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's I^2 statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=8 trials), saturated fat within treatment (n=7 trials) and saturated fat between treatments (n=8 trials), therefore analyses were not performed.

§ data available on 16 studies
¥ data available on 13 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.

2017-019438 on 20 April 2015. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For copyright, including for text and data mining, AI training, and similar technologies.



Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's I^2 statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup.


*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.


CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias

2017-019438 on 24 April 2015. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.




Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)


CHO, carbohydrate; MD, mean difference



Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)
MD, mean difference




Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure)




Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.




Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.

Supplemental Table S12: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5-6, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and	6, Supplemental

		simplifications made.	Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	7-9

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7-8
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	9-11, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10-11, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	11, Supplemental Figures S1-S2
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	11-12, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	11-12, Figures 2-3, Supplemental Figures S3-S7
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see item 15).	11, Supplemental

			Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	12-15, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	16-19
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	19-21
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	21-22
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	25-26

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097
For more information, visit: www.prisma-statement.org.

BMJ Open

Effect of Pasta in the Context of Low Glycemic Index Dietary Patterns on Body Weight and Markers of Adiposity: A Systematic Review and Meta-analysis of Randomized Controlled Trials in Adults

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019438.R1
Article Type:	Research
Date Submitted by the Author:	08-Dec-2017
Complete List of Authors:	Chiavaroli, Laura; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit; University of Toronto, Nutritional Sciences Kendall, Cyril; University of Toronto, Nutritional Sciences Braunstein, Catherine; University of Toronto, Nutritional Sciences; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit Blanco Mejia, Sonia; Clinical Nutrition and Risk Factor Modification Centre, St. Michael's Hospital; Toronto 3D Knowledge Synthesis and Clinical Trials Unit, St. Michael's Hospital Leiter, Lawrence; St. Michael's Hospital, Endocrinology; University of Toronto, Nutritional Sciences Jenkins, David; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Center; University of Toronto, Nutritional Sciences Sievenpiper, John; McMaster University, Department of Pathology and Molecular Medicine; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Centre
Primary Subject Heading:	Nutrition and metabolism
Secondary Subject Heading:	Cardiovascular medicine
Keywords:	body weight, pasta, glycemic index, glycaemic index, systematic review and meta-analysis, weight loss

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Effect of Pasta in the Context of Low Glycemic Index Dietary Patterns on Body Weight and Markers of Adiposity: A Systematic Review and Meta-analysis of Randomized Controlled Trials in Adults

Laura Chiavaroli^{1,2}, Cyril WC Kendall^{1,2,3}, Catherine R Braunstein^{1,2}, Sonia Blanco Mejia^{1,2}, Lawrence A Leiter^{1,2,4-6}, David JA Jenkins^{1,2,4-6}, John L Sievenpiper^{1,2,4-6}

- 1 Department of Nutritional Sciences, University of Toronto, Toronto, Canada
- 2 Toronto 3D Knowledge Synthesis and Clinical Trials Unit, St. Michael’s Hospital, Toronto, Canada
- 3 College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, Canada
- 4 Department of Medicine, University of Toronto, Toronto, Canada
- 5 Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Toronto, Canada
- 6 Division of Endocrinology and Metabolism, St. Michael’s Hospital, Toronto, Canada

Corresponding Author:
Dr. John L Sievenpiper, MD, PhD, FRCPC, St. Michael’s Hospital, #6137-61 Queen Street East, Toronto, ON, M5C 2T2, CANADA
Telephone: +1 416 867 7475; Fax: +1 416 867 7495; Email: john.sievenpiper@utoronto.ca

Word Count: 4,968
Tables: 1
Figures: 3
Online only supplementary material included

Keywords: body weight, pasta, glycemic index, glycaemic index, systematic review and meta-analysis, weight loss

ABSTRACT

Objective: Carbohydrates have been implicated in the obesity epidemic. To assess the effect of pasta alone or pasta in the context of low glycemic index (GI) dietary patterns on adiposity, we conducted a systematic review and meta-analysis.

Design: Systematic review and meta-analysis of randomized controlled trials (RCTs) with GRADE assessment.

Eligibility criteria for selecting studies: MEDLINE, Embase, CINAHL, and the Cochrane Library were searched through 07 February 2017. We included RCTs ≥ 3 -weeks assessing the effect of pasta alone or in the context of low-GI dietary patterns on measures of global (body weight, BMI, body fat) and regional (waist circumference [WC], waist-to-hip ratio [WHR], sagittal abdominal diameter [SAD]) adiposity in adults. Two independent reviewers extracted data and assessed risk of bias. Data were pooled using the generic inverse-variance method and expressed as mean differences (MDs) with 95% confidence intervals (95% CIs). Heterogeneity was assessed (Cochran Q statistic) and quantified (I^2 -statistic). GRADE assessed the overall certainty of the evidence.

Results: We identified no trials of the effect of pasta alone and 32 trial comparisons (n=2,448 participants) of the effect of pasta in the context of low-GI dietary patterns. Pasta in the context of low-GI dietary patterns significantly reduced body weight (MD=-0.63kg [95% CIs, -0.84, -0.42kg]) and BMI (MD=-0.26kg/m² [95% CIs, -0.36, -0.16kg/m²]) compared with higher-GI dietary patterns. There was no effect on other measures of adiposity. The overall certainty of the

evidence was graded as moderate for body weight, BMI, WHR, and SAD and low for WC and body fat.

Conclusions: Available RCTs demonstrate that pasta in the context of low-GI dietary patterns does not adversely affect adiposity and even reduces body weight and BMI compared to higher GI dietary patterns. Future trials should assess the effect of pasta in the context of other healthy dietary patterns.

Protocol registration: ClinicalTrials.gov Identifier, NCT02961088

Strengths and limitations of this study

- The present systematic review and meta-analysis was based on a comprehensive search and includes a large number of randomized controlled trials which provide the best protection against bias.
- We used the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) system to evaluate the strength and quality of the evidence.
- There was evidence of unexplained inconsistency in the intervention estimates across trials for waist circumference and body fat.
- The generalizability of our results is questionable with evidence of indirectness in the pooled estimates for all body and adiposity outcomes, as all of the trials assessed pasta in the context of low-GI dietary patterns (no trials assessed the effect of pasta alone or in the

74 context of other dietary patterns) and most of the available trials did not quantify the
75 amount of pasta consumed.

For peer review only

INTRODUCTION

As the role of saturated fat in chronic disease has been called into question, carbohydrates have come under attack in the media^{1,2}, popular books³⁻⁹, statements of health advocacy groups¹⁰ and commentaries in leading medical journals^{11,12}. Much of the attention has focused on sugars but traditional carbohydrate staples like pasta, rice, and breads are increasingly being implicated in the epidemics of overweight and obesity^{1,7}. Although advantages for weight related outcomes have been shown for dietary patterns that are high in these foods but low in glycemic index (GI)^{13,14}, high in whole grains^{15,16}, and/or high in dietary fibre^{17,18}, there has been a general lack of recognition of the importance of carbohydrate quality.

Pasta is an important example of a food which is considered a refined carbohydrate but has a low-GI, a property that has been exploited extensively in studies of low-GI dietary patterns. It remains unclear whether pasta alone or in the context of a low-GI dietary pattern contributes to weight gain^{1,7}. We undertook a systematic review and meta-analysis of randomized controlled trials (RCTs) with a full GRADE assessment to quantify the effect of pasta alone or in the context of low-GI dietary patterns on body weight and measures of adiposity relevant to the prevention and management of overweight and obesity.

METHODS

Design

Our protocol followed the guideline of the Cochrane Handbook for Systematic Reviews of Interventions¹⁹, and findings are reported according to the guidelines of the Preferred Reporting

Items for Systematic Reviews and Meta-Analyses²⁰ (**Supplemental Table S1**). The protocol is registered at clinicaltrials.gov (identifier, NCT02961088).

Data sources and searches

We searched MEDLINE (<http://www.nlm.nih.gov/bsd/pmresources.html>), Embase (<https://www.embase.com>), CINAHL (<https://health.ebsco.com/products/the-cinahl-database>), and the Cochrane Library (<http://www.cochranelibrary.com/>) from inception through 07 February 2017. The full search terms used in this study are presented in **Supplemental Tables S2-S3**. Briefly, we searched using variations of the terms pasta and glycemic index and glycemic load and body weight and BMI. The search was limited to human studies and had no language restrictions. Reference lists of selected studies and reviews were also searched to identify additional articles.

Study selection

We include RCTs that investigated the effect of pasta consumed alone or in the context of low-GI dietary patterns that emphasized pasta in comparison with higher GI diets that did not include pasta on body weight or other measures of global (BMI, body fat) or abdominal (waist circumference, waist-to-hip ratio, sagittal abdominal diameter, or visceral adipose tissue assessed by imaging modalities) adiposity in participants of all health backgrounds. Trials were included if the intervention arm assessed the effect of pasta consumed alone or assessed the effect of a low GI diet which emphasized pasta as part of the low GI dietary advice. Trials were excluded if they had diet duration of <3 weeks, did not intend to use a calorie- and macronutrient-matched comparator arm that was higher in GI; included pregnant or breast-feeding women or children; or

121 did not provide suitable end-point data. When multiple publications existed for the same study,
122 the article with the most information was included (n=6). Published abstracts were not included.

123
124 **Data extraction**

125 Two reviewers (LC, CB) assessed the titles and abstracts of all identified studies and
126 independently reviewed and extracted relevant data from each report, including study design,
127 blinding, sample size, participant characteristics, follow-up duration, identification of pasta in the
128 low-GI diet only, comparator diet, macronutrient profile, funding source, and outcome data. The
129 primary outcome was body weight, and secondary outcomes included markers of global (BMI,
130 body fat) and abdominal (waist circumference, waist-to-hip ratio, sagittal abdominal diameter, or
131 visceral adipose tissue assessed by imaging modalities) adiposity. Mean±SD differences between
132 test and control arms were extracted for each outcome.

133 In those trials where the data were included in figures and not provided numerically, we used the
134 software program Plot Digitizer (<http://plotdigitizer.sourceforge.net/>) to extract the data.

135 Additional information was requested from the authors of all included trials. Disagreement were
136 resolved by consensus or where necessary by a third author (SBM).

137
138 **Risk of bias assessment**

139 Risk of bias for each individual study was assessed using the Cochrane Risk of Bias tool¹⁹. The
140 level of bias was evaluated for sequence generation, allocation concealment, blinding,
141 incomplete outcome data, and selective reporting and determined overall as either low (proper
142 methods taken to reduce bias), high (improper methods creating bias), or unclear (insufficient
143 information provided to determine the bias level).

Statistical analysis

Baseline, end-of-study, and between-treatment differences in body weight, BMI, waist circumference, body fat, waist-to-hip ratio, sagittal abdominal diameter or visceral adipose tissue assessed by imaging modalities were recorded as means $\pm$ SDs. If not provided, between-treatment differences in change-from-baseline or end differences were calculated by subtracting means and variance measures such as SEs were imputed with the use of published formulas¹⁹. Missing SDs were imputed with the use of the pooled SD from other studies included in the analysis¹⁹.

Data analyses were conducted using Review Manager version 5.3 (RevMan) (Copenhagen, Denmark: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014) for primary analyses and Stata version 13 (College Station, TX: StataCorp LP) for subgroup analyses. A generic inverse-variance method with random-effects models was used to calculate pooled mean differences and 95% confidence intervals (CIs). Random-effects models were used even in the absence of statistically significant inter-study heterogeneity, as they yield more conservative summary effect estimates in the presence of residual heterogeneity. Change-from-baseline differences were preferred over end differences and paired analyses were applied to all crossover trials with the use of a within-individual correlation coefficient between treatments of 0.5 as described by Elbourne et al.²¹

Inter-study heterogeneity was assessed by the Cochran Q statistic, where $P < 0.10$ was considered statistically significant, and quantified by the I^2 statistic, where $I^2 \geq 50\%$ indicates substantial heterogeneity¹⁹. Sensitivity analysis were performed which included the removal of each single study from the meta-analyses one at a time and recalculation of the summary effect. An

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

167 influential study was considered a study whose removal changed the magnitude of the pooled
168 effect by >10%. Sensitivity analysis were also conducted using different correlation coefficient
169 values for crossover trials (0.25 and 0.75) to test for the robustness of the effect size, conducting
170 analyses using fixed effects models and restricting analyses to those trials for which pasta intake
171 could be quantified.

172 If ≥ 10 trial comparisons were available, then sources of heterogeneity were explored by
173 subgroup analyses. A priori categorical subgroups analyses were assessed by meta-regression
174 analyses. These included patient type (normal body weight, overweight or obese [average
175 baseline BMI $>27\text{kg/m}^2$]), diabetes, coronary heart disease), follow-up (<24 -weeks, ≥ 24 weeks),
176 baseline BMI (BMI ≤ 30 , $>30\text{kg/m}^2$), design (parallel, crossover), energy balance (negative on
177 both arms [weight loss diets], neutral on both arms [weight maintaining diets]) and dose of pasta
178 (based on the median). A priori categorical subgroup analyses also included the following dietary
179 factors: GI (absolute level [≤ 55 , >55 ; glucose scale], within-treatment change, between-treatment
180 change), fat intake (absolute level [$<30\%$, $\geq 30\%$ energy], within-treatment change, between-
181 treatment change), carbohydrate intake (absolute level [$<50\%$, $\geq 50\%$ energy], within-treatment
182 change, between-treatment change), protein intake (absolute level [$<20\%$, $\geq 20\%$ energy], within-
183 treatment change, between-treatment change), dietary fibre intake (absolute level [$<28\text{g/day}$,
184 $\geq 28\text{g/day}$], within-treatment change, between-treatment change), and risk of bias. A priori
185 continuous meta-regression analyses were conducted on the absolute levels and within- and
186 between-treatment changes of these same dietary factors in the intervention arms of pasta in the
187 context of low-GI dietary patterns. Linear and non-linear pasta intake dose-response analyses
188 were assessed by using continuous meta-regression analyses and spline curve modeling
189 (MKSPLINE procedure), respectively. Publication bias was assessed by visual inspection of

funnel plots and the Egger²² and Begg²³ tests, when ≥ 10 trial comparisons were available. If publication bias was suspected, then adjustment for funnel plot asymmetry was done by imputing missing study data using the Duval and Tweedie trim and fill method²⁴.

Grading the evidence

The grading of recommendations assessment, development, and evaluation (GRADE) approach was used to assess the certainty of the evidence²⁵. Evidence was graded as high, moderate, low or very low quality. The included RCTs were graded as high quality evidence by default and downgraded based on pre-specified criteria. Criteria to downgrade evidence included risk of bias (weight of studies show risk of bias assessed by the Cochrane Risk of Bias tool), inconsistency (substantial unexplained heterogeneity, $I^2 > 50\%$, $P < 0.10$), indirectness (presence of factors that limited the generalizability of the results), imprecision (the 95% CI for effect estimates were wide or crossed pre-specified minimally important differences [MIDs] for harm), and publication bias (significant evidence of small-study effects).

Patient involvement

No patients were directly involved in the development of the research question, selection of the outcome measures, design and implementation of the study, or interpretation of the results.

RESULTS

Search results

Figure 1 shows the flow of the literature. We identified 4876 reports of which 29 met eligibility criteria. No reports were identified that assessed the effect of pasta alone, while 29 reports (including 32 trial comparisons involving 2448 participants) were identified that assessed the effect of pasta in the context of low-GI dietary patterns on any adiposity outcome in adults²⁶⁻⁵⁴. Of the 32 trial comparisons that assessed the effect of pasta in the context of a low-GI dietary pattern, there were 32 trial comparisons for body weight, 18 trial comparisons for BMI^{27,28,31-33,35,36,39-41,43-46,48,49,52,53}, 18 trial comparisons for waist circumference^{27,28,31,33,34,36,38-40,42,44-47,52,53}, 10 trial comparisons for body fat^{27,28,31-33,36,38,41,43,53}, 6 trial comparisons for waist-to-hip ratio^{31,39,40,44,45,52}, and 3 trial comparisons for sagittal abdominal diameter^{34,39}. There was only one trial comparison identified³⁵ for visceral adipose tissue assessed by imaging modalities, thus a meta-analysis could not be undertaken for this outcome.

Trial characteristics

Table 1 and **Supplemental Table S4** show the characteristics of all included trials of the effect of pasta in the context of low-GI dietary patterns. The majority of trials had a parallel design (26/32) with a median follow-up of 12 weeks (IQR: 9–21 weeks) and a median number of participants per trial of 43 (IQR: 21–112). Most participants were middle aged (median age: 50 y; IQR: 40-58 y) men and women (median ratio of men to women: 0.4:1 in available trials). The median baseline BMI across studies was 30.4kg/m² (IQR: 28.2–32.0). Regarding metabolic phenotype, 21 (66%) trials included participants who were overweight or obese (had a baseline BMI≥27kg/m²), 10 (31%) had diabetes and one trial (3%) with coronary heart disease (CHD). We did not retrieve any trials where participants had a normal BMI at baseline (≤25kg/m²),

although 6 trials did not include BMI >25 kg/m² as part of criteria, the average baseline BMI was ≥ 27 kg/m², therefore categorized as overweight.

Risk of bias

Supplemental Figures S1 and S2 show the individual Cochrane Risk of Bias tool assessments for each of the included trials of the effect of pasta in the context of low-GI dietary patterns. No serious risk of bias was detected.

Effect of Pasta in the Context of Low-GI dietary patterns and Body Weight

Figure 2 shows the effect of pasta in the context of low-GI dietary patterns on the primary outcome body weight. Pooled analyses showed pasta in the context of low-GI dietary patterns had the effect of reducing body weight by -0.63 kg (95% CI: -0.84, -0.42 kg; $P < 0.001$) compared with higher GI control diets with no evidence of heterogeneity ($I^2 = 0\%$, P -heterogeneity = 0.51).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Global Adiposity

Figure 3 and Supplemental Figure S3-S4 show the effect of pasta in the context of low-GI dietary patterns on markers of global adiposity. Pooled analyses showed that pasta in the context of low-GI dietary patterns had the effect of reducing BMI by -0.26 kg/m² ($n = 18$ trials; MD = -0.26 kg/m²; 95% CI: -0.36, -0.16 kg/m²; $P < 0.001$) compared with higher GI control diets with no evidence of heterogeneity ($I^2 = 0\%$, P -heterogeneity = 0.90). There was no effect on body fat ($n = 10$ trials; MD = -0.01% , 95% CI: -0.58, 0.56%; $P = 0.98$) with evidence of substantial heterogeneity ($I^2 = 65\%$, P -heterogeneity < 0.01).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Abdominal Adiposity

Figure 3 and Supplemental Figures S5-S7 show the pooled estimates for the markers of abdominal adiposity. Pooled analyses did not show a significant effect of pasta in the context of low-GI dietary patterns compared with higher-GI control diets on waist circumference (n=18 trials; MD=-0.46cm, 95% CI:-1.05, 0.14cm; P=0.13), waist-to-hip ratio (n=6 trials; MD=-0.00, 95% CI:-0.01, 0.00; P=0.27), or sagittal abdominal diameter (n=3 trials, MD=-0.09cm, 95% CI:-0.34, 0.16cm; P=0.48) There was only evidence of substantial heterogeneity in the analyses for waist circumference ($I^2=62\%$, P-heterogeneity<0.01).

Sensitivity analyses

We conducted four sets of sensitivity analyses (**Supplemental Tables S5-6, Supplemental Figures S8-9**). The systematic removal of each trial did not modify the direction or significance of the effect estimates or the evidence of heterogeneity for any of the outcomes with the exception of waist circumference (**Supplemental Table S5**). In the sensitivity analysis for waist circumference, two studies were influential studies in that their removal altered the magnitude of the pooled effect in the remaining studies by >10%, where the removal of the studies of McMillan-Price et al. (high protein comparison)⁴² and Jenkins et al.⁴⁴ rendered the results for waist circumference statistically significant (MD= -0.62cm; 95% CI: -1.19, -0.05; P=0.03) and (MD= -0.61cm; 95% CI: -1.18, -0.04; P=0.04), respectively; forest plots not shown). Heterogeneity remained significant in both cases ($I^2= 55\%$, P-heterogeneity<0.01 and $I^2=50\%$, P-heterogeneity=0.01, respectively). Sensitivity analyses using correlation coefficients of 0.25 and 0.75 for paired analyses of crossover trials also did not modify the results (**Supplemental Table**

S6). In the sensitivity analyses where fixed effects models were applied (Supplemental Figure S8), the direction, magnitude and significance of the pooled estimates were very similar to those produced by the random effects models with the exception of the sensitivity analysis for waist circumference which was significant (MD= -0.62; 95% CI: -0.93, -0.32; P<0.001). Finally, restricting analyses to the 11 trials for which pasta intake could be quantified (median pasta intake, 3.33 servings/week [range, 1.75 – 7 servings/week]) showed a similar reduction in body weight (MD= -0.70 kg; 95% CI: -1.10, -0.29 kg; P<0.001) when pasta was consumed in the context of low-GI dietary patterns compared with the higher GI control arms without evidence of heterogeneity ($I^2=0\%$, P-heterogeneity=0.68) (Supplemental Figure S9).

Subgroup analyses

We were only able to conduct a priori categorical and continuous subgroup analyses for body weight, BMI, body fat and waist circumference. Subgroup analyses for waist-to-hip ratio and sagittal abdominal diameter could not be assessed, owing to <10 trial comparisons in each case. Supplemental Figures S10-S12 show the categorical a priori subgroup analyses for body weight. There was no evidence of significant effect modification in any of the subgroup analyses for body weight, including no effect modification of follow-up when comparing studies less than 24 weeks duration to those greater than or equal to 24 weeks (-0.63kg vs -0.57kg, respectively) (Supplemental Figure S10). Neither was there evidence of significant effect modification in any of the subgroup analyses for BMI, body fat or waist circumference (Supplemental Figures S13-20).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplemental Table S7 and **Supplemental Figures S21-22** show the continuous subgroup analyses for body weight. There was evidence of significant effect modification by carbohydrate and protein intake, where an increase in carbohydrate intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight loss ($\beta = -0.07$, 95% CI: -0.12, -0.01, $I^2 = 0.00\%$, $P = 0.02$), and an increase in protein intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight gain ($\beta = 0.15$, 95% CI: 0.03, 0.27, $I^2 = 0.00\%$, $P = 0.02$). None of the other continuous subgroup analyses were significant. There was no evidence of significant effect modification in any of the continuous subgroup analyses for BMI (**Supplemental Table S8**). For body fat, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of difference in GI between intervention and control groups, where greater difference in GI between the groups was associated with greater reduction in body fat in the intervention group ($\beta = -0.09$, 95% CI: -0.15, -0.03, $I^2 = 19.39\%$, $P = 0.01$) (**Supplemental Table S9**). None of the other continuous subgroup analyses were significant. For waist circumference, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of absolute carbohydrate level and absolute protein level, where greater carbohydrate level in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with greater loss in waist circumference ($\beta = -0.11$, 95% CI: -0.19, -0.04, $I^2 = 27.06\%$, $P < 0.01$) and a lower protein level in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with an increase in waist circumference ($\beta = 0.20$, 95% CI: 0.01, 0.38, $I^2 = 43.92\%$, $P = 0.04$) (**Supplemental Table S10**). None of the other continuous subgroup analyses were significant.

Dose-response analyses

Supplemental Tables S7 and S11 and Supplemental Figure S23 show the dose-response analysis for the 11 trials for which pasta intake could be quantified. No evidence of a linear dose response was seen for pasta intake by meta-regression analyses (Supplemental Table S7). There was also no evidence of a non-linear dose response by MKSPLINE ($P=0.85$) (Supplemental Figure S23) or piecewise linear meta-regression analyses (Supplemental Table S7).

Publication Bias

Supplemental Figures S24-S27 shows the funnel plots for body weight, BMI, body fat and waist circumference. There was no evidence of funnel-plot asymmetry. Formal testing with the Egger and Begg tests did not show evidence of small-study effects ($P>0.05$ for both). Publication bias was not assessed for waist-to-hip ratio and sagittal abdominal diameter, owing to <10 trial comparisons.

GRADE Assessment

Supplemental Table S12 shows a summary of the GRADE assessments for the effect of pasta in the context of low-GI dietary patterns on body weight and measures of adiposity. The evidence was graded as moderate for body weight, BMI, waist-to-hip ratio and sagittal abdominal diameter owing to downgrades for indirectness and low for waist circumference and body fat, owing to downgrades for indirectness and inconsistency ($I^2=59\%$, P -heterogeneity <0.001 ; $I^2=66\%$, P -heterogeneity <0.01 , respectively).

DISCUSSION

The present systematic review and meta-analysis quantified the effect of pasta alone and pasta in the context of low-GI dietary patterns on body weight and other markers of adiposity. We identified no trial comparisons for the effect of pasta alone and 32 trial comparisons involving 2448 participants who were predominantly middle-aged and overweight or obese for the effect of pasta in the context of low-GI dietary patterns. The primary pooled analysis demonstrated that pasta in the context of low-GI dietary patterns did not contribute to weight gain, resulting in a significant weight loss of -0.63kg when compared to diets higher in GI over a median follow-up of 12 weeks. The lack of harm was reflected in the established clinical secondary outcome measures of global (BMI and body fat) and abdominal adiposity (waist circumference, waist-to-hip ratio and sagittal abdominal diameter). These findings were robust across subgroups. The findings did not differ by metabolic phenotype in those who were overweight or obese or had diabetes, which is noteworthy since these are populations who would benefit from weight management strategies. There was also no effect modification by the energy balance of the design such that the weight loss was seen even under conditions of neutral energy balance (in which participants were instructed to consume dietary advice ad libitum), suggesting that encouragement of the consumption of pasta in the context of a low-GI dietary patterns does not cause harm and may even lead to spontaneous weight loss. There was also no effect modification by follow-up either in continuous meta-regression or categorical, where the 24 trials with <24weeks follow-up had a weight reduction similar to those 8 trials with ≥24weeks follow-up (-0.63kg vs -0.57kg, respectively). This is of relevance since many dietary studies are successful in demonstrating weight loss in the short term but not over the long term.

Findings in the context of existing studies

We are not aware of any RCTs directly assessing the effect of pasta intake on health parameters including body weight. Our findings, however, agree with earlier systematic reviews and meta-analyses of RCTs of the effect of low-GI dietary patterns irrespective of pasta intake on body weight and adiposity. The systematic review and meta-analysis by Thomas et al. in 2007 found a significant -1.1kg weight loss and -1.3kg/m² reduction in BMI favouring low-GI or glycemic load (GL) diets compared to control diets in 6 RCTs of 5 weeks to 6 months in duration in overweight or obese individuals¹³. Another systematic review and meta-analysis by Schwingshackl et al. in 2013 found a -0.62kg weight loss favouring low-GI/GL diets compared to higher GI/GL diets in 14 RCTs with greater than 6 months duration in overweight individuals (BMI>25kg/m²)¹⁴.

Our findings also agree with trials in which pasta was emphasized in the context of other healthy dietary patterns. One trial done in children in Spain given Mediterranean dietary advice which included increasing the intake of pasta found that approximately 11.3% of the participants in the Mediterranean diet group who were classified as overweight and obese changed their weight status to normal weight compared to only approximately 2.6% of the participants in the control group⁵⁵.

Other lines of evidence from observational studies have demonstrated benefits of pasta consumption on body weight and adiposity. Pasta intake was recently assessed in the Moli-sani study and the Italian Nutrition & HEalth Survey (INHES), a cross-sectional study of over 20,000 Italians from all over Italy⁵⁶. The study demonstrated that higher pasta consumption was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

394 associated with lower BMI, waist circumference and waist-to-hip ratio and with a lower
395 prevalence of overweight and obesity⁵⁶. Furthermore, greater pasta intake was associated with
396 better adherence to the Mediterranean diet, a dietary pattern which has a demonstrated
397 cardiovascular benefit⁵⁷. Similar associations between greater pasta intake and lower body
398 weight have also been observed in prospective U.S. cohorts⁵⁸. A pooled analysis of the 3
399 Harvard cohorts also showed that higher GI diets (which would preclude pasta) were associated
400 with weight gain⁵⁹.

401
402 Although the product form of pasta can vary widely, including in shape (e.g. macaroni, spaghetti,
403 linguine), ingredients (e.g. type of wheat, egg content), and processing technique (e.g. drying
404 temperature), studies have demonstrated that when comparing pastas varying in these
405 parameters, despite slight variations in glycemic response among pastas, glycemic responses are
406 still lower compared to a control, e.g. white bread^{60,61}. One concern of the choice of pasta as a
407 carbohydrate food is that it is a refined food low in fibre. Although there are whole grain pasta
408 options available, studies have demonstrated that fiber added to pasta, does not significantly
409 affect the glucose or insulin response, the secretion of gut hormones or satiety^{62,63}. Furthermore,
410 pasta has a similar glycemic response compared to many fiber-rich carbohydrates, including
411 barley, legumes and steel cut oats, and still a lower glycemic response compared to other fiber-
412 rich foods including whole wheat bread, breakfast cereals like bran flakes and potatoes with
413 skin⁶⁴. The typically consumed white wheat pasta also has a higher micronutrient content
414 compared to other white wheat products like bread since it contains the aleurone layer which is
415 preserved as a result of the use of harder wheats (durum wheat); even when durum wheats are
416 used in breads, pasta retains a lower glycemic response primarily because of the processing

techniques used in pasta making which give pasta a compact structure and reduced starch hydrolysis⁶¹.

The mechanism by which pasta in the context of low-GI dietary patterns lead to weight loss even under conditions of ad libitum dietary advice is unclear. Lower GI diets may result in greater body weight reduction compared to higher GI diets because lower GI foods may be more satiating⁶⁵ and have been shown to delay hunger and decrease subsequent energy intake¹³. Low-GI dietary patterns are also characterized by high fiber content^{64,66} which may also contribute to improvements in satiety and hunger¹⁷. Furthermore, studies which have compared ad libitum low-GI dietary patterns to standard energy-restricted low-fat diets, demonstrated equivalency or better weight loss when following the low-GI diet, despite the fact that they could eat as much as they desired^{13,67}. Thus, voluntary energy intake may be lower after low-GI meals, as has been previously demonstrated⁶⁸.

Strengths and limitations

The strengths of the present systematic review and meta-analysis include that it is comprehensive, includes RCTs which protects against bias and uses GRADE to evaluate the quality of evidence. Additionally, a large number of trials were identified (32 trials) for the primary outcome of body weight, the median follow-up period was 12 weeks which allows for the assessment of a moderate duration of intervention, none of the trials were rated as having a serious risk of bias, and there was no evidence of publication bias.

1
2
3 439 There are several limitations. First, we downgraded the certainty of the evidence for serious
4
5 440 inconsistency in the treatment estimates across trials for some of the outcomes assessed. There
6
7 441 was evidence of unexplained heterogeneity in waist circumference ($I^2=62\%$) and in body fat
8
9 442 ($I^2=65\%$). Although the inconsistency in these outcomes may have related to measurement
10
11 443 error⁶⁹ in the different techniques for measuring waist circumference and body fat, we were
12
13 444 unable to conduct sensitivity or subgroup analyses to explore this source of heterogeneity.
14
15 445 Second, we downgraded the certainty of the evidence for serious indirectness. Most of the
16
17 446 available trials did not quantify the amount of pasta consumed in the context of the low-GI
18
19 447 dietary patterns. Although sensitivity analyses in which analyses were restricted to the 11 trials
20
21 448 that did quantify (providing a median 3.33 servings/week) pasta intake did not meaningfully alter
22
23 449 our estimates (-0.70kg versus -0.63kg), it is difficult to quantify the effect of pasta in these diets.
24
25 450 There is also the question of indirectness in the translation to other background diets. None of the
26
27 451 available trials evaluated the effect of pasta alone or in the context of other dietary patterns.
28
29 452 Whether the observed effect of pasta in the context of low-GI dietary patterns will hold in the
30
31 453 context of other healthy dietary patterns, such as Mediterranean and Vegetarian dietary patterns,
32
33 454 is unclear. Although there is no biological reason to doubt that the findings would hold across
34
35 455 different dietary patterns, there was no direct evidence to support this conclusion. If the question
36
37 456 had been asked from the perspective of benefit as opposed to that of harm, then the relatively
38
39 457 short duration of the included trials is another reason to downgrade for serious indirectness. In
40
41 458 the absence of long-term trials (>1 year diet duration), it is difficult to conclude with certainty
42
43 459 that the observed lack of harm implies an actual sustainable benefit. Finally, there was some
44
45 460 evidence of imprecision for benefit. Whereas the 95% CI of the pooled estimates did not overlap
46
47 461 with our pre-specified MID for harm (that is, they did not contain evidence for harm) and so
48
49
50
51
52
53
54
55
56
57
58
59
60

were not downgraded for imprecision, the upper bound of the 95% CI did overlap with the lower bound of the same MID to assess the precision of the evidence for benefit for some outcomes.

Balancing these strengths and limitations, GRADE assessed the overall quality and strength of the available evidence of the effect of pasta in the context of low-GI dietary patterns as moderate for the primary outcome of body weight and the secondary outcomes of BMI, waist-to-hip ratio and sagittal abdominal diameter owing to downgrades for indirectness. The evidence was assessed as low for the other secondary outcomes of body fat and waist circumference owing to downgrades for indirectness and inconsistency.

Implications

These results are important considering the negative messages directed at the public regarding carbohydrates, which is influencing their food choices, as is evident in recent reductions in carbohydrate intake⁷⁰⁻⁷², and in particular reductions in pasta consumption^{70,73-76}. Contrary to these concerns, the available evidence shows that when pasta is consumed in the context of low-GI dietary patterns that there is not weight gain but rather marginally clinically significant weight loss ($>0.5\text{kg}$)⁷⁷.

Although we were able to approximate the amount of pasta consumed in one third of included trials, it is unclear what the effect of pasta would be in the context of other dietary patterns. Low-GI dietary patterns, however, shares many similarities with a Mediterranean diet, which emphasizes many low-GI foods and has demonstrated a CVD benefit⁵⁷.

1
2
3 485 Current clinical practice guidelines already suggest the replacement of high GI foods with low-
4
5 486 GI foods for improvement of glycemic control and cardiovascular risk factors^{78,79}. The present
6
7 487 evidence means that pasta may be highlighted as an important example of a low-GI food which
8
9
10 488 can contribute to a low-GI dietary pattern, a pattern which in turn may potentially improve
11
12 489 cardio-metabolic risk without an adverse effect on weight control.
13
14
15 490

16
17 491 **CONCLUSIONS**
18

19 492 In conclusion, the available evidence from RCTs does not allow us to conclude that pasta
20
21 493 consumed in the context of low-GI dietary patterns has an adverse effect on body weight and
22
23 494 adiposity outcomes of importance in the prevention and management of overweight and obesity.
24
25 495 On the contrary, pasta in the context of low-GI dietary patterns reduces body weight and BMI
26
27 496 compared with higher-GI dietary patterns. The results are generalizable in the context of a high
28
29 497 carbohydrate dietary pattern composed of low-GI foods with or without the intention of weight
30
31 498 loss in middle-aged individuals who are overweight or obese or have diabetes. Although the
32
33 499 clinical significance of the observed weight loss is debatable, this finding increases our
34
35 500 confidence that pasta in the context of low-GI dietary patterns does not result in weight gain.
36
37 501 Further research may change our confidence in the estimates for our primary outcome body
38
39 502 weight and several key secondary outcomes including BMI and two measures of abdominal
40
41 503 adiposity, waist-to-hip ratio and sagittal abdominal diameter. More research is needed, to
42
43 504 improve our estimates for the secondary outcomes, body fat and waist circumference and assess
44
45 505 whether our findings extend to related cardio-metabolic outcomes. There is also a need for more
46
47 506 randomized trials of >1 year diet duration to clarify whether the lack of harm for pasta in the
48
49
50
51
52
53 507 context of low-GI dietary patterns will translate into meaningful long-term benefits. Other
54
55
56
57
58
59
60

randomized trials should focus on whether pasta will have similar effects in the context of other healthy dietary patterns such as a Mediterranean diet.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

Figure 1: Literature Search

Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). n = 2448. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.
CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.
BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

538 Acknowledgements

539 We wish to thank Teruko Kishibe, an information specialist in the Scotiabank Health Sciences
540 Library, St Michael's Hospital, for her help in the development of search terms used. Aspects of
541 this work were presented at the Società Italiana di Nutrizione Umana (SINU) Congress, in
542 Parma, Italy, on April 12, 2017 and at the 35th International Symposium on Diabetes and
543 Nutrition, of the Diabetes and Nutrition Study Group (DNSG) of the European Association for
544 the Study of Diabetes (EASD) in Skagen, Denmark, on June 20 and 21, 2017.

545 Data Sharing

546 No additional data available.

548 Exclusive License

549 The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf
550 of all authors, an exclusive license (or non-exclusive for government employees) on a worldwide
551 basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in BMJ
552 editions and any other BMJ PGL products and sublicenses such use and exploit all subsidiary
553 rights, as set out in our license.

554 Ethics Approval

555 Not required.

556 Authorship:

557 All authors had full access to all of the data (including statistical reports and tables) in this study
558 and take full responsibility for the integrity of the data and the accuracy of the data analysis.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conception and design: C.W.C. Kendall, J.L. Sievenpiper.

Analysis and interpretation of the data: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Drafting of the article: L. Chiavaroli, J.L. Sievenpiper.

Critical revision of the article for important intellectual content: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Final approval of the article: L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

Obtaining of funding: C.W.C. Kendall, D.J.A. Jenkins, J.L. Sievenpiper.

Administrative, technical, or logistic support: S. Blanco Mejia.

Collection and assembly of data: L. Chiavaroli, C.R. Braunstein, S. Blanco Mejia.

Guarantor: J.L. Sievenpiper.

Transparency declaration: As guarantor, J.L. Sievenpiper affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Funding:

This work was funded by the Canadian Institutes of Health Research (funding reference number, 129920) through the Canada-wide Human Nutrition Trialists' Network (NTN). The Diet, Digestive tract, and Disease (3-D) Centre, funded through the Canada Foundation for Innovation (CFI) and the Ministry of Research and Innovation's Ontario Research Fund (ORF),

provided the infrastructure for the conduct of this project. **L Chiavaroli** was funded by a Toronto 3D Postdoctoral Fellowship Award. **DJA Jenkins** was funded by the Government of Canada through the Canada Research Chair Endowment. **JL Sievenpiper** was funded by a PSI Graham Farquharson Knowledge Translation Fellowship, Canadian Diabetes Association (CDA) Clinician Scientist award, CIHR INMD/CNS New Investigator Partnership Prize, and Banting & Best Diabetes Centre Sun Life Financial New Investigator Award. None of the sponsors had a role in any aspect of the present study, including design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, approval of the manuscript or decision to publish.

Conflicts of Interest:

All authors have completed the Unified Competing Interest form (available on request from the corresponding author) and declare: **CWC Kendall** has received research support from the Advanced Foods and Material Network, Agrifoods and Agriculture Canada, the Almond Board of California, the American Pistachio Growers, Barilla, the California Strawberry Commission, the Calorie Control Council, CIHR, the Canola Council of Canada, the Coca-Cola Company (investigator initiated, unrestricted grant), Hain Celestial, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Kraft, Loblaw Companies Ltd., Orafiti, Pulse Canada, Saskatchewan Pulse Growers, Solae and Unilever. He has received travel funding, consultant fees and/or honoraria from Abbott Laboratories, the Almond Board of California, the American Peanut Council, the American Pistachio Growers, Barilla, Bayer, the Canola Council of Canada, the Coca-Cola Company, Danone, General Mills, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Loblaw Companies Ltd., the Nutrition Foundation of Italy, Oldways Preservation Trust, Orafiti, Paramount Farms, the Peanut Institute,

1
2
3 604 PepsiCo, Pulse Canada, Sabra Dipping Co., Saskatchewan Pulse Growers, Solae, Sun-Maid, Tate
4
5 605 and Lyle, and Unilever. He is on the Clinical Practice Guidelines Expert Committee for Nutrition
6
7 606 Therapy of the European Association for the Study of Diabetes (EASD) and has served on the
8
9
10 607 scientific advisory board for the Almond Board of California, the International Tree Nut Council,
11
12 608 Oldways Preservation Trust, Paramount Farms and Pulse Canada. **DJAJ** has received research
13
14 609 grants from Saskatchewan Pulse Growers, the Agricultural Bioproducts Innovation Program
15
16 610 through the Pulse Research Network, the Advanced Foods and Material Network, Loblaw
17
18 611 Companies Ltd., Unilever, Barilla, the Almond Board of California, Agriculture and Agri-food
19
20 612 Canada, Pulse Canada, Kellogg's Company, Canada, Quaker Oats, Canada, Procter & Gamble
21
22 613 Technical Centre Ltd., Bayer Consumer Care, Springfield, NJ, Pepsi/Quaker, International Nut
23
24 614 & Dried Fruit (INC), Soy Foods Association of North America, the Coca-Cola Company
25
26 615 (investigator initiated, unrestricted grant), Solae, Haine Celestial, the Sanitarium Company,
27
28 616 Orafit, the International Tree Nut Council Nutrition Research and Education Foundation, the
29
30 617 Peanut Institute, the Canola and Flax Councils of Canada, the Calorie Control Council (CCC),
31
32 618 the CIHR, the Canada Foundation for Innovation and the Ontario Research Fund. He has
33
34 619 received in-kind supplies for trial as a research support from the Almond board of California,
35
36 620 Walnut Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo,
37
38 621 Loblaw Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada,
39
40 622 WhiteWave Foods. He has been on the speaker's panel, served on the scientific advisory board
41
42 623 and/or received travel support and/or honoraria from the Almond Board of California, Canadian
43
44 624 Agriculture Policy Institute, Loblaw Companies Ltd, the Griffin Hospital (for the development of
45
46 625 the NuVal scoring system, the Coca-Cola Company, EPICURE, Danone, Diet Quality Photo
47
48 626 Navigation (DQPN), FareWell, Verywell, True Health Initiative, Saskatchewan Pulse Growers,
49
50
51
52
53
54
55
56
57
58
59
60

Sanitarium Company, Orafiti, the Almond Board of California, the American Peanut Council, the International Tree Nut Council Nutrition Research and Education Foundation, the Peanut Institute, Herbalife International, Pacific Health Laboratories, Nutritional Fundamental for Health, Barilla, Metagenics, Bayer Consumer Care, Unilever Canada and Netherlands, Solae, Kellogg, Quaker Oats, Procter & Gamble, the Coca-Cola Company, the Griffin Hospital, Abbott Laboratories, the Canola Council of Canada, Dean Foods, the California Strawberry Commission, Haine Celestial, PepsiCo, the Alpro Foundation, Pioneer Hi-Bred International, DuPont Nutrition and Health, Spherix Consulting and WhiteWave Foods, the Advanced Foods and Material Network, the Canola and Flax Councils of Canada, the Nutritional Fundamentals for Health, Agri-Culture and Agri-Food Canada, the Canadian Agri-Food Policy Institute, Pulse Canada, the Saskatchewan Pulse Growers, the Soy Foods Association of North America, the Nutrition Foundation of Italy (NFI), Nutra-Source Diagnostics, the McDougall Program, the Toronto Knowledge Translation Group (St. Michael's Hospital), the Canadian College of Naturopathic Medicine, The Hospital for Sick Children, the Canadian Nutrition Society (CNS), the American Society of Nutrition (ASN), Arizona State University, Paolo Sorbini Foundation and the Institute of Nutrition, Metabolism and Diabetes. He received an honorarium from the United States Department of Agriculture to present the 2013 W.O. Atwater Memorial Lecture. He received the 2013 Award for Excellence in Research from the International Nut and Dried Fruit Council. He received funding and travel support from the Canadian Society of Endocrinology and Metabolism to produce mini cases for the Canadian Diabetes Association (CDA). He is a member of the International Carbohydrate Quality Consortium (ICQC). His wife, ALJ, is a director and partner of Glycemic Index Laboratories, Inc., and his sister received funding through a grant from the St. Michael's Hospital Foundation to develop a cookbook for

1
2
3 650 one of his studies. **JL Sievenpiper** has received research support from the Canadian Institutes of
4
5 651 health Research (CIHR), Canadian Diabetes Association (CDA), PSI Foundation, Banting and
6
7 652 Best Diabetes Centre (BBDC), Canadian Nutrition Society (CNS), American Society for
8
9
10 653 Nutrition (ASN), Calorie Control Council, INC International Nut and Dried Fruit Council
11
12 654 Foundation, National Dried Fruit Trade Association, The Coca-Cola Company (investigator
13
14 655 initiated, unrestricted), Dr. Pepper Snapple Group (investigator initiated, unrestricted), The Tate
15
16 656 and Lyle Nutritional Research Fund at the University of Toronto, and The Alberta Pulse Growers
17
18 657 Glycemic Control and Cardiovascular Disease in Type 2 Diabetes Fund at the University of
19
20 658 Toronto. He has received in-kind research support from the Almond board of California, Walnut
21
22 659 Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo, Loblaw
23
24 660 Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada, WhiteWave
25
26 661 Foods. He has received reimbursement of travel expenses, speaker fees, and/or honoraria from
27
28 662 the Canadian Diabetes Association (CDA), Canadian Nutrition Society (CNS), University of
29
30 663 Alabama at Birmingham, Oldways Preservation Trust, Nutrition Foundation of Italy (NFI),
31
32 664 Diabetes and Nutrition Study Group (DNSG) of the European Association for the Study of
33
34 665 Diabetes (EASD), International Life Sciences Institute (ILSI) North America, Dr. Pepper
35
36 666 Snapple Group, Corn Refiners Association, World Sugar Research Organization, Dairy Farmers
37
38 667 of Canada, Società Italiana di Nutrizione Umana (SINU), III World Congress of Public Health
39
40 668 Nutrition, C3 Collaborating for Health, Sprim Brasil, WhiteWave Foods, Rippe Lifestyle,
41
42 669 mdBriefcase, Federation of European Nutrition Societies (FENS), New York Academy of
43
44 670 Sciences, International Diabetes Federation, American Heart Association (AHA), American
45
46 671 Society for Nutrition (ASN), FoodMinds LLC, Memac Ogilvy & Mather LLC, Pulse Canada,
47
48 672 Pepsico, BCFN Foundation, The Ginger Network LLC, and Dietitians of Canada. He has ad hoc
49
50
51
52
53
54
55
56
57
58
59
60

consulting arrangements with Winston & Strawn LLP, Perkins Coie LLP, and Tate & Lyle. He is a member of the European Fruit Juice Association Scientific Expert Panel. He is on Clinical Practice Guidelines Expert Committees of the Canadian Diabetes Association (CDA) European Association for the study of Diabetes (EASD), and Canadian Cardiovascular Society (CCS), as well as an expert writing panel of the American Society for Nutrition (ASN). He serves as an unpaid scientific advisor for the Food, Nutrition, and Safety Program (FNSP) and the Technical Committee on Carbohydrates of the International Life Science Institute (ILSI) North America. He is a member of the International Carbohydrate Quality Consortium (ICQC), Executive Board Member of the Diabetes and Nutrition Study Group (DNSG) of the EASD, and Director of the Toronto 3D Knowledge Synthesis and Clinical Trials foundation. His wife is an employee of Unilever Canada. No competing interests were declared by **CR Braunstein, S Blanco Mejia, and LA Leiter.**

References

1. Malhotra A. Eating MORE fat while cutting carbs and quitting sugar can help you lose weight and be happier, says top cardiologist. Mail Online 2016.

2. Taubes G. Diet Advice That Ignores Hunger. The New York Times. Sunday Review ed 2015.

3. Ludwig D. Always hungry? Conquer cravings, retrain your fat cells and lose weight permanently. New York: Grand Central Life & Style; 2016.

4. Taubes G. Why We Get Fat And What To Do About It. . New York: Alfred A Knopf; 2011.

5. Atkins R. Dr Atkins' Diet Revolution: The High Calorie Way to Stay Thin Forever. : Bantam Books; 1973.

6. Perlmutter D. Grain Brain: The surprising truth about wheat, carbs, and sugar -- Your brain's silent killers: Little, Brown And Company; 2013.

7. David W. Wheat Belly: HarperCollins; 2012.

8. Cordain L. The Paleo Diet Revised: Lose weight and get healthy by eating the foods you were designed to eat: Houghton Mifflin Harcourt; 2010.

9. Teicholz N. The Big Fat Surprise: Why Butter, Meat and Cheese Belong in a Healthy Diet: Simon and Schuster; 2014.

10. National Obesity Forum, The Public Health Collaboration. Eat Fat, Cut The Carbs and Avoid Snacking To Reverse Obesity and Type 2 Diabetes.

11. Malhotra A. Saturated fat is not the major issue. BMJ 2013;347:f6340.

12. Malhotra A. Author's reply to Lichtenstein, Tedstone and Pyne, Mann and colleagues, Lim, and Clifton. BMJ 2013;347:f6855.

13. Thomas DE, Elliott EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. Cochrane Database Syst Rev 2007;CD005105.

14. Schwingshackl L, Hoffmann G. Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. Nutr Metab Cardiovasc Dis 2013;23:699-706.

15. Pol K, Christensen R, Bartels EM, Raben A, Tetens I, Kristensen M. Whole grain and body weight changes in apparently healthy adults: a systematic review and meta-analysis of randomized controlled studies. Am J Clin Nutr 2013;98:872-84.

16. Harland JI, Garton LE. Whole-grain intake as a marker of healthy body weight and adiposity. Public Health Nutr 2008;11:554-63.

17. Slavin JL. Dietary fiber and body weight. Nutrition 2005;21:411-8.

18. Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: Health Implications of Dietary Fiber. J Acad Nutr Diet 2015;115:1861-70.

19. Higgins JP, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions [Internet]. Version 5.1.0. The Cochrane Collaboration. Version current March 2011 [cited 2015 Sep 23]. Available from: www.cochrane-handbook.org.

20. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Ann Intern Med 2009;151:264-9, W64.

21. Elbourne DR, Altman DG, Higgins JP, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. Int J Epidemiol 2002;31:140-9.

22. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ 1997;315:629-34.

23. Begg CB. A measure to aid in the interpretation of published clinical trials. Stat Med 1985;4:1-9.

24. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 2000;56:455-63.

25. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383-94.
26. Jebb SA, Lovegrove JA, Griffin BA, et al. Effect of changing the amount and type of fat and carbohydrate on insulin sensitivity and cardiovascular risk: the RISCK (Reading, Imperial, Surrey, Cambridge, and Kings) trial. *Am J Clin Nutr* 2010;92:748-58.
27. Philippou E, Bovill-Taylor C, Rajkumar C, et al. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 2009;58:1703-8.
28. Philippou E, McGowan BM, Brynes AE, Dornhorst A, Leeds AR, Frost GS. The effect of a 12-week low glycaemic index diet on heart disease risk factors and 24 h glycaemic response in healthy middle-aged volunteers at risk of heart disease: a pilot study. *Eur J Clin Nutr* 2008;62:145-9.
29. Sichieri R, Moura AS, Genelhu V, Hu F, Willett WC. An 18-mo randomized trial of a low-glycemic-index diet and weight change in Brazilian women. *Am J Clin Nutr* 2007;86:707-13.
30. Wolever TM, Mehling C. High-carbohydrate-low-glycaemic index dietary advice improves glucose disposition index in subjects with impaired glucose tolerance. *Br J Nutr* 2002;87:477-87.
31. Pereira EV, Costa Jde A, Alfenas Rde C. Effect of glycemic index on obesity control. *Arch Endocrinol Metab* 2015;59:245-51.
32. Karl JP, Roberts SB, Schaefer EJ, et al. Effects of carbohydrate quantity and glycemic index on resting metabolic rate and body composition during weight loss. *Obesity (Silver Spring)* 2015;23:2190-8.
33. Buscemi S, Cosentino L, Rosafio G, et al. Effects of hypocaloric diets with different glycemic indexes on endothelial function and glycemic variability in overweight and in obese adult patients at increased cardiovascular risk. *Clin Nutr* 2013;32:346-52.
34. Larsen TM, Dalskov SM, van Baak M, et al. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med* 2010;363:2102-13.
35. Solomon TP, Haus JM, Kelly KR, et al. A low-glycemic index diet combined with exercise reduces insulin resistance, postprandial hyperinsulinemia, and glucose-dependent insulinotropic polypeptide responses in obese, prediabetic humans. *Am J Clin Nutr* 2010;92:1359-68.
36. Philippou E, Neary NM, Chaudhri O, et al. The effect of dietary glycemic index on weight maintenance in overweight subjects: a pilot study. *Obesity (Silver Spring)* 2009;17:396-401.
37. Abete I, Parra D, Martinez JA. Energy-restricted diets based on a distinct food selection affecting the glycemic index induce different weight loss and oxidative response. *Clin Nutr* 2008;27:545-51.
38. Aston LM, Stokes CS, Jebb SA. No effect of a diet with a reduced glycaemic index on satiety, energy intake and body weight in overweight and obese women. *Int J Obes (Lond)* 2008;32:160-5.
39. Jensen L, Sloth B, Krog-Mikkelsen I, et al. A low-glycemic-index diet reduces plasma plasminogen activator inhibitor-1 activity, but not tissue inhibitor of proteinases-1 or plasminogen activator inhibitor-1 protein, in overweight women. *Am J Clin Nutr* 2008;87:97-105.
40. Bellisle F, Dalix AM, De Assis MA, et al. Motivational effects of 12-week moderately restrictive diets with or without special attention to the Glycaemic Index of foods. *Br J Nutr* 2007;97:790-8.
41. de Rougemont A, Normand S, Nazare JA, et al. Beneficial effects of a 5-week low-glycaemic index regimen on weight control and cardiovascular risk factors in overweight non-diabetic subjects. *Br J Nutr* 2007;98:1288-98.
42. McMillan-Price J, Petocz P, Atkinson F, et al. Comparison of 4 diets of varying glycemic load on weight loss and cardiovascular risk reduction in overweight and obese young adults: a randomized controlled trial. *Arch Intern Med* 2006;166:1466-75.
43. Visek J, Lacigova S, Cechurova D, Rusavy Z. Comparison of a low-glycemic index vs standard diabetic diet. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:112-6.

1
2
3 790 44. Jenkins DJ, Kendall CW, Vuksan V, et al. Effect of lowering the glycemic load with canola oil on
4 791 glycemic control and cardiovascular risk factors: a randomized controlled trial. *Diabetes Care*
5 792 2014;37:1806-14.
6 793 45. Jenkins DJ, Kendall CW, Augustin LS, et al. Effect of legumes as part of a low glycemic index diet
7 794 on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled
8 795 trial. *Arch Intern Med* 2012;172:1653-60.
9 796 46. Yusof BN, Talib RA, Kamaruddin NA, Karim NA, Chinna K, Gilbertson H. A low-GI diet is
10 797 associated with a short-term improvement of glycaemic control in Asian patients with type 2 diabetes.
11 798 *Diabetes Obes Metab* 2009;11:387-96.
12 799 47. Wolever TM, Gibbs AL, Mehling C, et al. The Canadian Trial of Carbohydrates in Diabetes (CCD),
13 800 a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on
14 801 glycated hemoglobin but reduction in C-reactive protein. *Am J Clin Nutr* 2008;87:114-25.
15 802 48. Jenkins DJ, Kendall CW, McKeown-Eyssen G, et al. Effect of a low-glycemic index or a high-cereal
16 803 fiber diet on type 2 diabetes: a randomized trial. *JAMA* 2008;300:2742-53.
17 804 49. Jimenez-Cruz A, Bacardi-Gascon M, Turnbull WH, Rosales-Garay P, Severino-Lugo I. A flexible,
18 805 low-glycemic index mexican-style diet in overweight and obese subjects with type 2 diabetes improves
19 806 metabolic parameters during a 6-week treatment period. *Diabetes Care* 2003;26:1967-70.
20 807 50. Heilbronn LK, Noakes M, Clifton PM. The effect of high- and low-glycemic index energy
21 808 restricted diets on plasma lipid and glucose profiles in type 2 diabetic subjects with varying glycemic
22 809 control. *J Am Coll Nutr* 2002;21:120-7.
23 810 51. Fontvieille AM, Acosta M, Rizkalla SW, et al. A moderate switch from high to low glycemic-index
24 811 foods for 3 weeks improves metabolic control of type I (IDDM) diabetic subjects. . *Diabetes Nutr Metab*
25 812 1988;1:139-43.
26 813 52. Frost GS, Brynes AE, Bovill-Taylor C, Dornhorst A. A prospective randomised trial to determine
27 814 the efficacy of a low glycaemic index diet given in addition to healthy eating and weight loss advice in
28 815 patients with coronary heart disease. *Eur J Clin Nutr* 2004;58:121-7.
29 816 53. de Assis Costa J, de Cassia Goncalves Alfensa R. The consumption of low glycemic meals reduces
30 817 abdominal obesity in subjects with excess body weight. *Nutricion Hospitalaria* 2012;27:1178-83.
31 818 54. Fontvieille AM, Rizkalla SW, Penfornis A, Acosta M, Bornet FR, Slama G. The use of low
32 819 glycaemic index foods improves metabolic control of diabetic patients over five weeks. *Diabet Med*
33 820 1992;9:444-50.
34 821 55. Bibiloni MDM, Fernandez-Blanco J, Pujol-Plana N, et al. [Improving diet quality in children
35 822 through a new nutritional education programme: INFADIMED]. *Gac Sanit* 2017.
36 823 56. Pounis G, Castelnuovo AD, Costanzo S, et al. Association of pasta consumption with body mass
37 824 index and waist-to-hip ratio: results from Moli-sani and INHES studies. *Nutr Diabetes* 2016;6:e218.
38 825 57. Estruch R, Ros E, Salas-Salvado J, et al. Primary prevention of cardiovascular disease with a
39 826 Mediterranean diet. *N Engl J Med* 2013;368:1279-90.
40 827 58. Shay CM, Van Horn L, Stamler J, et al. Food and nutrient intakes and their associations with
41 828 lower BMI in middle-aged US adults: the International Study of Macro-/Micronutrients and Blood
42 829 Pressure (INTERMAP). *Am J Clin Nutr* 2012;96:483-91.
43 830 59. Smith JD, Hou T, Ludwig DS, et al. Changes in intake of protein foods, carbohydrate amount and
44 831 quality, and long-term weight change: results from 3 prospective cohorts. *Am J Clin Nutr* 2015;101:1216-
45 832 24.
46 833 60. Granfeldt Y, Bjorck I. Glycemic response to starch in pasta: a study of mechanisms of limited
47 834 enzyme availability. . *J Cereal Sci* 1991;14:47-61.
48 835 61. Granfeldt Y, Bjorck I, Hagander B. On the importance of processing conditions, product thickness
49 836 and egg addition for the glycaemic and hormonal responses to pasta: a comparison with bread made
50 837 from 'pasta ingredients'. *Eur J Clin Nutr* 1991;45:489-99.

62. Frost GS, Brynes AE, Dhillon WS, Bloom SR, McBurney MI. The effects of fiber enrichment of pasta and fat content on gastric emptying, GLP-1, glucose, and insulin responses to a meal. *Eur J Clin Nutr* 2003;57:293-8.
63. Korczak R, Timm D, Ahnen R, Thomas W, Slavin JL. High Protein Pasta is Not More Satiating than High Fiber Pasta at a Lunch Meal, Nor Does it Decrease Mid-Afternoon Snacking in Healthy Men and Women. *J Food Sci* 2016;81:S2240-5.
64. Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care* 2008;31:2281-3.
65. Ludwig DS. Dietary glycemic index and obesity. *J Nutr* 2000;130:280S-3S.
66. Riccardi G, Rivellese AA, Giacco R. Role of glycemic index and glycemic load in the healthy state, in prediabetes, and in diabetes. *Am J Clin Nutr* 2008;87:269S-74S.
67. Saris WH, Astrup A, Prentice AM, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the CARMEN study. The Carbohydrate Ratio Management in European National diets. *Int J Obes Relat Metab Disord* 2000;24:1310-8.
68. Ludwig DS, Majzoub JA, Al-Zahrani A, Dallal GE, Blanco I, Roberts SB. High glycemic index foods, overeating, and obesity. *Pediatrics* 1999;103:E26.
69. Wang J, Thornton JC, Bari S, et al. Comparisons of waist circumferences measured at 4 sites. *Am J Clin Nutr* 2003;77:379-84.
70. Makarem N, Scott M, Quatromoni P, Jacques P, Parekh N. Trends in dietary carbohydrate consumption from 1991 to 2008 in the Framingham Heart Study Offspring Cohort. *Br J Nutr* 2014;111:2010-23.
71. Rehm CD, Penalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults, 1999-2012. *JAMA* 2016;315:2542-53.
72. Golzarand M, Mirmiran P, Jessri M, Toolabi K, Mojarad M, Azizi F. Dietary trends in the Middle East and North Africa: an ecological study (1961 to 2007). *Public Health Nutr* 2012;15:1835-44.
73. Leclercq C, Arcella D, Piccinelli R, et al. The Italian National Food Consumption Survey INRAN-SCAI 2005-06: main results in terms of food consumption. *Public Health Nutr* 2009;12:2504-32.
74. Bonaccio M, Di Castelnuovo A, Bonanni A, et al. Decline of the Mediterranean diet at a time of economic crisis. Results from the Moli-sani study. *Nutr Metab Cardiovasc Dis* 2014;24:853-60.
75. Sofi F, Innocenti G, Dini C, et al. Low adherence of a clinically healthy Italian population to nutritional recommendations for primary prevention of chronic diseases. *Nutr Metab Cardiovasc Dis* 2006;16:436-44.
76. Ferdman RA. This is what happens when everyone is terrified of carbs. *The Washington Post* 2015 September 2, 2015.
77. Johnston BC, Kanters S, Bandayrel K, et al. Comparison of weight loss among named diet programs in overweight and obese adults: a meta-analysis. *JAMA* 2014;312:923-33.
78. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;129:S1-45.
79. Anderson TJ, Gregoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. *The Canadian journal of cardiology* 2013;29:151-67.

Table 1: Summary of trial characteristics

Trial Characteristics*	ALL	Neutral Energy Balance	Negative Energy Balance
Trial Number (n)	32	23	9
Trial Size (total, range)	2448 (8 - 250)	1989 (8 - 250)	459 (13 - 123)
Male: Female ^a (%)	40:60	47:53	27:73
Age (years)	50 (40 - 58)	52.0 (42.1 – 59.5)	49.5 (34.2 – 53.0)
Metabolic Phenotype (OW/OB:DM:CHD) (%)	66:31:3	57:39:4	89:11:0
Setting (IP:OP) (%)	3:97	4:96	0:100
Baseline Body Weight (kg) ^b	85.5 (80.0 - 91.9)	84.1 (79.5 – 87.5)	92.5 (86.1 – 93.9)
Baseline BMI (kg/m ²) ^c	30.4 (28.2 - 32.0)	29.5 (27.4 – 31.4)	31.7 (30.1 – 32.9)
Study Design (C:P) (%)	19:81	26:74	0:100
Dose Pasta (servings/week) ^d	3.3 (2.3 - 3.5)	3.4 (2.9 – 4.1)	2.3 (2.3-3.5)
GI in Pasta/LGI group	49.0 (44.0 - 55.1)	46.5 (49.9 – 55.5)	44.0 (42.3 - 49.4)
GI in Higher GI group	62.5 (61.6 - 63.2)	63.3 (60.1 – 64.4)	61.0 (59.2 – 66.6)
Calorie reduction in Pasta/LGI group (kcal) ^e	-179 (-90 - -448)	-165 (-74 - -313)	-447 (-134 - -594)
Calorie reduction in Higher GI group (kcal) ^e	-181 (-93 - -401)	-160 (-40 - -248)	-470 (-172 - -561)
Feeding Control (Met:Supp:DA) (%)	6:44:50	4:48:48	11:33:56
Follow-Up Duration (Weeks)	12 (9-21)	12 (6-24)	12 (10 - 21)
Funding Sources (A:I:AI:NR) (%)	47:9:25:19	44:13:26:17	56:0:22:22

* median (inter quartile range), unless otherwise indicated

^a24/32 trials provided data on sex

^b 30/32 trials reported baseline body weight

^c 28/32 trials reported baseline BMI

^d 11/32 trials provided data from which dose could be approximated

^e 20/32 trials provided data from which to approximate changes in caloric intake

A, agency; AI, agency and industry; BMI, body mass index; C, crossover design; CHD, coronary heart disease; DA, dietary advice; DM, diabetes; GI, glycemic index; I, industry; IP, inpatient; LGI, low glycemic index; Met, metabolic; NR, not reported; OB, obese; OP, outpatient; OW, overweight; P, parallel design; Suppl, supplemented/provision of certain food




Figure 1: Literature Search
40x54mm (300 x 300 DPI)




Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). $n = 2448$. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

40x54mm (300 x 300 DPI)

For peer review only




Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I² statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

40x54mm (300 x 300 DPI)

Online Supplemental Information

Table of Contents

Supplemental Tables..... 4

Supplemental Table S1: PRISMA Checklist 4

Supplemental Table S2: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials 7

Supplemental Table S3: PICO framework of the search strategy..... 9

Supplement Table S4a: Trial characteristics 10

Supplement Table S4b: Trial characteristics continued..... 11

Supplement Table S4c: Trial characteristics continued 12

Supplement Table S4d: Trial characteristics continued..... 13

Supplement Table S4e: Trial characteristics continued..... 14

Supplemental Table S5: Sensitivity analysis of the systematic removal of each trial* 15

Supplemental Table S6: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 for crossover trials 18

Supplement Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹..... 19

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹..... 21

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹..... 23

Supplemental Table S10. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹..... 25

Supplementary Table S11. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg). 27

Supplementary Table S12: GRADE assessment of study quality 28

Supplemental Figures 31

Supplemental Figure S1: Cochrane risk of bias summary for all included trials..... 32

Supplemental Figure S2: Risk of bias proportion graph for all included trials..... 32

Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038) 34

Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285) 35

Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	36
Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)	37
Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)	38
Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448) ..	39
Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)	40
Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	41
Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)	43
Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	45
Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	47
Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) continued (n = 1038)	49
Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	51
Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	53
Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	55
Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	57
Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)	59
Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380) ..	61
Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)	63

Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19) 64

Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure) 65

Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg) 66

Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²) 67

Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%) 68

Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) 69

Supplemental Tables

Supplemental Table S1: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	6
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6-7, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6-7, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7

Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7, Supplemental Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ²) for each meta-analysis.	8-10

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8-10
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11-12, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	12, Supplemental Figures S1-S2
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	12-13, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	12-13, Figures 2-3, Supplemental Figures S3-S7
Risk of bias	22	Present results of any assessment of risk of bias across studies	12, 16

across studies		(see Item 15).	Supplemental Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	13-16, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	17-20
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	20-22
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	22-24
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	27-28

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Supplemental Table S2: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials

Database	Search Period	Search Terms
Medline	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw. 17. overweight.tw. 18. weight*.tw. 19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
Embase	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw. 17. overweight.tw. 18. weight*.tw.

		19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
The Cochrane Library	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycemic index/ 8. glycaemic ind*.tw. 9. glycemic ind*.tw. 10. glycemic load*.tw. 11. glycaemic load*.tw. 12. exp body weight/ 13. body weight*.tw. 14. BMI.tw. 15. body mass index/ 16. body mass index.tw. 17. weight*.tw. 18. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 19. 12 or 13 or 14 or 15 or 16 or 17 20. 18 and 19 21. limit 20 to animals 22. 20 not 21

Supplemental Table S3: PICO framework of the search strategy

PICO framework ^a defined in the present systematic review and meta-analysis			
Participants	Interventions	Comparators	Outcomes
Adult men and women excluding pregnant or breastfeeding women	Low glycemic index interventions where pasta is included as part of the intervention	Higher glycemic index diets where pasta is not included as part of the intervention	Body weight Body mass index (BMI) Body Fat (%) Waist circumference Waist-to-hip ratio

a Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA and PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 2015; 4:1. <https://doi.org/10.1186/2046-4053-4-1>

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
All rights reserved. No reuse allowed without permission. For copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4a: Trial characteristics

Overweight/Obese Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)§§	Dietary Prescription	Funding Source
Karl et al. 2015 †††	39 (19M:20F)**	OB, OP					2.33	USA	P	Negative+ Neutral	17		CR to 2/3kcal; Metabolic	Agency
Low GI	20		56 (5)*	92.9 (13.6)*	32.3 (3.4)*	42:133						68:15:16		
Higher GI	19		56 (5)*	94 (9.7)*	33.4 (2.6)*	61:201						70:16:14		
Pereira et al. 2015	19 (4M:15F)**	OW, IP/OP					NR	Brazil	P	Neutral	6.4		Ad libitum	Unknown
Low GI	10		28(5)	80.0(12.6)	29.9 (2.1)	41.2(2.2) ¥ **						48.3:16.1:32.8		
Higher GI	9		26(3)	79.1(12.2)	29.1 (2.0)	74.1(2.9) ¥ **						54.6:12.7:34.4		
Buscemi et al. 2013	40 (19M:21F)**	OW/OB, high CVD risk, OP					NR	Italy	P	Negative	12		CR to 20kcal/kg/d; Ad libitum	Unknown
Low GI	19		51 (8)	93.8 (17.3)	34.3 (6.6)	48.1: 138						56:18:26		
Higher GI	21		49 (8)	93.2 (14.4)	34.5 (5.1)	59.3: 174						57:16:27		
Costa et al. 2012	17 (7M:10F)	OW, IP/OP	25.4 (5.8)	84.1(16.3)	26.3(3.2)		NR	Brazil	C	Neutral	4		Ab libitum, 2 meals+3 fruit/d provided	NR
Low GI						47.5(3.8)						58.6:13.9:25.5		
Higher GI						61.6(2.8)						55.4:14.2:30.3		
Jebb et al. 2010 - High MUFA	225**	OH, some CV risk factors, OP	~52		~28.5*		1.75	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	115			83.7 (69.6-93.1)¶¶¶		~55.2						44.6:16.4:35.7****		
Higher GI	110			80.5 (70.0-92.1)¶¶¶		~63.3						44.9:15.3:35.6****		
Jebb et al. 2010 - Low Fat	250**	OH, some CV risk factors, OP	~52		~28.5*		3.5	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	117			79.4 (70.1- 91.8)¶¶¶		~56.3						41.5:14.2:26.1****		
Higher GI	111			80.7 (71.4- 91.4)¶¶¶		~64.4						41.1:15.7:27.5****		
Larsen et al. 2010-High Pro	231**	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	124		42.1 (6.5)	88.5 (15.6)		~56.5: 108.9						~43:22:32		
Higher GI	107		42 (5.7)	89.5 (17.1)		~61.4: 113.1						~45:23:31		
Larsen et al. 2010 - Low Pro	203 **	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	106		42.2 (5.7)	88.4 (15.7)		~56.2: 121						~51:18:30		
Higher GI	97		42 (5.9)	86.6 (13.8)		~61.6: 137.9						~51:17:31		

Supplement Table S4b: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Solomon et al. 2010	22 (8M:14F)**	OB, Pre-T2DM, OP					7	USA	P	Neutral	12		Metabolic plus exercise program	Agency
Low GI	10 (3M:7F)		67 (6)	97.4 (12.0)	34.9 (1.1)	39.8 (0.9)						28.3(0.3):17.0(0.3)		
Higher GI	12 (5M:7F)		64 (3)	94.7 (15.2)	34.1 (1.1)	80.0 (2.1)						27.8(0.7):16.6(0.3)		
Philippou et al. 2009- 6 mo	38**	≥1 CHD risk factors, OP	(35-65) ¶				NR	UK	P	Negative (for all but n=2)	24		500kcal CR; Ad libitum	Agency
Low GI	22			91.3(14.8)***	29.1 (3.6)***	50.6 (4.6):114.4(31.5)								
Higher GI	16			97.5(16.4)***	30.5 (3.5)***	63.2(5.6):175.0(45.6)								
Philippou et al. 2009- 4 mo	42**	OW, OP	(18-65)¶				NR	UK	P	Neutral	16		Ad libitum	Unknown
Low GI	23			87.2 (15.3)	32.5 (4.8)	49.7(5.7):89.7(27.5)						19.5(4.2):31.8(5.8)		
Higher GI	19			83.6 (13.4)	31.3 (4.8)	63.7(9.4):136.8(56.3)						19.3(4.9):30.9(9)		
Abete et al. 2008	32 (18M:14F)	OB, OP	36(7)				2.33	Spain	P	Negative	8		30% CR; Ad libitum, 3-day menus	Agency
Low GI	16			94.3(16.1)	32.8 (4.3)	(40-45)¶						18.3(1.6):31.5(1.6)		
Higher GI	16			94.4(13.1)	32.2 (4.4)	(60-65)¶						19.6(5.6):32.6(4.3)		
Aston et al. 2008	19 (0M:19F)**	OW/OB, OP	51.9(7.6)	87.5 (15)	33.1 (4.9)		3.33	UK	C	Neutral	12		Ad libitum, key CHO foods provided	Agency
Low GI						55.5(3.8):133.8(22.9)****						17.0(2.4):32.2(5.1)*		
Higher GI						63.9(3):138.8(30.5)****						17.6(3.3):34.1(5.7)*		
Jensen et al. 2008	44 (0M:44F)**	OW, OP	(20-40)¶				3	Denmark	P	Neutral	10		Ad libitum, partial provision, menu plans	Agency, Industry
Low GI	22 (0M:22F)			77.9(6.9)	27.4 (1.5)	72¥						17(5):17(0):23(5) ‡		
Higher GI	22 (0M:22F)			80.2(1.4)	27.6 (0.3)	95¥						17(5):17(0):22(5) ‡		
Philippou et al. 2008	13 (5M:8F)**	≥1 CHD risk factors, OP					NR	UK	P	Negative	12		500kcal CR; Ad libitum	Agency
Low GI	7 (3M:4F)		54 (49-58)¶¶	81.5 (4.7)***	28.7 (2.1)***	51.3(51.0-52.0):105.6 (76.9-110.1)¶¶						37.8-51.0): 17.1(15.7-37.1): 32.8(31.3-37.1)¶¶		
Higher GI	6 (2M:4F)		45 (39-50)¶¶	89.7 (12.8)***	31.5 (4.4)***	59.3(59.2-64.0):114.7(98.5-134.9)¶¶						47.8-51.7):19.6(14.0-22.4):29.2(25.2-34.5)¶¶		

Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department of GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4c: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Bellisle et al. 2007	65 (OM:65F)**	OW/OB, OP					NR	France	P	Neutral	12		Ad libitum	Industry
Low GI	35		46.1 (13.6)	80 (13.2)	30.2 (4.1)	na								
Higher GI	30		45.3 (12.0)	79 (13.1)	30.4 (4.4)	na								
de Rougemont et al. 2007	38 (22M:16F)**	OW, OP					2.8	France	P	Neutral	5		Ad libitum, some foods provided	Agency, Industry
Low GI	19		36.3 (8.7)	77.2 (9.6)	27.5 (1.3)	46.5 (1.3)						46.5 (1.3):19.8 (1.7):37.7 (4.4)		
Higher GI	19		40.4 (9.6)	77.3 (9.2)	27.2 (1.3)	66.3 (2.6)						66.3 (2.6):17.6 (1.7):38.4 (3.1)		
Sichieri et al. 2007	123 (OM:123F)**	OW, OP					NR	Brazil	P	Negative	72		100-300kcal CR; 6-d menu and exchange lists provided	Agency
Low GI	63		37.2 (5.4)*	67.7 (6.6)*	na	21(38): 74(84)						21(38): 74(84):13.3 (6.3): 13.3: 27.2(4.6)		
Higher GI	60		37.5 (5.6)*	68.5 (7.5)*	na	51(28): 199(43)						51(28): 199(43):12.3 (6.2): 12.3: 26.1(4.7)		
McMillian-Price et al. 2006-High Carb	64(16M:48F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	32		30.5 (7.9)	87.1 (15.3)	30.6 (4.5)	45 (6):89 (28)						45 (6):89 (28):6 (6):19 (0):22 (6)		
Higher GI	32		31.8 (9.6)	86 (10.7)	30.9 (3.4)	70 (6):129 (45)						70 (6):129 (45):20 (6):18 (6):19 (6)		
McMillian-Price et al. 2006-High Protein	65(15M:50F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	33		34.6 (8.6)	88.4 (17.2)	32.1 (5.2)	44 (6):59 (23)						44 (6):59 (23):20 (11):26 (6):28 (6)		
Higher GI	32		30.2 (8.5)	87.7 (16.4)	31.3 (4.5)	59 (6):75 (17)						59 (6):75 (17):22 (6):28 (6):27 (6)		
Wolever et al. 2002	24 (5M:19F)**	IGT, OP					NR	Canada	P	Neutral	16		Ad libitum, partial provision	Agency
Low GI	13(3M:10F)		55.2 (10.8)	79.7 (13.1)***	29.7 (4.3)	54.4 (2.5):91.8 (9.4)						54.4 (2.5):91.8 (9.4):19.4 (1.8):24.7 (5.8)		
Higher GI	11(2M:9F)		58.8 (13.3)	76.4 (20.4)***	29.3 (7.3)	59.3 (2.0):96.8 (11.6)						59.3 (2.0):96.8 (11.6):17.4 (2.3):27.9 (6.3)		

Supplement Table S4d: Trial characteristics continued

Diabetes Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Jenkins et al. 2014	141(77M:64F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum, bread supplement	Industry Association
Low GI	70 (38M:32F)		59 (10)	85 (20)	30 (5)	~51:53						~38.5:19.8:37.2		
Higher GI	71 (39M:32F)		59 (10)	84 (19)	31 (6)	~62:89						~49.2:19.8:27.4		
Visek et al. 2014	20 (12M:8F)	T2DM, OP	62.7 (5.8)	91.9 (14.1)	32 (4.2)		NR	Czech Republic	C	Neutral	12		Ad libitum	Agency
Low GI						49 (48-51)¶¶						~37.2:18.0:36.0		
Higher GI						68 (61-72)¶¶						~36.2:17.3:40.0		
Jenkins et al. 2012	121 (61M: 60F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum	Agency
Low GI	60		58 (10.1)	85.6 (20.1)	31.4 (7.0)	47: 80						45.4:22.8:30.5		
Higher GI	61		61 (7.8)	82.5 (17.2)	29.9 (5.5)	58: 100						48.3:21.4:28.5		
Yusof et al. 2009	100**	T2DM, OP	NR				NR	Malaysia	P	Neutral	12		Ad libitum, key foods provided to lowGI group	Agency
Low GI	51			69.12 (13.33)	27.05 (4.91)	57(6): 108(32)						52(4):18(3):30(4)		
Higher GI	49			66.83 (11.50)	26.79 (4.65)	64(5): 131(30)						54(4):17(3):28(5)		
Jenkins et al 2008	210 (125M:82F)	T2DM, OP					NR	Canada	P	Neutral	24		Ad libitum	Agency
Low GI	106 (65M:41F)		60 (10)	87.0 (20.0)	30.6 (6.0)	49.4: 91.5						44.0:21.2:33.3		
Higher GI	104 (63M:41F)		61 (9)	87.8 (19.4)	31.2 (5.8)	59.3: 117.9						47.5:20.7:30.5		
Wolever et al. 2008	103	T2DM, OW/OB, OP					NR	Canada	P	Neutral	52		Ad libitum, key foods provided	Agency
Low GI	55		60.6 (7.5)*	81.1 (18.7)*	31.6 (4.5)*	55.1 (3.0): 133 (14.8)						46.7(3.7):20.6(3.0):26.5 (5.9)		
Higher GI	48		60.4 (7.9)*	84.4(18.0)*	30.1 (4.3)*	63.2 (2.8): 135 (20.8)						46.9(3.2):20.4 (2.8):30.8 (4.8)		

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4e: Trial characteristics continued

Diabetes Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶¶¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance§	Duration (wks)	Dietary Composition % (SD)§§	Dietary Prescription	Funding Source
Jimenez-Cruz et al. 2003	14 (6M:8F)**	T2DM, OP	59 (34)											
Low GI				91.6 (24.3)	32.4 (6.0)	44(3.4): 86(19.8)		NR	Mexico	C	Neutral	60:21:23	Ad libitum	Industry
Higher GI				92.6 (25.4)	32.3 (6.0)	56(4.9): 139(27.3)						64:18:20		
Heilbronn et al. 2002	45 (23M:22F)**	T2DM, OW, OP			NR			3.5	Australia	P	Negative		CR to 1500kcal/d; Ad libitum, partial provision	Unknown
Low GI	24 (11M:13F)		56.0(9.4)	91.7(16.2)		43						58.9(9.9):22.2 (1.5):17.9 (3.9)		
Higher GI	21 (12M:9F)		57.5(9.6)	93.2 (13.3)		75						60.3(12.3):21.7 (0.9):17.1 (2.3)		
Fontvieille et al. 1992	18 (12M:6F)	T1DM/T2DM, OP	47.2(11.6)	NR	24.8(2.6)			4.7	France	C	Neutral		Ad libitum	Agency, Industry
Low GI						38.1(5.3)						58.2(2.2):18.0(2.5):36.2(6.8)		
Higher GI						64.2(3.1)						67.3(3.3):18.8(1.6):36.3(6.0)		
Fontvieille et al. 1988	8 (4M:4F)	T1DM, OP	43.5 (9.9)	NR	24.1 (6.8)			3.5	France	C	Neutral		Ad libitum, partial provision	Agency, Industry
Low GI						46.5(2.5)						58.1(1.5):17.4 (1.4):35.0 (2.8)		
Higher GI						60.1 (5.1)						63.5(2.5):16.9 (1.7):36.0 (2.8)		
CHD Trial														
Frost et al. 2004	55 (48M:7F)**	CHD, OP						NR	UK	P	Neutral §§§		Ad Libitum	Unknown
Low GI	26 (23M:3F)		63.6 (9.4)	81.2 (12.2)	26.9 (3.3)	50(4):115(39)						59 (5):18 (5):31 (5)		
Higher GI	29 (25M:4F)		61.8 (9)	81.7 (16.7)	28.7 (4.6)	57(4):106(34)						67 (4):10(10):18 (5):32 (10)		

Abbreviations: BMI=Body Mass Index; C=Crossover design; Carb=Carbohydrate; CED= carbohydrate exchange diet; CHD= Coronary Heart Disease; CR= calorie restriction; CV=cardiovascular; GI= glycemic index; GL=glycemic load; IGT= impaired glucose tolerance; IP=inpatient; IR= insulin resistant; High Carb= high carbohydrate; HI= hyperinsulinemic; M = male; mo=months; MUFA=monounsaturated fatty acids; na=not available; T1DM=type 1 diabetes mellitus; T2DM = type 2 diabetes mellitus; OB= obese; OH=otherwise healthy; OP = outpatient; OW= overweight; P=Parallel design; SD= standard deviation; F = female; UK= United Kingdom; USA= United States of America; wks=weeks; y= years

§Negative energy balance diets were designed for weight loss, Neutral energy balance diets for weight maintenance, where all diets are isocaloric between test and control groups; §§Energy from carbohydrate: protein: fat for the planned diet or if not planned; §§§ Participants were advised to lose weight if they had a BMI>28kg/m².

¶ Range of values; ¶¶ median and interquartile range (IQR); ¶¶¶ Actual GI/GL or if not available, planned;

* Calculated before dropout; **Completer Analysis, as used in data analysis; *** completer analysis, data obtained from authors; **** based on less participants than in analysis;

+ approximate based on all study arms; ¥ approximate based on test meals; ‡ approximate based on n=45 from Slovic et al. 2004, the original publication of this study;

‡‡‡ analysis includes weight loss and weight maintenance phases

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplemental Table S5: Sensitivity analysis of the systematic removal of each trial*

	MD [95% CI], P-value I ² , P-value					
	Body Weight (kg) N=32	BMI (kg/m ²) N=18	Body Fat (%) N=10	Waist Circumference (cm) N=18	Waist-to-hip Ratio N=6	Sagittal Abdominal Diameter (cm) N=3
All studies	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.01 [-0.58, 0.56], P=0.98 65%, P=0.003	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92
Removal of:						
OW/OB						
Pereira et al. 2015	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.54	-0.24 [-0.34, -0.14], P<0.01 0.00%, P=0.93	0.19 [-0.25, 0.63], P=0.39 36%, P=0.13	-0.37 [-1.04, 0.31], P=0.28 62%, P<0.01	0.00 [-0.01, 0.01], P=0.67 0.00%, P=0.58	n/a
Karl et al. 2015	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.50	-0.25 [-0.35, -0.16], P<0.01 0.00%, P=0.91	0.10 [-0.51, 0.71], P=0.74 64%, P<0.01	n/a	n/a	n/a
Buscemi et al. 2013	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.86	-0.08 [-0.63, 0.48], P=0.79 64%, P<0.01	-0.46 [-1.06, 0.15], P=0.14 64%, P<0.01	n/a	n/a
Costa et al. 2012	-0.62 [-0.84, -0.41], P<0.01 0.00%, P=0.46	-0.26 [-0.37, -0.16], P<0.01 0.00%, P=0.86	-0.06 [-0.73, 0.61], P=0.86 66%, P<0.01	-0.33 [-0.94, 0.28], P=0.29 59%, P<0.01	n/a	n/a
Jebb et al. 2010 - LowFat	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - LowPro	-0.60 [-0.81, -0.38], P<0.01 0.00%, P=0.57	n/a	n/a	-0.45 [-1.09, 0.18], P=0.16 64%, P<0.01	n/a	-0.05 [-0.39, 0.29], P=0.79 0.00%, P=0.86
Jebb et al. 2010- HighMUFA	-0.64 [-0.85, -0.43], P<0.01 0.00%, P=0.65	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - HighPro	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.48	n/a	n/a	-0.37 [-1.00, 0.26], P=0.26 62%, P<0.01	n/a	-0.13 [-0.45, 0.19], P=0.42 0.00%, P=0.92
Solomon et al. 2010	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	n/a	n/a	n/a	n/a

	0.00%, P=0.47	0.00%, P=0.88				
Philippou et al. 2009 - 6mnh	-0.65 [-0.86, -0.44], P<0.01 0.00%, P=0.59	-0.28 [-0.37, -0.18], P<0.01 0.00%, P=0.97	-0.12 [-0.71, 0.48], P=0.70 65%, P<0.01	-0.55 [-1.15, 0.04], P=0.07 61%, P<0.01	n/a	n/a
Philippou et al. 2009 - 4mnh	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.89	-0.09 [-0.72, 0.54], P=0.78 66%, P<0.01	-0.44 [-1.08, 0.19], P=0.17 64%, P<0.01	n/a	n/a
Abete et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a	n/a
Philippou et al. 2008	-0.62 [-0.83, -0.42], P<0.01 0.00%, P=0.48	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.89	-0.06 [-0.68, 0.55], P=0.84 68%, P<0.01	-0.41 [-1.02, 0.19], P=0.18 63%, P<0.01	n/a	n/a
Aston et al. 2008	-0.66 [-0.87, -0.44], P<0.01 0.00%, P=0.52	n/a	-0.00 [-0.70, 0.69], P=0.99 68%, P<0.01	-0.54 [-1.14, 0.07], P=0.08 62%, P<0.01	n/a	n/a
Jensen et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.27 [-0.36, -0.17], P<0.01 0.00%, P=0.88	n/a	-0.44 [-1.05, 0.18], P=0.16 64%, P<0.01	0.00 [-0.01, 0.00], P=0.24 0.00%, P=0.41	-0.09 [-0.35, 0.18], P=0.51 0.00%, P=0.69
de Rougemont et al. 2007	-0.57 [-0.80, -0.34], P<0.01 0.00%, P=0.53	-0.25 [-0.36, -0.14], P<0.01 0.00%, P=0.87	0.06 [-0.57, 0.70], P=0.84 67%, P<0.01	n/a	n/a	n/a
Sichieri et al. 2007	-0.66 [-0.88, -0.45], P<0.01 0.00%, P=0.53	n/a	n/a	n/a	n/a	n/a
Bellisle et al. 2007	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.88	n/a	-0.47 [-1.08, 0.13], P=0.13 64%, P<0.01	0.00 [-0.01, 0.00], P=0.14 0.00%, P=0.70	n/a
McMillan-Price et al. 2006 - HighCHO	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.51	n/a	n/a	-0.38 [-1.01, 0.25], P=0.23 63%, P<0.01	n/a	n/a
McMillan-Price et al. 2006 - HighPro	-0.70 [-0.91, -0.49], P<0.01 0.00%, P=0.91	n/a	n/a	-0.62 [-1.19, -0.05], P=0.03 55%, P<0.01	n/a	n/a
Wolever et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Diabetes:						
Visek et al. 2014	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	0.01 [-0.57, 0.60], P=0.96	n/a	n/a	n/a

	0.00%, P=0.46	0.00%, P=0.86	68%, P<0.01			
Jenkins et al. 2014	-0.66 [-0.88, -0.43], P<0.01 0.00%, P=0.47	-0.29 [-0.39, -0.18], P<0.01 0.00%, P=0.94	n/a	-0.61 [-1.18, -0.04], P=0.04 50%, P=0.01	0.00 [-0.01, 0.00], P=0.21 0.00%, P=0.43	n/a
Jenkins et al. 2012	-0.62 [-0.84, -0.40], P<0.01 0.00%, P=0.46	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.87	n/a	-0.44 [-1.05, 0.17], P=0.16 64%, P<0.01	0.00 [-0.01, 0.01], P=0.64 0.00%, P=0.53	n/a
Yusof et al. 2009	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.87	n/a	-0.33 [-0.95, 0.28], P=0.29 58%, P<0.01	n/a	n/a
Jenkins et al. 2008	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.36, -0.15], P<0.01 0.00%, P=0.87	n/a	n/a	n/a	n/a
Wolever et al. 2008	-0.64 [-0.84, -0.43], P<0.01 0.00%, P=0.57	n/a	n/a	-0.50 [-1.10, 0.09], P=0.10 62%, P<0.01	n/a	n/a
Jimenez-Cruz et al. 2003	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.86	n/a	n/a	n/a	n/a
Heilbronn et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a	n/a
Fontvieille et al. 1992	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Fontvielle et al. 1988	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
CHD						
Frost et al. 2004	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.17], P<0.01 0.00%, P=0.89	n/a	-0.48 [-1.09, 0.12], P=0.12 63%, P<0.01	0.00 [-0.01, 0.00], P=0.25 1%, P=0.40	n/a

*Sensitivity analysis included the removal of each single study from the meta-analyses one at a time and the summary effect was recalculated. An influential outlier was considered a study whose removal changed the magnitude of the pooled effect by >10%.

BMI, body mass index; CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; MD, mean difference; mnt, month; MUFA, monounsaturated fatty acids; n/a, not applicable; OB, obese; OW, overweight; Pro, protein

Supplemental Table S6: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 in crossover trials

	MD (95% CI), P-value I ² , P-value		
	Correlation Coefficient used in the Primary Analysis	Correlation coefficient used in Sensitivity Analyses	
Outcome (no. crossover trials/total)	0.5	0.25	0.75
Body Weight, kg (6*/32)	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.43], P<0.01 0.00%, P=0.51
BMI, kg/m ² (3/18)	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.90	-0.26 [-0.35, -0.17], P<0.01 0.00%, P=0.90
Body Fat, % (3*/10)	-0.01 [-0.58, 0.56], P=0.98 65%, P<0.01	-0.00 [-0.58, 0.58], P=0.99 64%, P<0.01	-0.02 [-0.57, 0.54], P=0.96 66%, P<0.01
Waist Circumference, cm (2*/18)	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	-0.44 [-1.04, 0.15], P=0.14 60%, P<0.01	-0.46 [-1.07, 0.14], P=0.13 65%, P<0.01
Waist-to-hip Ratio (0/6)	-0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	n/a	n/a
Sagittal Abdominal Diameter, cm (0/6)	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92	n/a	n/a

* one of these crossover trials, however, did not require the use of a correlation coefficient as complete data was available

BMI, body mass index; CI, confidence interval; MD, mean difference; no., number

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplement Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	24.1 – 37.1 kg/m ²	32	2448	-0.04 (-0.13, 0.06)	0.00%	0.454
Follow-up	3 – 72 wks	32	2448	0.01 (-0.01, 0.02)	0.00%	0.314
Dose Pasta	1.75 – 7.0 serv/wk	11	740	0.29 (-0.91, 1.50)	0.00%	0.595
GI ³	21 - 72	31	2383	-0.01 (-0.04, 0.02)	0.45%	0.482
Difference in GI ²	4.9 - 40.2	31	2383	-0.00 (-0.04, 0.03)	2.75%	0.811
Fiber ³	8.0 - 39.4g/d	27	1851	0.02 (-0.01, 0.06)	0.00%	0.221
Change in Fiber ⁴	-7.5 - +13.8g/d	17	1571	-0.02 (-0.07, 0.04)	23.13%	0.595
Difference in Fiber ²	-8.5 - +15.3g/d	27	1851	-0.01 (-0.06, 0.04)	0.00%	0.701
Saturated Fat ³	5.1 - 12.5%	14	1396	-0.05 (-0.30, 0.20)	34.32%	0.648
Change in Saturated Fat ⁴	-8.2 - -1.2%	12	1309	-0.01 (-0.29, 0.28)	46.27%	0.965
Difference in Saturated Fat ²	-2.0 - +2.3%	14	1396	-0.34 (-1.08, 0.39)	32.80%	0.329
CHO ³	37.2 - 68.0%	30	2345	-0.01 (-0.05, 0.03)	0.00%	0.541
Change in CHO ⁴	-7.2 - +10.1%	19	2046	-0.07 (-0.12, -0.01)	0.00%	0.016
Difference in CHO ²	-11.1 - +5.4%	30	2345	0.02 (-0.05, 0.10)	0.00%	0.547
Protein ³	13.3 - 26.1%	30	2345	0.04 (-0.04, 0.12)	0.00%	0.332

Change in Protein ⁴	-0.5 - +8.6%	19	2046	0.15 (0.03, 0.27)	0.00%	0.017
Difference in Protein ²	-2.5 - +3.4%	30	2345	-0.18 (-0.38, 0.02)	0.00%	0.069
Fat ³	16.0 - 37.7%	30	2345	0.01 (-0.05, 0.06)	0.00%	0.852
Change in Fat ⁴	-12.2 - +5.4%	19	2046	0.05 (-0.02, 0.12)	17.73%	0.122
Difference in Fat ²	-4.4 - +10.6%	30	2345	0.02 (-0.05, 0.09)	0.00%	0.610

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body weight on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body weight. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 37.1 kg/m ²	18	1038	-0.01 (-0.06, 0.03)	0.00%	0.559
Follow-up	4 – 24 wks	18	1038	0.00 (-0.01, 0.02)	0.00%	0.559
Dose Pasta*	2.33 – 7.0 serv/wk	4	143			
GI ³	39.8 - 72	17	973	0.01 (-0.01, 0.03)	0.00%	0.153
Difference in GI ²	7.0 - 40.2	17	973	-0.01 (-0.03, 0.01)	0.00%	0.275
Fiber ³	8.0 - 39.4g/d	16	935	0.01 (-0.01, 0.02)	0.00%	0.366
Change in Fiber ⁴	-1.8 - +12.4g/d	10	758	0.00 (-0.03, 0.03)	0.00%	0.846
Difference in Fiber ²	-4.9 - +13.0g/d	16	935	-0.00 (-0.02, 0.02)	0.00%	0.821
Saturated Fat* ³	7.6 - 12.5%	7				
Change in Saturated Fat* ⁴	-2.4 - -1.2%	6				
Difference in Saturated Fat* ²	-1.0 - +2.3%	7				
CHO ³	37.2 - 68.0%	16	935	-0.00 (-0.02, 0.01)	0.00%	0.594
Change in CHO ⁴	-5.6 - +3.2%	10	758	-0.02 (-0.06, 0.02)	0.00%	0.325
Difference in CHO ²	-11.1 - +2.0%	16	935	-0.01 (-0.03, 0.02)	0.00%	0.531
Protein ³	13.9 – 22.8%	16	935	0.00 (-0.04, 0.04)	0.00%	0.905

Change in Protein ⁴	-0.2 - +3.0%	10	758	0.01 (-0.11, 0.14)	0.00%	0.786
Difference in Protein ²	-2.5 - +3.4%	16	935	-0.03 (-0.12, 0.06)	0.00%	0.441
Fat ³	16.0 - 37.7%	16	935	0.00 (-0.02, 0.02)	0.00%	0.717
Change in Fat ⁴	-4.8 - +5.4%	10	758	0.02 (-0.02, 0.06)	0.00%	0.240
Difference in Fat ²	-4.4 - +10.6%	16	935	0.01 (-0.01, 0.04)	0.00%	0.299

*For Dose, there were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in BMI on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in BMI. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	10	285	-0.02 (-0.26, 0.23)	67.69%	0.890
Follow-up	4 – 24 wks	10	285	0.06 (-0.06, 0.17)	66.00%	0.303
Dose Pasta*	2.33 - 3.33 serv/wk	3	96			
GI ³	39.8 - 72	10	285	0.12 (-0.00, 0.25)	52.21%	0.053
Difference in GI ²	7.0 - 40.2	10	285	-0.09 (-0.15, -0.03)	19.39%	0.008
Fiber* ³	8.0 - 39.4g/d	8	228			
Change in Fiber* ⁴	-1.8 - +12.4g/d	4	87			
Difference in Fiber* ²	-4.9 - +13.0g/d	8	228			
Saturated Fat* ³	7.6 - 12.5%	3	93			
Change in Saturated Fat* ⁴	-2.4 - -1.2%	2	51			
Difference in Saturated Fat* ²	-1.0 - +2.3%	3	93			
CHO* ³	37.2 - 68.0%	9	247			
Change in CHO* ⁴	-5.6 - +3.2%	4	87			
Difference in CHO* ²	-11.1 - +2.0%	9	247			
Protein* ³	13.9 – 22.8%	9	247			

Change in Protein* ⁴	-0.2 - +3.0%	4	87		
Difference in Protein* ²	-2.5 - +3.4%	9	247		
Fat* ³	16.0 - 37.7%	9	247		
Change in Fat* ⁴	-4.8 - +5.4%	4	87		
Difference in Fat* ²	-4.4 - +10.6%	9	247		

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body fat on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body fat. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplemental Table S10. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	18	1380	0.10 (-0.13, 0.32)	59.02%	0.372
Follow-up	4 – 52 wks	18	1380	0.05 (-0.03, 0.13)	60.50%	0.225
Dose Pasta*	3.0 – 3.33 serv/wk	2	63			
GI ³	41.2 – 72.0	17	1315	-0.01 (-0.12, 0.10)	66.00%	0.841
Difference in GI ²	4.9 – 32.9	17	1315	-0.02 (-0.10, 0.06)	64.51%	0.596
Fiber ³	8.0 - 39.4g/d	15	1258	0.05 (-0.05, 0.14)	59.60%	0.342
Change in Fiber ⁴	-1.8 - +13.8g/d	13	1176	0.01 (-0.17, 0.20)	70.59%	0.861
Difference in Fiber ²	-4.9 - +15.3g/d	15	1258	-0.00 (-0.14, 0.13)	64.86%	0.939
Saturated Fat* ³	6.1 – 10.1%	8	604			
Change in Saturated Fat* ⁴	-7.6 - -1.2%	7	562			
Difference in Saturated Fat* ²	-2.0 - +2.3%	8	604			
CHO ³	38.5 - 58.6%	16	1277	-0.11 (-0.19, -0.04)	27.06%	0.007
Change in CHO ⁴	-7.2 - +10.1%	13	1176	-0.09 (-0.23, 0.04)	57.19%	0.148
Difference in CHO ²	-11.1 - +5.4%	16	1277	-0.01 (-0.18, 0.17)	60.81%	0.947
Protein ³	13.9 – 26.1%	16	1277	0.20 (0.01, 0.38)	43.92%	0.038

Change in Protein ⁴	-0.3 - +8.6%	13	1176	0.19 (-0.09, 0.46)	63.38%	0.165
Difference in Protein ²	-2.5 - +3.4%	16	1277	-0.14 (-0.66, 0.38)	62.79%	0.570
Fat ³	22.0 - 37.2%	16	1277	0.10 (-0.05, 0.26)	53.48%	0.185
Change in Fat ⁴	-12.2 - +5.4%	13	1176	0.08 (-0.07, 0.24)	61.15%	0.274
Difference in Fat ²	-4.4 - +10.6%	16	1277	0.09 (-0.06, 0.25)	47.17%	0.220

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in waist circumference on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in waist circumference. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplementary Table S11. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg)

Dose threshold of servings of pasta consumed per week	Dose ranges of servings of pasta consumed per week	β (95% CIs) *	Residual I^2 †	p -value
3.0	≤ 3.0	-0.70 (-3.27, 1.86)	0.00%	0.890
	> 3.0	0.91 (-0.89, 2.70)		
3.33	≤ 3.33	0.05 (-1.80, 1.89)	0.00%	0.518
	> 3.33	0.44 (-1.75, 2.63)		
3.5	≤ 3.5	0.09 (-1.65, 1.82)	0.00%	0.888
	> 3.5	0.46 (-1.89, 2.81)		

* β is the slope derived from the piecewise linear meta-regression analyses and represents the treatment effect on body weight for doses above and below each dose-threshold representing servings of pasta consumption per week. † The residual I^2 value indicates heterogeneity unexplained by each dose-threshold.

-2017-019438 on 24 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplementary Table S12: GRADE assessment of study quality

Quality assessment*							№ of patients		Effect	Quality Importance	
№ of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)		
Body Weight (follow up: median 12 weeks)										Downloaded from https://open.bmj.com/ on June 5, 2025 at Department of Clinical Nutrition, AI training, and similar technologies.	
32	randomised trials	not serious	not serious	serious ^a	not serious	none	1294	1228	MD -0.63 kg (-0.84 to -0.42)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
BMI (follow up: median 12 weeks)											
18	randomised trials	not serious	not serious	serious ^a	not serious	none	551	538	MD -0.26 kg/m ² (-0.36 to -0.16)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Waist Circumference (follow up: median 12 weeks)											
18	randomised trials	not serious	serious ^b	serious ^a	not serious	none	717	624	MD -0.46 cm (-0.05 to 0.14)		⊕⊕○○ LOW ^{a,b} Due to downgrade for inconsistency and indirectness

Quality assessment*							№ of patients		Effect	Quality Importance	
№ of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)		
Body Fat (follow up: median 12 weeks)										Downloaded from https://bmjopen.bmj.com/ on June 5, 2025 at Department of Health, Behavior, and Society, Johns Hopkins University. All rights reserved. For uses related to text and data mining, AI training, and similar technologies.	
10	randomised trials	not serious	serious ^c	serious ^a	not serious	none	176	165	MD -0.01 % (-0.58 to 0.56)		⊕⊕○○ LOW ^{a,c} Due to downgrade for inconsistency and indirectness
Waist-to-hip Ratio (follow up: median 12 weeks)											
6	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	223	222	MD -0.00 (-0.01 to 0.00)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Sagittal Abdominal Diameter (follow up: median 26 weeks)											
3	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	237	214	MD -0.09 cm (-0.34 to 0.16)	⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness	

CI, Confidence interval; GI, glycemic index; LGI, low glycemic index; MD, Mean difference

*All outcomes started with high quality evidence since all studies were randomized controlled trials. Risk of Bias – We rated down for risk of bias if the majority of studies were considered to be at high risk of bias. Inconsistency – We assessed inconsistency using I^2 estimates where an I^2 of 50% or higher indicates substantial heterogeneity. I^2 is the percentage of variability in the treatment estimates that is attributable to heterogeneity between studies. We rated down for inconsistency if there was substantial heterogeneity that was unexplained by any a priori sensitivity or subgroup analyses. Indirectness – We rated down for indirectness if there were factors present relating to the population, interventions, and outcomes that limited the generalizability of the results. Imprecision – We rated down for imprecision if the 95% confidence interval (95% CI) crossed the minimally important difference (MID) for harm. MIDs used for each outcome are: 0.5kg for body weight (based on Johnston et al. JAMA 2014;312(9):923-33); 0.2m² for BMI; 2.0cm for waist circumference; 2.0% for body fat; 0.02 for waist-to-hip ratio; 2.0cm for sagittal abdominal diameter.

a. Downgrade for serious indirectness, since most of the available trials did not quantify the amount of data consumed in the context of the low-GI dietary patterns

b. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2=62\%$, P -heterogeneity<0.001), which could not be explained

c. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2= 65\%$, P -heterogeneity=0.003), which could not be explained

d. No downgrade for publication bias, as publication bias could not be assessed due to lack of power for assessing funnel plot asymmetry and small study effects (<10 trials included in the meta-analysis)


Supplemental Figures

	Random Sequence Generation	Allocation Concealment	Blinding of Participants, Personnel and Outcome Assessment	Incomplete Outcome Data	Selective Reporting
Abete et al. 2008	?	?	?	+	+
Aston et al. 2008	+	?	+	+	+
Bellisle et al. 2007	?	?	?	?	+
Buscemi et al. 2012	+	+	+	?	+
Cost et al. 2012	?	?	+	+	+
de Rougemont et al. 2007	?	+	+	+	+
Fontvielle et al. 1992	?	?	+	+	+
Fontvielle et al. 1988	?	?	+	+	+
Frost et al. 2004	?	?	+	+	+
Heilbronn et al. 2002	?	?	+	?	+
Jebb et al. 2010 - HighMUFA	+	?	+	?	+
Jebb et al. 2010 - LowFat	+	?	+	?	+
Jenkins et al. 2014	+	+	+	+	+
Jenkins et al. 2012	+	+	+	+	+
Jenkins et al. 2008	+	+	+	+	+
Jensen et al. 2008	?	?	+	+	+

	Random Sequence Generation	Allocation Concealment	Blinding of Participants, Personnel and Outcome Assessment	Incomplete Outcome Data	Selective Reporting
Jimenez-Cruz et al. 2003	?	?	?	?	+
Karl et al. 2015	+	+	+	+	+
Larsen et al. 2010 -LowPro	+	+	+	+	+
Larsen et al. 2010 -HighPro	+	+	+	+	+
McMillan-Price et al. 2006- HighCHO	?	?	+	+	+
McMillan-Price et al. 2006- HighPro	?	?	+	+	+
Pereira et al. 2015	?	?	+	?	+
Philippou et al. 2008	?	?	+	?	+
Philippou et al. 2009-4mo	?	?	+	?	+
Philippou et al. 2009-6mo	?	?	?	?	+
Sicheri et al. 2007	+	?	+	?	+
Solomon et al. 2010	?	?	+	+	+
Vissek et al. 2014	?	?	+	+	+
Wolever et al. 2008	+	+	?	+	+
Wolever et al. 2002	+	+	+	+	+
Yusof et al. 2009	+	?	+	+	+


Supplemental Figure S1: Cochrane risk of bias summary for all included trials

Summary of risk of bias ratings for each individual study included in the meta-analysis.



Supplemental Figure S2: Risk of bias proportion graph for all included trials


For peer review only



Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


BMI, body mass index; CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot’s bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran’s Q statistic at a significance level of $P<0.10$ and quantified with the use of the I^2 statistic where $I^2\geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot’s bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance fixed effects models. Between-study heterogeneity was detected with the use of the Cochran’s Q statistic at a significance level of $P<0.10$ and quantified with the use of the I^2 statistic where $I^2\geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.




Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and it indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p = 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.

BMI, body mass index; btw, between; CHD, coronary heart disease; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; serv, serving; trt, treatment; wk, week.

*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.02 kg (-0.08, 0.65) (1 vs. 2) to 1.32 kg (-4.28, 6.91) (1 vs. 3) to -1.29 kg (-6.91, 4.32) (2 vs. 3).


† data available on 31 studies

‖ data available on 17 studies

‡ data available on 27 studies

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and it indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.


BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); trt, treatment.

§ data available on 30 studies

¥ data available on 19 studies

¶ data available on 14 studies

Ψ data available on 12 studies




Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup.

LRB=Low Risk of Bias, URB= Unclear Risk of Bias, HRB=High Risk of Bias. *Within and between subgroup analysis could not be performed against HRB since no values for HRB subgroup. ** no values for URB subgroup. § Blinding of Participants, Personnel, and Outcome Assessors.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I² was estimated with the use of the Cochran's Q statistic and indicates the between study

heterogeneity unexplained by the variability in response because of the specific between studies factor. 0.05 indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=9 trials), therefore analyses were not performed.

*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.00kg/m² (-0.156, 0.274) (1 vs. 2) to 0.584kg/m² (-1.009, 2.178) (1 vs. 3) to -0.525kg/m² (-2.122, 1.072) (2 vs. 3).


† data available on 17 studies

‡ data available on 16 studies

|| data available on 10 studies

BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47




Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) continued (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the degree of heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=7 trials), saturated fat within treatment (n=6 trials) and saturated fat between treatments (n=7 trials), therefore analyses were not performed.

§ data available on 16 studies

¥ data available on 10 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.



Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)


The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the degree of heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup.

*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.


BMI, body mass index; CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias.



Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=9), test carbohydrate, protein and fat (n=9), carbohydrate, protein and fat within treatment (n=4), carbohydrate, protein and fat between treatment (n=9), test fibre (n=8), fibre within treatment (n=4), fibre between treatment (n=8), test saturated fat (n=3 trials), saturated fat within treatment (n=2 trials) and saturated fat between treatments (n=3 trials), therefore analyses were not performed.

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.



Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)


The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup.

*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.

CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias



Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=2 trials), therefore analyses were not performed.


*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.158cm (-1.492, 1.818) (1 vs. 2) to 1.608cm (-3.432, 6.648) (1 vs. 3) to -1.445cm (-6.630, 3.740) (2 vs. 3).

† data available on 17 studies

‡ data available on 15 studies

BMI, body mass index; btw, between; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 9, 2025 at Department GEZ-LTA Erasmus Hogeschool. by copyright, including for uses related to text and data mining, AI training, and similar technologies.




Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's I^2 statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=8 trials), saturated fat within treatment (n=7 trials) and saturated fat between treatments (n=8 trials), therefore analyses were not performed.

§ data available on 16 studies

¥ data available on 13 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.



Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)


The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's I^2 statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup.

*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.


** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.


CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias




Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)
CHO, carbohydrate; MD, mean difference



Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)
MD, mean difference




Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure)




Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.




Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.

Supplemental Table S1: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	6
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6-7, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6-7, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and	7, Supplemental

		simplifications made.	Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	8-10

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8-10
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11-12, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	12, Supplemental Figures S1-S2
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	12-13, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	12-13, Figures 2-3, Supplemental Figures S3-S7
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see item 15).	12,16 Supplemental

			Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	13-16, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	17-20
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	20-22
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	22-24
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	27-28

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

BMJ Open

Effect of Pasta in the Context of Low Glycemic Index Dietary Patterns on Body Weight and Markers of Adiposity: A Systematic Review and Meta-analysis of Randomized Controlled Trials in Adults

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019438.R2
Article Type:	Research
Date Submitted by the Author:	26-Jan-2018
Complete List of Authors:	Chiavaroli, Laura; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit; University of Toronto, Nutritional Sciences Kendall, Cyril; University of Toronto, Nutritional Sciences Braunstein, Catherine; University of Toronto, Nutritional Sciences; St. Michael's Hospital, Toronto 3D Knowledge Synthesis and Clinical Trials Unit Blanco Mejia, Sonia; Clinical Nutrition and Risk Factor Modification Centre, St. Michael's Hospital; Toronto 3D Knowledge Synthesis and Clinical Trials Unit, St. Michael's Hospital Leiter, Lawrence; St. Michael's Hospital, Endocrinology; University of Toronto, Nutritional Sciences Jenkins, David; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Center; University of Toronto, Nutritional Sciences Sievenpiper, John; McMaster University, Department of Pathology and Molecular Medicine; St. Michael's Hospital, Clinical Nutrition and Risk Factor Modification Centre
Primary Subject Heading:	Nutrition and metabolism
Secondary Subject Heading:	Cardiovascular medicine
Keywords:	body weight, pasta, glycemic index, glycaemic index, systematic review and meta-analysis, weight loss

SCHOLARONE™
Manuscripts

Laura Chiavaroli^{1,2}, Cyril WC Kendall^{1,2,3}, Catherine R Braunstein^{1,2}, Sonia Blanco Mejia^{1,2},
Lawrence A Leiter^{1,2,4-6}, David JA Jenkins^{1,2,4-6}, John L Sievenpiper^{1,2,4-6}

ABSTRACT

Objective: Carbohydrate staples such as pasta have been implicated in the obesity epidemic. It is unclear whether pasta contributes to weight gain or like other low-glycemic index (GI) foods contributes to weight loss. We synthesized the evidence of the effect of pasta on measures of adiposity.

Design: Systematic review and meta-analysis using the GRADE approach.

Data sources: MEDLINE, Embase, CINAHL, and the Cochrane Library were searched through 07 February 2017.

Eligibility criteria for selecting studies: We included randomized controlled trials ≥ 3 -weeks assessing the effect of pasta alone or in the context of low-GI dietary patterns on measures of global (body weight, BMI, body fat) and regional (waist circumference, waist-to-hip ratio, sagittal abdominal diameter) adiposity in adults.

Data extraction and synthesis: Two independent reviewers extracted data and assessed risk of bias. Data were pooled using the generic inverse-variance method and expressed as mean differences (MDs) with 95% confidence intervals (95% CIs). Heterogeneity was assessed (Cochran Q statistic) and quantified (I^2 -statistic). GRADE assessed the certainty of the evidence.

Results: We identified no trials of the effect of pasta alone and 32 trial comparisons (n=2,448 participants) of the effect of pasta in the context of low-GI dietary patterns. Pasta in the context of low-GI dietary patterns significantly reduced body weight (MD=-0.63kg [95% CIs,-0.84,-0.42kg]) and BMI (MD=-0.26kg/m² [95% CIs,-0.36,-0.16kg/m²]) compared with higher-GI dietary patterns. There was no effect on other measures of adiposity. The certainty of the evidence was graded as moderate for body weight, BMI, WHR, and SAD and low for WC and body fat.

Conclusions: Pasta in the context of low-GI dietary patterns does not adversely affect adiposity and even reduces body weight and BMI compared to higher GI dietary patterns. Future trials should assess the effect of pasta in the context of other healthy dietary patterns.

Protocol registration: ClinicalTrials.gov Identifier, NCT02961088

Strengths and limitations of this study

- The present systematic review and meta-analysis was based on a comprehensive search and includes a large number of randomized controlled trials which provide the best protection against bias.
- We used the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach to evaluate the strength and quality of the evidence.
- There was evidence of unexplained inconsistency in the intervention estimates across trials for waist circumference and body fat.
- The generalizability of our results is questionable with evidence of indirectness in the pooled estimates for all body and adiposity outcomes, as all of the trials assessed pasta in the context of low-GI dietary patterns (no trials assessed the effect of pasta alone or in the context of other dietary patterns) and most of the available trials did not quantify the amount of pasta consumed.

INTRODUCTION

As the role of saturated fat in chronic disease has been called into question, carbohydrates have come under attack in the media^{1,2}, popular books³⁻⁹, statements of health advocacy groups¹⁰ and commentaries in leading medical journals^{11,12}. Much of the attention has focused on sugars but traditional carbohydrate staples like pasta, rice, and breads are increasingly being implicated in the epidemics of overweight and obesity^{1,7}. Although systematic reviews and meta-analyses of randomized controlled trials of dietary patterns that are high in these foods but low in glycemic index (GI)^{13,14}, high in whole grains^{15,16}, and/or high in dietary fibre have shown advantages for weight related outcomes^{17,18}, there has been a general lack of recognition of the importance of carbohydrate quality.

Pasta is an important example of a food which is considered a refined carbohydrate but has a low-GI, a property that has been exploited extensively in studies of low-GI dietary patterns. It remains unclear whether pasta alone or in the context of a low-GI dietary pattern shares the advantages of other low-GI foods or on the contrary contributes to weight gain. We are not aware of any systematic reviews and meta-analyses which have synthesized the evidence of the effect of pasta on body weight outcomes. We undertook a systematic review and meta-analysis of randomized controlled trials (RCTs) using the GRADE approach to quantify the effect of pasta alone or in the context of low-GI dietary patterns on body weight and measures of adiposity relevant to the prevention and management of overweight and obesity.

METHODS

Design

Our protocol followed the guideline of the Cochrane Handbook for Systematic Reviews of Interventions¹⁹, and findings are reported according to the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses²⁰ (**Supplemental Table S1**). The protocol is registered at clinicaltrials.gov (identifier, NCT02961088).

Data sources and searches

We searched MEDLINE (<http://www.nlm.nih.gov/bsd/pmresources.html>), Embase (<https://www.embase.com>), CINAHL (<https://health.ebsco.com/products/the-cinahl-database>), and the Cochrane Library (<http://www.cochranelibrary.com/>) from inception through 07 February 2017. The full search terms used in this study are presented in **Supplemental Tables S2-S3**. Briefly, we searched using variations of the terms pasta and glycemic index and glycemic load and body weight and BMI. The search was limited to human studies and had no language restrictions. Reference lists of selected studies and reviews were also searched to identify additional articles.

Study selection

We include RCTs that investigated the effect of pasta consumed alone or in the context of low-GI dietary patterns that emphasized pasta in comparison with higher GI diets that did not include pasta on body weight or other measures of global (BMI, body fat) or abdominal (waist circumference, waist-to-hip ratio, sagittal abdominal diameter, or visceral adipose tissue assessed by imaging modalities) adiposity in participants of all health backgrounds. Trials were included if the intervention arm assessed the effect of pasta consumed alone or assessed the effect of a low GI diet which emphasized pasta as part of the low GI dietary advice. Trials were excluded if they

had diet duration of <3 weeks, did not intend to use a calorie- and macronutrient-matched comparator arm that was higher in GI; included pregnant or breast-feeding women or children; or did not provide suitable end-point data. When multiple publications existed for the same study, the article with the most information was included (n=6). Published abstracts were not included.

Data extraction

Two reviewers (LC, CB) assessed the titles and abstracts of all identified studies and independently reviewed and extracted relevant data from each report, including study design, blinding, sample size, participant characteristics, follow-up duration, identification of pasta in the low-GI diet only, comparator diet, macronutrient profile, funding source, and outcome data. The primary outcome was body weight, and secondary outcomes included markers of global (BMI, body fat) and abdominal (waist circumference, waist-to-hip ratio, sagittal abdominal diameter, or visceral adipose tissue assessed by imaging modalities) adiposity. Mean±SD differences between test and control arms were extracted for each outcome.

In those trials where the data were included in figures and not provided numerically, we used the software program Plot Digitizer version 2.6.8 (<http://plotdigitizer.sourceforge.net/>) to extract the data. This program is a JAVA program that digitizes scanned figures of X and Y plots from GIF, JPEG, or PNG image file formats and allows one to calibrate the X and Y axes for the estimation data points. Additional information was requested from the authors of all included trials.

Disagreement were resolved by consensus or where necessary by a third author (SBM).

Risk of bias assessment

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Risk of bias for each individual study was assessed using the Cochrane Risk of Bias tool¹⁹. The level of bias was evaluated for sequence generation, allocation concealment, blinding, incomplete outcome data, and selective reporting and determined overall as either low (proper methods taken to reduce bias), high (improper methods creating bias), or unclear (insufficient information provided to determine the bias level).

Statistical analysis

Baseline, end-of-study, and between-treatment differences in body weight, BMI, waist circumference, body fat, waist-to-hip ratio, sagittal abdominal diameter or visceral adipose tissue assessed by imaging modalities were recorded as means±SDs. If not provided, between-treatment differences in change-from-baseline or end differences were calculated by subtracting means and variance measures such as SEs were imputed with the use of published formulas¹⁹. Missing SDs were imputed with the use of the pooled SD from other studies included in the analysis¹⁹. Data analyses were conducted using Review Manager (RevMan) version 5.3 (Copenhagen, Denmark: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014) for primary analyses and Stata version 13 (College Station, TX: StataCorp LP) for subgroup analyses. A generic inverse-variance method with random-effects models was used to calculate pooled mean differences and 95% confidence intervals (CIs). Random-effects models were used even in the absence of statistically significant inter-study heterogeneity, as they yield more conservative summary effect estimates in the presence of residual heterogeneity. Change-from-baseline differences were preferred over end differences and paired analyses were applied to all crossover

160 trials with the use of a within-individual correlation coefficient between treatments of 0.5 as
161 described by Elbourne et al.²¹

162 Inter-study heterogeneity was assessed by the Cochran Q statistic, where $P < 0.10$ was considered
163 statistically significant, and quantified by the I^2 statistic, where $I^2 \geq 50\%$ indicates substantial
164 heterogeneity¹⁹. Sensitivity analysis were performed which included the removal of each single
165 study from the meta-analyses one at a time and recalculation of the summary effect. An
166 influential study was considered a study whose removal changed the magnitude of the pooled
167 effect by $>10\%$. Sensitivity analysis were also conducted using different correlation coefficient
168 values for crossover trials (0.25 and 0.75) to test for the robustness of the effect size, conducting
169 analyses using fixed effects models and restricting analyses to those trials for which pasta intake
170 could be quantified.

171 If ≥ 10 trial comparisons were available, then sources of heterogeneity were explored by
172 subgroup analyses. A priori categorical subgroups analyses were assessed by meta-regression
173 analyses. These included patient type (normal body weight, overweight or obese [average
174 baseline BMI $>27\text{kg/m}^2$]), diabetes, coronary heart disease), follow-up (<24 -weeks, ≥ 24 weeks),
175 baseline BMI (BMI ≤ 30 , $>30\text{kg/m}^2$), design (parallel, crossover), energy balance (negative on
176 both arms [weight loss diets], neutral on both arms [weight maintaining diets]) and dose of pasta
177 (based on the median). A priori categorical subgroup analyses also included the following dietary
178 factors: GI (absolute level [≤ 55 , >55 ; glucose scale], within-treatment change, between-treatment
179 change), fat intake (absolute level [$<30\%$, $\geq 30\%$ energy], within-treatment change, between-
180 treatment change), carbohydrate intake (absolute level [$<50\%$, $\geq 50\%$ energy], within-treatment
181 change, between-treatment change), protein intake (absolute level [$<20\%$, $\geq 20\%$ energy], within-
182 treatment change, between-treatment change), dietary fibre intake (absolute level [$<28\text{g/day}$,

183 $\geq 28\text{g/day}$], within-treatment change, between-treatment change), and risk of bias. A priori
184 continuous meta-regression analyses were conducted on the absolute levels and within- and
185 between-treatment changes of these same dietary factors in the intervention arms of pasta in the
186 context of low-GI dietary patterns. Linear and non-linear pasta intake dose-response analyses
187 were assessed by using continuous meta-regression analyses and spline curve modeling
188 (MKSPLINE procedure), respectively. Publication bias was assessed by visual inspection of
189 funnel plots and the Egger²² and Begg²³ tests, when ≥ 10 trial comparisons were available. If
190 publication bias was suspected, then adjustment for funnel plot asymmetry was done by imputing
191 missing study data using the Duval and Tweedie trim and fill method ²⁴.

193 **Grading the evidence**

194 The grading of recommendations assessment, development, and evaluation (GRADE) approach
195 was used to assess the certainty of the evidence ²⁵. Evidence was graded as high, moderate, low
196 or very low quality. The included RCTs were graded as high quality evidence by default and
197 downgraded based on pre-specified criteria. Criteria to downgrade evidence included risk of bias
198 (weight of studies show risk of bias assessed by the Cochrane Risk of Bias tool), inconsistency
199 (substantial unexplained heterogeneity, $I^2 > 50\%$, $P < 0.10$), indirectness (presence of factors that
200 limited the generalizability of the results), imprecision (the 95% CI for effect estimates were
201 wide or crossed pre-specified minimally important differences [MIDs] for harm), and publication
202 bias (significant evidence of small-study effects).

204 **Patient involvement**

No patients were directly involved in the development of the research question, selection of the outcome measures, design and implementation of the study, or interpretation of the results.

RESULTS

Search results

Figure 1 shows the flow of the literature. We identified 4876 reports of which 29 met eligibility criteria. No reports were identified that assessed the effect of pasta alone, while 29 reports (including 32 trial comparisons involving 2448 participants) were identified that assessed the effect of pasta in the context of low-GI dietary patterns on any adiposity outcome in adults²⁶⁻⁵⁴. Of the 32 trial comparisons that assessed the effect of pasta in the context of a low-GI dietary pattern, there were 32 trial comparisons for body weight, 18 trial comparisons for BMI^{27,28,31-33,35,36,39-41,43-46,48,49,52,53}, 18 trial comparisons for waist circumference^{27,28,31,33,34,36,38-40,42,44-47,52,53}, 10 trial comparisons for body fat^{27,28,31-33,36,38,41,43,53}, 6 trial comparisons for waist-to-hip ratio^{31,39,40,44,45,52}, and 3 trial comparisons for sagittal abdominal diameter^{34,39}. There was only one trial comparison identified³⁵ for visceral adipose tissue assessed by imaging modalities, thus a meta-analysis could not be undertaken for this outcome.

Trial characteristics

Table 1 and **Supplemental Table S4** show the characteristics of all included trials of the effect of pasta in the context of low-GI dietary patterns. The majority of trials had a parallel design (26/32) with a median follow-up of 12 weeks (IQR: 9–21 weeks) and a median number of participants per trial of 43 (IQR: 21–112). Most participants were middle aged (median age: 50

y; IQR: 40-58 y) men and women (median ratio of men to women: 0.4:1 in available trials). The median baseline BMI across studies was 30.4kg/m² (IQR: 28.2–32.0). Regarding metabolic phenotype, 21 (66%) trials included participants who were overweight or obese (had a baseline BMI≥27kg/m²), 10 (31%) had diabetes and one trial (3%) with coronary heart disease (CHD). We did not retrieve any trials where participants had a normal BMI at baseline (≤25kg/m²), although 6 trials did not include BMI >25 kg/m² as part of criteria, the average baseline BMI was ≥27 kg/m², therefore categorized as overweight.

Risk of bias

Supplemental Figures S1 and S2 show the individual Cochrane Risk of Bias tool assessments for each of the included trials of the effect of pasta in the context of low-GI dietary patterns. No serious risk of bias was detected.

Effect of Pasta in the Context of Low-GI dietary patterns and Body Weight

Figure 2 shows the effect of pasta in the context of low-GI dietary patterns on the primary outcome body weight. Pooled analyses showed pasta in the context of low-GI dietary patterns had the effect of reducing body weight by -0.63 kg (95% CI:-0.84, -0.42 kg; P<0.001) compared with higher GI control diets with no evidence of heterogeneity (I²=0%, P-heterogeneity=0.51).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Global Adiposity

Figure 3 and Supplemental Figure S3-S4 show the effect of pasta in the context of low-GI dietary patterns on markers of global adiposity. Pooled analyses showed that pasta in the context of low-GI dietary patterns had the effect of reducing BMI by -0.26kg/m² (n=18 trials; MD=-

0.26kg/m²; 95% CI:-0.36, -0.16 kg/m²; P<0.001) compared with higher GI control diets with no evidence of heterogeneity (I²=0%, P-heterogeneity=0.90). There was no effect on body fat (n=10 trials; MD=-0.01%, 95% CI:-0.58, 0.56%; P=0.98) with evidence of substantial heterogeneity (I²=65%, P-heterogeneity<0.01).

Effect of Pasta in the Context of Low-GI dietary patterns on Markers of Abdominal Adiposity

Figure 3 and Supplemental Figures S5-S7 show the pooled estimates for the markers of abdominal adiposity. Pooled analyses did not show a significant effect of pasta in the context of low-GI dietary patterns compared with higher-GI control diets on waist circumference (n=18 trials; MD=-0.46cm, 95% CI:-1.05, 0.14cm; P=0.13), waist-to-hip ratio (n=6 trials; MD=-0.00, 95% CI:-0.01, 0.00; P=0.27), or sagittal abdominal diameter (n=3 trials, MD=-0.09cm, 95% CI:-0.34, 0.16cm; P=0.48) There was only evidence of substantial heterogeneity in the analyses for waist circumference (I²=62%, P-heterogeneity<0.01).

Sensitivity analyses

We conducted four sets of sensitivity analyses (**Supplemental Tables S5-6, Supplemental Figures S8-9**). The systematic removal of each trial did not modify the direction or significance of the effect estimates or the evidence of heterogeneity for any of the outcomes with the exception of waist circumference (**Supplemental Table S5**). In the sensitivity analysis for waist circumference, two studies were influential studies in that their removal altered the magnitude of the pooled effect in the remaining studies by >10%, where the removal of the studies of McMillan-Price et al. (high protein comparison)⁴² and Jenkins et al.⁴⁴ rendered the results for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

waist circumference statistically significant (MD= -0.62cm; 95% CI: -1.19, -0.05; P=0.03) and (MD= -0.61cm; 95% CI: -1.18, -0.04; P=0.04), respectively; forest plots not shown). Heterogeneity remained significant in both cases ($I^2= 55\%$, P-heterogeneity<0.01 and $I^2=50\%$, P-heterogeneity=0.01, respectively). Sensitivity analyses using correlation coefficients of 0.25 and 0.75 for paired analyses of crossover trials also did not modify the results (**Supplemental Table S6**). In the sensitivity analyses where fixed effects models were applied (**Supplemental Figure S8**), the direction, magnitude and significance of the pooled estimates were very similar to those produced by the random effects models with the exception of the sensitivity analysis for waist circumference which was significant (MD= -0.62; 95% CI: -0.93, -0.32; P<0.001). Finally, restricting analyses to the 11 trials for which pasta intake could be quantified (median pasta intake, 3.33 servings/week [range, 1.75 – 7 servings/week]) showed a similar reduction in body weight (MD= -0.70 kg; 95% CI:-1.10, -0.29 kg; P<0.001) when pasta was consumed in the context of low-GI dietary patterns compared with the higher GI control arms without evidence of heterogeneity ($I^2=0\%$, P-heterogeneity=0.68) (**Supplemental Figure S9**).

Subgroup analyses

We were only able to conduct a priori categorical and continuous subgroup analyses for body weight, BMI, body fat and waist circumference. Subgroup analyses for waist-to-hip ratio and sagittal abdominal diameter could not be assessed, owing to <10 trial comparisons in each case. **Supplemental Figures S10-S12** show the categorical a priori subgroup analyses for body weight. There was no evidence of significant effect modification in any of the subgroup analyses for body weight, including no effect modification of follow-up when comparing studies less than 24 weeks duration to those greater than or equal to 24 weeks (-0.63kg vs -0.57kg, respectively)

(**Supplemental Figure S10**). Neither was there evidence of significant effect modification in any of the subgroup analyses for BMI, body fat or waist circumference (**Supplemental Figures S13-20**).

Supplemental Table S7 and **Supplemental Figures S21-22** show the continuous subgroup analyses for body weight. There was evidence of significant effect modification by carbohydrate and protein intake, where an increase in carbohydrate intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight loss ($\beta = -0.07$, 95% CI: -0.12, -0.01, $I^2 = 0.00\%$, $P = 0.02$), and an increase in protein intake in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with weight gain ($\beta = 0.15$, 95% CI: 0.03, 0.27, $I^2 = 0.00\%$, $P = 0.02$). None of the other continuous subgroup analyses were significant. There was no evidence of significant effect modification in any of the continuous subgroup analyses for BMI (**Supplemental Table S8**). For body fat, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of difference in GI between intervention and control groups, where greater difference in GI between the groups was associated with greater reduction in body fat in the intervention group ($\beta = -0.09$, 95% CI: -0.15, -0.03, $I^2 = 19.39\%$, $P = 0.01$) (**Supplemental Table S9**). None of the other continuous subgroup analyses were significant. For waist circumference, there was evidence of significant effect modification in the continuous meta-regression subgroup analysis of absolute carbohydrate level and absolute protein level, where greater carbohydrate level in the intervention group in which pasta was consumed in the context of low-GI dietary patterns was associated with greater loss in waist circumference ($\beta = -0.11$, 95% CI: -0.19, -0.04, $I^2 = 27.06\%$, $P < 0.01$) and a lower protein level in the intervention group in which pasta was

1
2
3 320 consumed in the context of low-GI dietary patterns was associated with an increase in waist
4
5 321 circumference ($\beta= 0.20$, 95% CI: 0.01, 0.38, $I^2=43.92\%$, $P=0.04$) (**Supplemental Table S10**).
6
7
8 322 None of the other continuous subgroup analyses were significant.
9

10 323
11
12 324 **Dose-response analyses**
13
14 325 **Supplemental Tables S7 and S11 and Supplemental Figure S23** show the dose-response
15
16 326 analysis for the 11 trials for which pasta intake could be quantified. No evidence of a linear dose
17
18 327 response was seen for pasta intake by meta-regression analyses (**Supplemental Table S7**).
19
20 328 There was also no evidence of a non-linear dose response by MKSPLINE ($P=0.85$)
21
22 329 (**Supplemental Figure S23**) or piecewise linear meta-regression analyses (**Supplemental Table**
23
24 330 **S11**).
25
26 331

27
28 332 **Publication Bias**
29
30 333 **Supplemental Figures S24-S27** shows the funnel plots for body weight, BMI, body fat and
31
32 334 waist circumference. There was no evidence of funnel-plot asymmetry. Formal testing with the
33
34 335 Egger and Begg tests did not show evidence of small-study effects ($P>0.05$ for both). Publication
35
36 336 bias was not assessed for waist-to-hip ratio and sagittal abdominal diameter, owing to <10 trial
37
38 337 comparisons.
39
40 338

41
42 339 **GRADE Assessment**
43
44 340 **Supplemental Table S12** shows a summary of the GRADE assessments for the effect of pasta in
45
46 341 the context of low-GI dietary patterns on body weight and measures of adiposity. The evidence
47
48 342 was graded as moderate for body weight, BMI, waist-to-hip ratio and sagittal abdominal
49
50
51
52
53
54
55
56
57
58
59
60

diameter owing to downgrades for indirectness and low for waist circumference and body fat, owing to downgrades for indirectness and inconsistency ($I^2=59\%$, P -heterogeneity <0.001 ; $I^2=66\%$, P -heterogeneity <0.01 , respectively).

DISCUSSION

The present systematic review and meta-analysis quantified the effect of pasta alone and pasta in the context of low-GI dietary patterns on body weight and other markers of adiposity. We failed to identify any trial comparisons for the effect of pasta alone but did identify 32 trial comparisons for the effect of pasta in the context of low-GI dietary patterns in 2448 participants who were predominantly middle-aged and overweight or obese. The primary pooled analysis demonstrated that pasta in the context of low-GI dietary patterns did not contribute to weight gain, resulting in a significant weight loss of -0.63kg when compared to diets higher in GI over a median follow-up of 12 weeks. The lack of harm was reflected in the established clinical secondary outcome measures of global (BMI and body fat) and abdominal adiposity (waist circumference, waist-to-hip ratio and sagittal abdominal diameter). These findings were robust across subgroups. The findings did not differ by metabolic phenotype in those who were overweight or obese or had diabetes, which is noteworthy since these are populations who would benefit from weight management strategies. There was also no effect modification by the energy balance of the design such that the weight loss was seen even under conditions of neutral energy balance (in which participants were instructed to consume dietary advice ad libitum), suggesting that encouragement of the consumption of pasta in the context of a low-GI dietary patterns does not cause harm and may even lead to spontaneous weight loss. There was also no effect modification by follow-up either in continuous meta-regression or categorical, where the 24 trials with

<24weeks follow-up had a weight reduction similar to those 8 trials with ≥ 24 weeks follow-up (-0.63kg vs -0.57kg, respectively). This is of relevance since many dietary studies are successful in demonstrating weight loss in the short term but not over the long term.

Findings in the context of existing studies

We are not aware of any RCTs directly assessing the effect of pasta intake on any health parameters including body weight. Our findings, however, agree with earlier systematic reviews and meta-analyses of RCTs of the effect of low-GI dietary patterns irrespective of pasta intake on body weight and adiposity. The systematic review and meta-analysis by Thomas et al. in 2007 found a significant -1.1kg weight loss and -1.3kg/m² reduction in BMI favouring low-GI or glycemic load (GL) diets compared to control diets in 6 RCTs of 5 weeks to 6 months in duration in overweight or obese individuals¹³. Another systematic review and meta-analysis by Schwingshackl et al. in 2013 found a -0.62kg weight loss favouring low-GI/GL diets compared to higher GI/GL diets in 14 RCTs with greater than 6 months duration in overweight individuals (BMI>25kg/m²)¹⁴.

Our findings also agree with trials in which pasta was emphasized in the context of other healthy dietary patterns. One trial done in children in Spain given Mediterranean dietary advice which included increasing the intake of pasta found that approximately 11.3% of the participants in the Mediterranean diet group who were classified as overweight and obese changed their weight status to normal weight compared to only approximately 2.6% of the participants in the control group⁵⁵.

Other lines of evidence from observational studies have demonstrated benefits of pasta consumption on body weight and adiposity. Pasta intake was recently assessed in the Moli-sani study and the Italian Nutrition & HEalth Survey (INHES), a cross-sectional study of over 20,000 Italians from all over Italy⁵⁶. The study demonstrated that higher pasta consumption was associated with lower BMI, waist circumference and waist-to-hip ratio and with a lower prevalence of overweight and obesity⁵⁶. Furthermore, greater pasta intake was associated with better adherence to the Mediterranean diet, a dietary pattern which has a demonstrated cardiovascular benefit⁵⁷. Similar associations between greater pasta intake and lower body weight have also been observed in prospective U.S. cohorts⁵⁸. A pooled analysis of the 3 Harvard cohorts also showed that higher GI diets (which would preclude pasta) were associated with weight gain⁵⁹.

Although the product form of pasta can vary widely, including in shape (e.g. macaroni, spaghetti, linguine), ingredients (e.g. type of wheat, egg content), and processing technique (e.g. drying temperature), studies have demonstrated that when comparing pastas varying in these parameters, despite slight variations in glycemic response among pastas, glycemic responses are still lower compared to a control, e.g. white bread^{60,61}. One concern of the choice of pasta as a carbohydrate food is that it is a refined food low in fibre. Although there are whole grain pasta options available, studies have demonstrated that fiber added to pasta, does not significantly affect the glucose or insulin response, the secretion of gut hormones or satiety^{62,63}. Furthermore, pasta has a similar glycemic response compared to many fiber-rich carbohydrates, including barley, legumes and steel cut oats, and still a lower glycemic response compared to other fiber-rich foods including whole wheat bread, breakfast cereals like bran flakes and potatoes with

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

412 skin⁶⁴. The typically consumed white wheat pasta also has a higher micronutrient content
413 compared to other white wheat products like bread since it contains the aleurone layer which is
414 preserved as a result of the use of harder wheats (durum wheat); even when durum wheats are
415 used in breads, pasta retains a lower glycemic response primarily because of the processing
416 techniques used in pasta making which give pasta a compact structure and reduced starch
417 hydrolysis⁶¹.

418
419 The mechanism by which pasta in the context of low-GI dietary patterns lead to weight loss even
420 under conditions of ad libitum dietary advice is unclear. Lower GI diets may result in greater
421 body weight reduction compared to higher GI diets because lower GI foods may be more
422 satiating⁶⁵ and have been shown to delay hunger and decrease subsequent energy intake¹³. Low-
423 GI dietary patterns are also characterized by high fiber content^{64,66} which may also contribute to
424 improvements in satiety and hunger¹⁷. Furthermore, studies which have compared ad libitum
425 low-GI dietary patterns to standard energy-restricted low-fat diets, demonstrated equivalency or
426 better weight loss when following the low-GI diet, despite the fact that they could eat as much as
427 they desired^{13,67}. Thus, voluntary energy intake may be lower after low-GI meals, as has been
428 previously demonstrated⁶⁸.

429
430 **Strengths and limitations**

431 The strengths of the present systematic review and meta-analysis include that it is
432 comprehensive, includes RCTs which protects against bias and uses the GRADE approach to
433 evaluate the quality of evidence. Additionally, a large number of trials were identified (32 trials)
434 for the primary outcome of body weight, the median follow-up period was 12 weeks which

allows for the assessment of a moderate duration of intervention, none of the trials were rated as having a serious risk of bias, and there was no evidence of publication bias.

There are several limitations. First, we downgraded the certainty of the evidence for serious inconsistency in the treatment estimates across trials for some of the outcomes assessed. There was evidence of unexplained heterogeneity in waist circumference ($I^2=62\%$) and in body fat ($I^2=65\%$). Although the inconsistency in these outcomes may have related to measurement error⁶⁹ in the different techniques for measuring waist circumference and body fat, we were unable to conduct sensitivity or subgroup analyses to explore this source of heterogeneity. Second, we downgraded the certainty of the evidence for serious indirectness. Most of the available trials did not quantify the amount of pasta consumed in the context of the low-GI dietary patterns. Although sensitivity analyses in which analyses were restricted to the 11 trials that did quantify (providing a median 3.33 servings/week) pasta intake did not meaningfully alter our estimates (-0.70kg versus -0.63kg), it is difficult to quantify the effect of pasta in these diets. There is also the question of indirectness in the translation to other background diets. None of the available trials evaluated the effect of pasta alone or in the context of other dietary patterns. Whether the observed effect of pasta in the context of low-GI dietary patterns will hold in the context of other healthy dietary patterns, such as Mediterranean and Vegetarian dietary patterns, is unclear. Although there is no biological reason to doubt that the findings would hold across different dietary patterns, there was no direct evidence to support this conclusion. If the question had been asked from the perspective of benefit as opposed to that of harm, then the relatively short duration of the included trials is another reason to downgrade for serious indirectness. In the absence of long-term trials (>1 year diet duration), it is difficult to conclude with certainty

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

458 that the observed lack of harm implies an actual sustainable benefit. Finally, there was some
459 evidence of imprecision for benefit. Whereas the 95% CI of the pooled estimates did not overlap
460 with our pre-specified MID for harm (that is, they did not contain evidence for harm) and so
461 were not downgraded for imprecision, the upper bound of the 95% CI did overlap with the lower
462 bound of the same MID to assess the precision of the evidence for benefit for some outcomes.
463
464 Balancing these strengths and limitations, the GRADE approach assessed the overall quality and
465 strength of the available evidence of the effect of pasta in the context of low-GI dietary patterns
466 as moderate for the primary outcome of body weight and the secondary outcomes of BMI, waist-
467 to-hip ratio and sagittal abdominal diameter owing to downgrades for indirectness. The evidence
468 was assessed as low for the other secondary outcomes of body fat and waist circumference
469 owing to downgrades for indirectness and inconsistency.

471 **Implications**

472 These results are important considering the negative messages directed at the public regarding
473 carbohydrates, which is influencing their food choices, as is evident in recent reductions in
474 carbohydrate intake⁷⁰⁻⁷², and in particular reductions in pasta consumption^{70,73-76}. Contrary to
475 these concerns, the available evidence shows that when pasta is consumed in the context of low-
476 GI dietary patterns that there is not weight gain but rather marginally clinically significant weight
477 loss (>0.5kg)⁷⁷.

479 Although we were able to approximate the amount of pasta consumed in one third of included
480 trials, it is unclear what the effect of pasta would be in the context of other dietary patterns. Low-

GI dietary patterns, however, shares many similarities with a Mediterranean diet, which emphasizes many low-GI foods and has demonstrated a CVD benefit⁵⁷.

Current clinical practice guidelines already suggest the replacement of high GI foods with low-GI foods for improvement of glycemic control and cardiovascular risk factors^{78,79}. The present evidence means that pasta may be highlighted as an important example of a low-GI food which can contribute to a low-GI dietary pattern, a pattern which in turn may potentially improve cardio-metabolic risk without an adverse effect on weight control.

CONCLUSIONS

In conclusion, the available evidence from RCTs does not allow us to conclude that pasta consumed in the context of low-GI dietary patterns has an adverse effect on body weight and adiposity outcomes of importance in the prevention and management of overweight and obesity. On the contrary, pasta in the context of low-GI dietary patterns reduces body weight and BMI compared with higher-GI dietary patterns. The results are generalizable in the context of a high carbohydrate dietary pattern composed of low-GI foods with or without the intention of weight loss in middle-aged individuals who are overweight or obese or have diabetes. Although the clinical significance of the observed weight loss is debatable, this finding increases our confidence that pasta in the context of low-GI dietary patterns does not result in weight gain. Further research may change our confidence in the estimates for our primary outcome body weight and several key secondary outcomes including BMI and two measures of abdominal adiposity, waist-to-hip ratio and sagittal abdominal diameter. More research is needed, to improve our estimates for the secondary outcomes, body fat and waist circumference and assess

1
2
3 504 whether our findings extend to related cardio-metabolic outcomes. There is also a need for more
4
5 505 randomized trials of >1 year diet duration to clarify whether the lack of harm for pasta in the
6
7
8 506 context of low-GI dietary patterns will translate into meaningful long-term benefits. Other
9
10 507 randomized trials should focus on whether pasta will have similar effects in the context of other
11
12 508 healthy dietary patterns such as a Mediterranean diet.
13
14
15
16 509

17
18 510 **Acknowledgements**

19
20 511 We wish to thank Teruko Kishibe, an information specialist in the Scotiabank Health Sciences
21
22 512 Library, St Michael's Hospital, for her help in the development of search terms used. Aspects of
23
24 513 this work were presented at the Società Italiana di Nutrizione Umana (SINU) Congress, in
25
26 514 Parma, Italy, on April 12, 2017 and at the 35th International Symposium on Diabetes and
27
28 515 Nutrition, of the Diabetes and Nutrition Study Group (DNSG) of the European Association for
29
30 516 the Study of Diabetes (EASD) in Skagen, Denmark, on June 20 and 21, 2017.
31
32
33

34 517 **Data Sharing**

35
36
37 518 No additional data available.
38
39
40 519

41
42
43 520 **Exclusive License**

44
45
46 521 The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf
47
48 522 of all authors, an exclusive license (or non-exclusive for government employees) on a worldwide
49
50 523 basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in BMJ
51
52 524 editions and any other BMJ PGL products and sublicenses such use and exploit all subsidiary
53
54 525 rights, as set out in our license.
55
56
57
58
59
60

526 **Ethics Approval**

527 Not required.

528 **Authorship:**

529 All authors had full access to all of the data (including statistical reports and tables) in this study
530 and take full responsibility for the integrity of the data and the accuracy of the data analysis.

531 **Conception and design:** C.W.C. Kendall, J.L. Sievenpiper.

532 **Analysis and interpretation of the data:** L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S.

533 Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

534 **Drafting of the article:** L. Chiavaroli, J.L. Sievenpiper.

535 **Critical revision of the article for important intellectual content:** L. Chiavaroli, C.W.C.

536 Kendall, C.R. Braunstein, S. Blanco Mejia, L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

537 **Final approval of the article:** L. Chiavaroli, C.W.C. Kendall, C.R. Braunstein, S. Blanco Mejia,

538 L.A. Leiter, D.J.A. Jenkins, J.L. Sievenpiper.

539 **Obtaining of funding:** C.W.C. Kendall, D.J.A. Jenkins, J.L. Sievenpiper.

540 **Administrative, technical, or logistic support:** S. Blanco Mejia.

541 **Collection and assembly of data:** L. Chiavaroli, C.R. Braunstein, S. Blanco Mejia.

542 **Guarantor:** J.L. Sievenpiper.

543 **Transparency declaration:** As guarantor, J.L. Sievenpiper affirms that the manuscript is an

544 honest, accurate, and transparent account of the study being reported; that no important aspects

of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Funding:

This work was funded by the Canadian Institutes of Health Research (funding reference number, 129920) through the Canada-wide Human Nutrition Trialists' Network (NTN). The Diet, Digestive tract, and Disease (3-D) Centre, funded through the Canada Foundation for Innovation (CFI) and the Ministry of Research and Innovation's Ontario Research Fund (ORF), provided the infrastructure for the conduct of this project. **L Chiavaroli** was funded by a Toronto 3D Postdoctoral Fellowship Award. **DJA Jenkins** was funded by the Government of Canada through the Canada Research Chair Endowment. **JL Sievenpiper** was funded by a PSI Graham Farquharson Knowledge Translation Fellowship, Canadian Diabetes Association (CDA) Clinician Scientist award, CIHR INMD/CNS New Investigator Partnership Prize, and Banting & Best Diabetes Centre Sun Life Financial New Investigator Award. None of the sponsors had a role in any aspect of the present study, including design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, approval of the manuscript or decision to publish.

Conflicts of Interest:

All authors have completed the Unified Competing Interest form (available on request from the corresponding author) and declare: **CWC Kendall** has received research support from the Advanced Foods and Material Network, Agrifoods and Agriculture Canada, the Almond Board of California, the American Pistachio Growers, Barilla, the California Strawberry Commission, the Calorie Control Council, CIHR, the Canola Council of Canada, the Coca-Cola Company (investigator initiated, unrestricted grant), Hain Celestial, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Kraft, Loblaw Companies Ltd., Orafit, Pulse Canada, Saskatchewan Pulse Growers, Solae and Unilever. He has received travel funding, consultant fees and/or honoraria from Abbott Laboratories, the Almond Board of California, the American Peanut Council, the American Pistachio Growers, Barilla, Bayer, the Canola Council

of Canada, the Coca-Cola Company, Danone, General Mills, the International Tree Nut Council Nutrition Research and Education Foundation, Kellogg, Loblaw Companies Ltd., the Nutrition Foundation of Italy, Oldways Preservation Trust, Orafiti, Paramount Farms, the Peanut Institute, PepsiCo, Pulse Canada, Sabra Dipping Co., Saskatchewan Pulse Growers, Solae, Sun-Maid, Tate and Lyle, and Unilever. He is on the Clinical Practice Guidelines Expert Committee for Nutrition Therapy of the European Association for the Study of Diabetes (EASD) and has served on the scientific advisory board for the Almond Board of California, the International Tree Nut Council, Oldways Preservation Trust, Paramount Farms and Pulse Canada. **DJAJ** has received research grants from Saskatchewan Pulse Growers, the Agricultural Bioproducts Innovation Program through the Pulse Research Network, the Advanced Foods and Material Network, Loblaw Companies Ltd., Unilever, Barilla, the Almond Board of California, Agriculture and Agri-food Canada, Pulse Canada, Kellogg's Company, Canada, Quaker Oats, Canada, Procter & Gamble Technical Centre Ltd., Bayer Consumer Care, Springfield, NJ, Pepsi/Quaker, International Nut & Dried Fruit (INC), Soy Foods Association of North America, the Coca-Cola Company (investigator initiated, unrestricted grant), Solae, Haine Celestial, the Sanitarium Company, Orafiti, the International Tree Nut Council Nutrition Research and Education Foundation, the Peanut Institute, the Canola and Flax Councils of Canada, the Calorie Control Council (CCC), the CIHR, the Canada Foundation for Innovation and the Ontario Research Fund. He has received in-kind supplies for trial as a research support from the Almond board of California, Walnut Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo, Loblaw Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada, WhiteWave Foods. He has been on the speaker's panel, served on the scientific advisory board and/or received travel support and/or honoraria from the Almond Board of California, Canadian Agriculture Policy Institute, Loblaw Companies Ltd, the Griffin Hospital (for the development of the NuVal scoring system, the Coca-Cola Company, EPICURE, Danone, Diet Quality Photo Navigation (DQPN), FareWell, Verywell, True Health Initiative, Saskatchewan Pulse Growers, Sanitarium Company, Orafiti, the Almond Board of California, the American Peanut Council, the International Tree Nut Council Nutrition Research and Education Foundation, the Peanut Institute, Herbalife International, Pacific Health Laboratories, Nutritional Fundamental for Health, Barilla, Metagenics, Bayer Consumer Care, Unilever Canada and Netherlands, Solae, Kellogg, Quaker Oats, Procter & Gamble, the Coca-Cola Company, the Griffin Hospital, Abbott Laboratories, the Canola Council of Canada, Dean Foods, the California Strawberry Commission, Haine Celestial, PepsiCo, the Alpro Foundation, Pioneer Hi-Bred International, DuPont Nutrition and Health, Spherix Consulting and WhiteWave Foods, the Advanced Foods and Material Network, the Canola and Flax Councils of Canada, the Nutritional Fundamentals for Health, Agri-Culture and Agri-Food Canada, the Canadian Agri-Food Policy Institute, Pulse Canada, the Saskatchewan Pulse Growers, the Soy Foods Association of North America, the Nutrition Foundation of Italy (NFI), Nutra-Source Diagnostics, the McDougall Program, the Toronto Knowledge Translation Group (St. Michael's Hospital), the Canadian College of Naturopathic Medicine, The Hospital for Sick Children, the Canadian Nutrition Society (CNS), the American Society of Nutrition (ASN), Arizona State University, Paolo Sorbini Foundation and the Institute of Nutrition, Metabolism and Diabetes. He received an honorarium from the United States Department of Agriculture to present the 2013 W.O. Atwater Memorial Lecture. He received the 2013 Award for Excellence in Research from the International Nut and Dried Fruit Council. He received funding and travel support from the Canadian Society of Endocrinology and Metabolism to produce mini cases for the Canadian Diabetes Association

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(CDA). He is a member of the International Carbohydrate Quality Consortium (ICQC). His wife, ALJ, is a director and partner of Glycemic Index Laboratories, Inc., and his sister received funding through a grant from the St. Michael's Hospital Foundation to develop a cookbook for one of his studies. **JL Sievenpiper** has received research support from the Canadian Institutes of health Research (CIHR), Canadian Diabetes Association (CDA), PSI Foundation, Banting and Best Diabetes Centre (BBDC), Canadian Nutrition Society (CNS), American Society for Nutrition (ASN), Calorie Control Council, INC International Nut and Dried Fruit Council Foundation, National Dried Fruit Trade Association, The Coca-Cola Company (investigator initiated, unrestricted), Dr. Pepper Snapple Group (investigator initiated, unrestricted), The Tate and Lyle Nutritional Research Fund at the University of Toronto, and The Alberta Pulse Growers Glycemic Control and Cardiovascular Disease in Type 2 Diabetes Fund at the University of Toronto. He has received in-kind research support from the Almond board of California, Walnut Council of California, American Peanut Council, Barilla, Unilever, Unico, Primo, Loblaw Companies, Quaker (Pepsico), Pristine Gourmet, Bunge Limited, Kellogg Canada, WhiteWave Foods. He has received reimbursement of travel expenses, speaker fees, and/or honoraria from the Canadian Diabetes Association (CDA), Canadian Nutrition Society (CNS), University of Alabama at Birmingham, Oldways Preservation Trust, Nutrition Foundation of Italy (NFI), Diabetes and Nutrition Study Group (DNSG) of the European Association for the Study of Diabetes (EASD), International Life Sciences Institute (ILSI) North America, Dr. Pepper Snapple Group, Corn Refiners Association, World Sugar Research Organization, Dairy Farmers of Canada, Società Italiana di Nutrizione Umana (SINU), III World Congress of Public Health Nutrition, C3 Collaborating for Health, Sprim Brasil, WhiteWave Foods, Rippe Lifestyle, mdBriefcase, Federation of European Nutrition Societies (FENS), New York Academy of Sciences, International Diabetes Federation, American Heart Association (AHA), American Society for Nutrition (ASN), FoodMinds LLC, Memac Ogilvy & Mather LLC, Pulse Canada, Pepsico, BCFN Foundation, The Ginger Network LLC, and Dietitians of Canada. He has ad hoc consulting arrangements with Winston & Strawn LLP, Perkins Coie LLP, and Tate & Lyle. He is a member of the European Fruit Juice Association Scientific Expert Panel. He is on Clinical Practice Guidelines Expert Committees of the Canadian Diabetes Association (CDA) European Association for the study of Diabetes (EASD), and Canadian Cardiovascular Society (CCS), as well as an expert writing panel of the American Society for Nutrition (ASN). He serves as an unpaid scientific advisor for the Food, Nutrition, and Safety Program (FNSP) and the Technical Committee on Carbohydrates of the International Life Science Institute (ILSI) North America. He is a member of the International Carbohydrate Quality Consortium (ICQC), Executive Board Member of the Diabetes and Nutrition Study Group (DNSG) of the EASD, and Director of the Toronto 3D Knowledge Synthesis and Clinical Trials foundation. His wife is an employee of Unilever Canada. No competing interests were declared by **CR Braunstein, S Blanco Mejia, and LA Leiter.**

References

1. Malhotra A. Eating MORE fat while cutting carbs and quitting sugar can help you lose weight and be happier, says top cardiologist. Mail Online 2016.
2. Taubes G. Diet Advice That Ignores Hunger. The New York Times. Sunday Review ed 2015.
3. Ludwig D. Always hungry? Conquer cravings, retrain your fat cells and lose weight permanently. New York: Grand Central Life & Style; 2016.
4. Taubes G. Why We Get Fat And What To Do About It. . New York: Alfred A Knopf; 2011.
5. Atkins R. Dr Atkins' Diet Revolution: The High Calorie Way to Stay Thin Forever. : Bantam Books; 1973.
6. Perlmutter D. Grain Brain: The surprising truth about wheat, carbs, and sugar -- Your brain's silent killers: Little, Brown And Company; 2013.
7. David W. Wheat Belly: HarperCollins; 2012.
8. Cordain L. The Paleo Diet Revised: Lose weight and get healthy by eating the foods you were designed to eat: Houghton Mifflin Harcourt; 2010.
9. Teicholz N. The Big Fat Surprise: Why Butter, Meat and Cheese Belong in a Healthy Diet: Simon and Schuster; 2014.
10. National Obesity Forum, The Public Health Collaboration. Eat Fat, Cut The Carbs and Avoid Snacking To Reverse Obesity and Type 2 Diabetes.
11. Malhotra A. Saturated fat is not the major issue. BMJ 2013;347:f6340.
12. Malhotra A. Author's reply to Lichtenstein, Tedstone and Pyne, Mann and colleagues, Lim, and Clifton. BMJ 2013;347:f6855.
13. Thomas DE, Elliott EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. Cochrane Database Syst Rev 2007;CD005105.
14. Schwingshackl L, Hoffmann G. Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. Nutr Metab Cardiovasc Dis 2013;23:699-706.
15. Pol K, Christensen R, Bartels EM, Raben A, Tetens I, Kristensen M. Whole grain and body weight changes in apparently healthy adults: a systematic review and meta-analysis of randomized controlled studies. Am J Clin Nutr 2013;98:872-84.
16. Harland JI, Garton LE. Whole-grain intake as a marker of healthy body weight and adiposity. Public Health Nutr 2008;11:554-63.
17. Slavin JL. Dietary fiber and body weight. Nutrition 2005;21:411-8.
18. Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: Health Implications of Dietary Fiber. J Acad Nutr Diet 2015;115:1861-70.
19. Higgins JP, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions [Internet]. Version 5.1.0. The Cochrane Collaboration. Version current March 2011 [cited 2015 Sep 23]. Available from: www.cochrane-handbook.org.
20. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Ann Intern Med 2009;151:264-9, W64.
21. Elbourne DR, Altman DG, Higgins JP, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. Int J Epidemiol 2002;31:140-9.
22. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ 1997;315:629-34.
23. Begg CB. A measure to aid in the interpretation of published clinical trials. Stat Med 1985;4:1-9.
24. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 2000;56:455-63.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

25. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383-94.

26. Jebb SA, Lovegrove JA, Griffin BA, et al. Effect of changing the amount and type of fat and carbohydrate on insulin sensitivity and cardiovascular risk: the RISCK (Reading, Imperial, Surrey, Cambridge, and Kings) trial. *Am J Clin Nutr* 2010;92:748-58.

27. Philippou E, Bovill-Taylor C, Rajkumar C, et al. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 2009;58:1703-8.

28. Philippou E, McGowan BM, Brynes AE, Dornhorst A, Leeds AR, Frost GS. The effect of a 12-week low glycaemic index diet on heart disease risk factors and 24 h glycaemic response in healthy middle-aged volunteers at risk of heart disease: a pilot study. *Eur J Clin Nutr* 2008;62:145-9.

29. Sichieri R, Moura AS, Genelhu V, Hu F, Willett WC. An 18-mo randomized trial of a low-glycemic-index diet and weight change in Brazilian women. *Am J Clin Nutr* 2007;86:707-13.

30. Wolever TM, Mehling C. High-carbohydrate-low-glycaemic index dietary advice improves glucose disposition index in subjects with impaired glucose tolerance. *Br J Nutr* 2002;87:477-87.

31. Pereira EV, Costa Jde A, Alfenas Rde C. Effect of glycemic index on obesity control. *Arch Endocrinol Metab* 2015;59:245-51.

32. Karl JP, Roberts SB, Schaefer EJ, et al. Effects of carbohydrate quantity and glycemic index on resting metabolic rate and body composition during weight loss. *Obesity (Silver Spring)* 2015;23:2190-8.

33. Buscemi S, Cosentino L, Rosafio G, et al. Effects of hypocaloric diets with different glycemic indexes on endothelial function and glycemic variability in overweight and in obese adult patients at increased cardiovascular risk. *Clin Nutr* 2013;32:346-52.

34. Larsen TM, Dalskov SM, van Baak M, et al. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med* 2010;363:2102-13.

35. Solomon TP, Haus JM, Kelly KR, et al. A low-glycemic index diet combined with exercise reduces insulin resistance, postprandial hyperinsulinemia, and glucose-dependent insulinotropic polypeptide responses in obese, prediabetic humans. *Am J Clin Nutr* 2010;92:1359-68.

36. Philippou E, Neary NM, Chaudhri O, et al. The effect of dietary glycemic index on weight maintenance in overweight subjects: a pilot study. *Obesity (Silver Spring)* 2009;17:396-401.

37. Abete I, Parra D, Martinez JA. Energy-restricted diets based on a distinct food selection affecting the glycemic index induce different weight loss and oxidative response. *Clin Nutr* 2008;27:545-51.

38. Aston LM, Stokes CS, Jebb SA. No effect of a diet with a reduced glycaemic index on satiety, energy intake and body weight in overweight and obese women. *Int J Obes (Lond)* 2008;32:160-5.

39. Jensen L, Sloth B, Krog-Mikkelsen I, et al. A low-glycemic-index diet reduces plasma plasminogen activator inhibitor-1 activity, but not tissue inhibitor of proteinases-1 or plasminogen activator inhibitor-1 protein, in overweight women. *Am J Clin Nutr* 2008;87:97-105.

40. Bellisle F, Dalix AM, De Assis MA, et al. Motivational effects of 12-week moderately restrictive diets with or without special attention to the Glycaemic Index of foods. *Br J Nutr* 2007;97:790-8.

41. de Rougemont A, Normand S, Nazare JA, et al. Beneficial effects of a 5-week low-glycaemic index regimen on weight control and cardiovascular risk factors in overweight non-diabetic subjects. *Br J Nutr* 2007;98:1288-98.

42. McMillan-Price J, Petocz P, Atkinson F, et al. Comparison of 4 diets of varying glycemic load on weight loss and cardiovascular risk reduction in overweight and obese young adults: a randomized controlled trial. *Arch Intern Med* 2006;166:1466-75.

43. Visek J, Lacigova S, Cechurova D, Rusavy Z. Comparison of a low-glycemic index vs standard diabetic diet. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:112-6.

44. Jenkins DJ, Kendall CW, Vuksan V, et al. Effect of lowering the glycemic load with canola oil on glycemic control and cardiovascular risk factors: a randomized controlled trial. *Diabetes Care* 2014;37:1806-14.
45. Jenkins DJ, Kendall CW, Augustin LS, et al. Effect of legumes as part of a low glycemic index diet on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 2012;172:1653-60.
46. Yusof BN, Talib RA, Kamaruddin NA, Karim NA, Chinna K, Gilbertson H. A low-GI diet is associated with a short-term improvement of glycaemic control in Asian patients with type 2 diabetes. *Diabetes Obes Metab* 2009;11:387-96.
47. Wolever TM, Gibbs AL, Mehling C, et al. The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on glycated hemoglobin but reduction in C-reactive protein. *Am J Clin Nutr* 2008;87:114-25.
48. Jenkins DJ, Kendall CW, McKeown-Eyssen G, et al. Effect of a low-glycemic index or a high-cereal fiber diet on type 2 diabetes: a randomized trial. *JAMA* 2008;300:2742-53.
49. Jimenez-Cruz A, Bacardi-Gascon M, Turnbull WH, Rosales-Garay P, Severino-Lugo I. A flexible, low-glycemic index mexican-style diet in overweight and obese subjects with type 2 diabetes improves metabolic parameters during a 6-week treatment period. *Diabetes Care* 2003;26:1967-70.
50. Heilbronn LK, Noakes M, Clifton PM. The effect of high- and low-glycemic index energy restricted diets on plasma lipid and glucose profiles in type 2 diabetic subjects with varying glycemic control. *J Am Coll Nutr* 2002;21:120-7.
51. Fontvieille AM, Acosta M, Rizkalla SW, et al. A moderate switch from high to low glycemic-index foods for 3 weeks improves metabolic control of type I (IDDM) diabetic subjects. *Diabetes Nutr Metab* 1988;1:139-43.
52. Frost GS, Brynes AE, Bovill-Taylor C, Dornhorst A. A prospective randomised trial to determine the efficacy of a low glycaemic index diet given in addition to healthy eating and weight loss advice in patients with coronary heart disease. *Eur J Clin Nutr* 2004;58:121-7.
53. de Assis Costa J, de Cassia Goncalves Alfensa R. The consumption of low glycemic meals reduces abdominal obesity in subjects with excess body weight. *Nutricion Hospitalaria* 2012;27:1178-83.
54. Fontvieille AM, Rizkalla SW, Penfornis A, Acosta M, Bornet FR, Slama G. The use of low glycaemic index foods improves metabolic control of diabetic patients over five weeks. *Diabet Med* 1992;9:444-50.
55. Bibiloni MDM, Fernandez-Blanco J, Pujol-Plana N, et al. [Improving diet quality in children through a new nutritional education programme: INFADIMED]. *Gac Sanit* 2017.
56. Pounis G, Castelnuovo AD, Costanzo S, et al. Association of pasta consumption with body mass index and waist-to-hip ratio: results from Moli-sani and INHES studies. *Nutr Diabetes* 2016;6:e218.
57. Estruch R, Ros E, Salas-Salvado J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013;368:1279-90.
58. Shay CM, Van Horn L, Stamler J, et al. Food and nutrient intakes and their associations with lower BMI in middle-aged US adults: the International Study of Macro-/Micronutrients and Blood Pressure (INTERMAP). *Am J Clin Nutr* 2012;96:483-91.
59. Smith JD, Hou T, Ludwig DS, et al. Changes in intake of protein foods, carbohydrate amount and quality, and long-term weight change: results from 3 prospective cohorts. *Am J Clin Nutr* 2015;101:1216-24.
60. Granfeldt Y, Bjorck I. Glycemic response to starch in pasta: a study of mechanisms of limited enzyme availability. *J Cereal Sci* 1991;14:47-61.
61. Granfeldt Y, Bjorck I, Hagander B. On the importance of processing conditions, product thickness and egg addition for the glycaemic and hormonal responses to pasta: a comparison with bread made from 'pasta ingredients'. *Eur J Clin Nutr* 1991;45:489-99.

1
2
3 798 62. Frost GS, Brynes AE, Dhillon WS, Bloom SR, McBurney MI. The effects of fiber enrichment of pasta
4 799 and fat content on gastric emptying, GLP-1, glucose, and insulin responses to a meal. *Eur J Clin Nutr*
5 800 2003;57:293-8.
6 801 63. Korczak R, Timm D, Ahnen R, Thomas W, Slavin JL. High Protein Pasta is Not More Satiating than
7 802 High Fiber Pasta at a Lunch Meal, Nor Does it Decrease Mid-Afternoon Snacking in Healthy Men and
8 803 Women. *J Food Sci* 2016;81:S2240-5.
9 804 64. Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and
10 805 glycemic load values: 2008. *Diabetes Care* 2008;31:2281-3.
11 806 65. Ludwig DS. Dietary glycemic index and obesity. *J Nutr* 2000;130:280S-3S.
12 807 66. Riccardi G, Rivellese AA, Giacco R. Role of glycemic index and glycemic load in the healthy state,
13 808 in prediabetes, and in diabetes. *Am J Clin Nutr* 2008;87:269S-74S.
14 809 67. Saris WH, Astrup A, Prentice AM, et al. Randomized controlled trial of changes in dietary
15 810 carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the
16 811 CARMEN study. The Carbohydrate Ratio Management in European National diets. *Int J Obes Relat*
17 812 *Metab Disord* 2000;24:1310-8.
18 813 68. Ludwig DS, Majzoub JA, Al-Zahrani A, Dallal GE, Blanco I, Roberts SB. High glycemic index foods,
19 814 overeating, and obesity. *Pediatrics* 1999;103:E26.
20 815 69. Wang J, Thornton JC, Bari S, et al. Comparisons of waist circumferences measured at 4 sites. *Am*
21 816 *J Clin Nutr* 2003;77:379-84.
22 817 70. Makarem N, Scott M, Quatromoni P, Jacques P, Parekh N. Trends in dietary carbohydrate
23 818 consumption from 1991 to 2008 in the Framingham Heart Study Offspring Cohort. *Br J Nutr*
24 819 2014;111:2010-23.
25 820 71. Rehm CD, Penalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults, 1999-2012.
26 821 *JAMA* 2016;315:2542-53.
27 822 72. Golzarand M, Mirmiran P, Jessri M, Toolabi K, Mojarad M, Azizi F. Dietary trends in the Middle
28 823 East and North Africa: an ecological study (1961 to 2007). *Public Health Nutr* 2012;15:1835-44.
29 824 73. Leclercq C, Arcella D, Piccinelli R, et al. The Italian National Food Consumption Survey INRAN-
30 825 SCAI 2005-06: main results in terms of food consumption. *Public Health Nutr* 2009;12:2504-32.
31 826 74. Bonaccio M, Di Castelnuovo A, Bonanni A, et al. Decline of the Mediterranean diet at a time of
32 827 economic crisis. Results from the Moli-sani study. *Nutr Metab Cardiovasc Dis* 2014;24:853-60.
33 828 75. Sofi F, Innocenti G, Dini C, et al. Low adherence of a clinically healthy Italian population to
34 829 nutritional recommendations for primary prevention of chronic diseases. *Nutr Metab Cardiovasc Dis*
35 830 2006;16:436-44.
36 831 76. Ferdman RA. This is what happens when everyone is terrified of carbs. *The Washington Post*
37 832 2015 September 2, 2015.
38 833 77. Johnston BC, Kanters S, Bandayrel K, et al. Comparison of weight loss among named diet
39 834 programs in overweight and obese adults: a meta-analysis. *JAMA* 2014;312:923-33.
40 835 78. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of
41 836 blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American
42 837 College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*
43 838 2014;129:S1-45.
44 839 79. Anderson TJ, Gregoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society
45 840 guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease
46 841 in the adult. *The Canadian journal of cardiology* 2013;29:151-67.
47 842
48 843

Figure Legend

Figure 1: Literature Search

Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). n = 2448. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

Table 1: Summary of trial characteristics

Trial Characteristics*	ALL	Neutral Energy Balance	Negative Energy Balance
Trial Number (n)	32	23	9
Trial Size (total, range)	2448 (8 - 250)	1989 (8 - 250)	459 (13 - 123)
Male: Female^a (%)	40:60	47:53	27:73
Age (years)	50 (40 - 58)	52.0 (42.1 – 59.5)	49.5 (34.2 – 53.0)
Metabolic Phenotype (OW/OB:DM:CHD) (%)	66:31:3	57:39:4	89:11:0
Setting (IP:OP) (%)	3:97	4:96	0:100
Baseline Body Weight (kg)^b	85.5 (80.0 - 91.9)	84.1 (79.5 – 87.5)	92.5 (86.1 – 93.9)
Baseline BMI (kg/m²)^c	30.4 (28.2 - 32.0)	29.5 (27.4 – 31.4)	31.7 (30.1 – 32.9)
Study Design (C:P) (%)	19:81	26:74	0:100
Dose Pasta (servings/week)^d	3.3 (2.3 - 3.5)	3.4 (2.9 – 4.1)	2.3 (2.3-3.5)
GI in Pasta/LGI group	49.0 (44.0 - 55.1)	46.5 (49.9 – 55.5)	44.0 (42.3 - 49.4)
GI in Higher GI group	62.5 (61.6 - 63.2)	63.3 (60.1 – 64.4)	61.0 (59.2 – 66.6)
Calorie reduction in Pasta/LGI group (kcal)^e	-179 (-90 - -448)	-165 (-74 - -313)	-447 (-134 - -594)
Calorie reduction in Higher GI group (kcal)^e	-181 (-93 - -401)	-160 (-40 - -248)	-470 (-172 - -561)
Feeding Control (Met:Supp:DA) (%)	6:44:50	4:48:48	11:33:56
Follow-Up Duration (Weeks)	12 (9-21)	12 (6-24)	12 (10 - 21)
Funding Sources (A:I:AI:NR) (%)	47:9:25:19	44:13:26:17	56:0:22:22

* median (inter quartile range), unless otherwise indicated

^a24/32 trials provided data on sex

^b 30/32 trials reported baseline body weight

^c 28/32 trials reported baseline BMI

^d 11/32 trials provided data from which dose could be approximated

^e 20/32 trials provided data from which to approximate changes in caloric intake

A, agency; AI, agency and industry; BMI, body mass index; C, crossover design; CHD, coronary heart disease; DA, dietary advice; DM, diabetes; GI, glycemic index; I, industry; IP, inpatient; LGI, low glycemic index; Met, metabolic; NR, not reported; OB, obese; OP, outpatient; OW, overweight; P, parallel design; Suppl, supplemented/provision of certain food

For peer review only




Figure 1: Literature Search
40x54mm (300 x 300 DPI)




Figure 2: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg). $n = 2448$. Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.

40x54mm (300 x 300 DPI)

For peer review only




Figure 3: Plot of pooled effect estimates from randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on global and abdominal markers of adiposity. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I² statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

BMI; body mass index; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet.

40x54mm (300 x 300 DPI)

Online Supplemental Information

Table of Contents

Supplemental Tables..... 4

Supplemental Table S1: PRISMA Checklist 4

Supplemental Table S2: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials 7

Supplemental Table S3: PICO framework of the search strategy..... 9

Supplement Table S4a: Trial characteristics 10

Supplement Table S4b: Trial characteristics continued..... 11

Supplement Table S4c: Trial characteristics continued 12

Supplement Table S4d: Trial characteristics continued..... 13

Supplement Table S4e: Trial characteristics continued..... 14

Supplemental Table S5: Sensitivity analysis of the systematic removal of each trial* 15

Supplemental Table S6: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 for crossover trials 18

Supplement Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹..... 19

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹..... 21

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹..... 23

Supplemental Table S10. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹..... 25

Supplementary Table S11. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg). 27

Supplementary Table S12: GRADE assessment of study quality 28

Supplemental Figures 32

Supplemental Figure S1: Cochrane risk of bias summary for all included trials..... 32

Supplemental Figure S2: Risk of bias proportion graph for all included trials..... 33

Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038) 34

Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285) 35

Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	36
Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)	37
Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)	38
Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448) ..	39
Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)	40
Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	41
Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)	43
Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)	45
Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	47
Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) continued (n = 1038)	49
Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m ²) (n = 1038)	51
Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	53
Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)	55
Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)	57
Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)	59
Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CIs) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380). 61	
Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)	63

Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19) 64

Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure) 65

Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg) 66

Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²) 67

Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%) 68

Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) 69

Supplemental Tables

Supplemental Table S1: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5-6, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5-6, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6

Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6, Supplemental Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6-7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7-8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ²) for each meta-analysis.	8
Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7-10
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10-11, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	11, Supplemental Figures S1-S2
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	11-13, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	11-13, Figures 2-3, Supplemental Figures S3-S7
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	11,15 Supplemental

			Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	12-16, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	16-22
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	19-21
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	22-23
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	26-31

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Supplemental Table S2: Search strategy for studies assessing the effect of pasta in the context of low-GI dietary patterns on body weight in randomized controlled trials

Database	Search Period	Search Terms
Medline	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw. 17. overweight.tw. 18. weight*.tw. 19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
Embase	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycaemic index.tw. 8. glycemic index.tw. 9. glycaemic ind*.tw. 10. glycemic ind*.tw. 11. glycemic load*.tw. 12. glycaemic load*.tw. 13. glycemic index/ 14. body mass index/ 15. body mass index.tw. 16. BMI.tw. 17. overweight.tw. 18. weight*.tw.

		19. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 20. 14 or 15 or 16 or 17 or 18 21. 19 and 20 22. limit 21 to animals 23. 21 not 22
The Cochrane Library	1946 to February 07, 2017	1. pasta/ 2. spaghetti/ 3. macaroni/ 4. lasagna/ 5. fusilli/ 6. noodle/ 7. glycemic index/ 8. glycaemic ind*.tw. 9. glycemic ind*.tw. 10. glycemic load*.tw. 11. glycaemic load*.tw. 12. exp body weight/ 13. body weight*.tw. 14. BMI.tw. 15. body mass index/ 16. body mass index.tw. 17. weight*.tw. 18. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 19. 12 or 13 or 14 or 15 or 16 or 17 20. 18 and 19 21. limit 20 to animals 22. 20 not 21

Supplemental Table S3: PICO framework of the search strategy

PICO framework ^a defined in the present systematic review and meta-analysis			
Participants	Interventions	Comparators	Outcomes
Adult men and women excluding pregnant or breastfeeding women	Low glycemic index interventions where pasta is included as part of the intervention	Higher glycemic index diets where pasta is not included as part of the intervention	Body weight Body mass index (BMI) Body Fat (%) Waist circumference Waist-to-hip ratio

a Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA and PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 2015; 4:1. <https://doi.org/10.1186/2046-4053-4-1>

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4a: Trial characteristics

Overweight/Obese Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)§§	Dietary Prescription	Funding Source
Karl et al. 2015 †††	39 (19M:20F)**	OB, OP					2.33	USA	P	Negative+ Neutral	17		CR to 2/3kcal; Metabolic	Agency
Low GI	20		56 (5)*	92.9 (13.6)*	32.3 (3.4)*	42:133						68:15:16		
Higher GI	19		56 (5)*	94 (9.7)*	33.4 (2.6)*	61:201						70:16:14		
Pereira et al. 2015	19 (4M:15F)**	OW, IP/OP					NR	Brazil	P	Neutral	6.4		Ad libitum	Unknown
Low GI	10		28(5)	80.0(12.6)	29.9 (2.1)	41.2(2.2) ¥ **						48.3:16.1:32.8		
Higher GI	9		26(3)	79.1(12.2)	29.1 (2.0)	74.1(2.9) ¥ **						54.6:12.7:34.4		
Buscemi et al. 2013	40 (19M:21F)**	OW/OB, high CVD risk, OP					NR	Italy	P	Negative	12		CR to 20kcal/kg/d; Ad libitum	Unknown
Low GI	19		51 (8)	93.8 (17.3)	34.3 (6.6)	48.1: 138						56:18:26		
Higher GI	21		49 (8)	93.2 (14.4)	34.5 (5.1)	59.3: 174						57:16:27		
Costa et al. 2012	17 (7M:10F)	OW, IP/OP	25.4 (5.8)	84.1(16.3)	26.3(3.2)		NR	Brazil	C	Neutral	4		Ab libitum,2 meals+3 fruit/d provided	NR
Low GI						47.5(3.8)						58.6:13.9:25.5		
Higher GI						61.6(2.8)						55.4:14.2:30.3		
Jebb et al. 2010 - High MUFA	225**	OH, some CV risk factors, OP	~52		~28.5 ⁺		1.75	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	115			83.7 (69.6-93.1)¶¶¶		~55.2						44.6:16.4:35.7****		
Higher GI	110			80.5 (70.0-92.1)¶¶¶		~63.3						44.9:15.3:35.6****		
Jebb et al. 2010 - Low Fat	250**	OH, some CV risk factors, OP	~52		~28.5 ⁺		3.5	UK	P	Neutral	24		Ad Libitum, key foods provided	Agency, foods by industry
Low GI	117			79.4 (70.1- 91.8)¶¶¶		~56.3						41.5:14.2:26.1****		
Higher GI	111			80.7 (71.4- 91.4)¶¶¶		~64.4						41.1:15.7:27.5****		
Larsen et al. 2010-High Pro	231**	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	124		42.1 (6.5)	88.5 (15.6)		~56.5: 108.9						~43:22:32		
Higher GI	107		42 (5.7)	89.5 (17.1)		~61.4: 113.1						~45:23:31		
Larsen et al. 2010 - Low Pro	203 **	OW/OB, OP			NR		NR	Europe	P	Neutral	26		Ad libitum	Agency
Low GI	106		42.2 (5.7)	88.4 (15.7)		~56.2: 121						~51:18:30		
Higher GI	97		42 (5.9)	86.6 (13.8)		~61.6: 137.9						~51:17:31		

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool.
 Copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4b: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Body composition % (SD)\$§	Dietary Prescription	Funding Source
Solomon et al. 2010	22 (8M:14F)**	OB, Pre-T2DM, OP					7	USA	P	Neutral	12		Metabolic plus exercise program	Agency
Low GI	10 (3M:7F)		67 (6)	97.4 (12.0)	34.9 (1.1)	39.8 (0.9)						33.3(3.3):28.3(0.3):17.0(0.3)		
Higher GI	12 (5M:7F)		64 (3)	94.7 (15.2)	34.1 (1.1)	80.0 (2.1)						30.7(0.7):27.8(0.7):16.6(0.3)		
Philippou et al. 2009- 6 mo	38**	≥1 CHD risk factors, OP	(35-65) ¶				NR	UK	P	Negative (for all but n=2)	24		500kcal CR; Ad libitum	Agency
Low GI	22			91.3(14.8)***	29.1 (3.6)***	50.6 (4.6):114.4(31.5)								
Higher GI	16			97.5(16.4)***	30.5 (3.5)***	63.2(5.6):175.0(45.6)								
Philippou et al. 2009- 4 mo	42**	OW, OP	(18-65)¶				NR	UK	P	Neutral	16		Ad libitum	Unknown
Low GI	23			87.2 (15.3)	32.5 (4.8)	49.7(5.7):89.7(27.5)						34.7(4.7):19.5(4.2):31.8(5.8)		
Higher GI	19			83.6 (13.4)	31.3 (4.8)	63.7(9.4):136.8(56.3)						31.1(4.7):19.3(4.9):30.9(9)		
Abete et al. 2008	32 (18M:14F)	OB, OP	36(7)				2.33	Spain	P	Negative	8		30% CR; Ad libitum, 3-day menus	Agency
Low GI	16			94.3(16.1)	32.8 (4.3)	(40-45)¶						34.8(4.8):18.3(1.6):31.5(1.6)		
Higher GI	16			94.4(13.1)	32.2 (4.4)	(60-65)¶						33.8(3.8):19.6(5.6):32.6(4.3)		
Aston et al. 2008	19 (0M:19F)**	OW/OB, OP	51.9(7.6)	87.5 (15)	33.1 (4.9)		3.33	UK	C	Neutral	12		Ad libitum, key CHO foods provided	Agency
Low GI						55.5(3.8):133.8(22.9)*****						34.0(4.0):17.0(2.4):32.2(5.1)*		
Higher GI						63.9(3):138.8(30.5)*****						36.5(5.1):17.6(3.3):34.1(5.7)*		
Jensen et al. 2008	44 (0M:44F)**	OW, OP	(20-40)¶				3	Denmark	P	Neutral	10		Ad libitum, partial provision, menu plans	Agency, Industry
Low GI	22 (0M:22F)			77.9(6.9)	27.4 (1.5)	72¥						37.7(5.7):17(0):23(5) ±		
Higher GI	22 (0M:22F)			80.2(1.4)	27.6 (0.3)	95¥						37.7(5.7):17(0):22(5) ±		
Philippou et al. 2008	13 (5M:8F)**	≥1 CHD risk factors, OP					NR	UK	P	Negative	12		500kcal CR; Ad libitum	Agency
Low GI	7 (3M:4F)		54 (49-58)¶¶	81.5 (4.7)***	28.7 (2.1)***	51.3(51.0-52.0):105.6 (76.9-110.1)¶¶						36.6(37.8-51.0): 17.1(15.7-22.4): 32.8(31.3-37.1)¶¶		
Higher GI	6 (2M:4F)		45 (39-50)¶¶	89.7 (12.8)***	31.5 (4.4)***	59.3(59.2-64.0):114.7(98.5-134.9)¶¶						49.4(47.8-51.7):19.6(14.0-22.4):29.2(25.2-34.5)¶¶		

Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department of GEZ-LTA Erasmus Hogeschool .
For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Supplement Table S4c: Trial characteristics continued

Overweight/Obese Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance\$	Duration (wks)	Composition % (SD)\$§	Dietary Prescription	Funding Source
Bellisle et al. 2007	65 (OM:65F)**	OW/OB, OP					NR	France	P	Neutral	12		Ad libitum	Industry
Low GI	35		46.1 (13.6)	80 (13.2)	30.2 (4.1)	na								
Higher GI	30		45.3 (12.0)	79 (13.1)	30.4 (4.4)	na								
de Rougemont et al. 2007	38 (22M:16F)**	OW, OP					2.8	France	P	Neutral	5		Ad libitum, some foods provided	Agency, Industry
Low GI	19		36.3 (8.7)	77.2 (9.6)	27.5 (1.3)	46.5 (1.3)						6 (6):19.8 (1.7):37.7 (4.4)		
Higher GI	19		40.4 (9.6)	77.3 (9.2)	27.2 (1.3)	66.3 (2.6)						6 (6):19.8 (1.7):38.4 (3.1)		
Sichieri et al. 2007	123 (OM:123F)**	OW, OP					NR	Brazil	P	Negative	72		100-300kcal CR; 6-d menu and exchange lists provided	Agency
Low GI	63		37.2 (5.4)*	67.7 (6.6)*	na	21(38): 74(84)						6 (6.3): 13.3: 27.2(4.6)		
Higher GI	60		37.5 (5.6)*	68.5 (7.5)*	na	51(28): 199(43)						6 (6.2): 12.3: 26.1(4.7)		
McMillian-Price et al. 2006-High Carb	64(16M:48F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	32		30.5 (7.9)	87.1 (15.3)	30.6 (4.5)	45 (6):89 (28)						6 (6):19 (0):22 (6)		
Higher GI	32		31.8 (9.6)	86 (10.7)	30.9 (3.4)	70 (6):129 (45)						6 (6):18 (6):19 (6)		
McMillian-Price et al. 2006-High Protein	65(15M:50F)	OW/OB, OP					NR	Australia	P	Negative	12		Ad libitum, key foods and meals provided	Agency-Industry
Low GI	33		34.6 (8.6)	88.4 (17.2)	32.1 (5.2)	44 (6):59 (23)						6 (11):26 (6):28 (6)		
Higher GI	32		30.2 (8.5)	87.7 (16.4)	31.3 (4.5)	59 (6):75 (17)						6 (6):28 (6):27 (6)		
Wolever et al. 2002	24 (5M:19F)**	IGT, OP					NR	Canada	P	Neutral	16		Ad libitum, partial provision	Agency
Low GI	13(3M:10F)		55.2 (10.8)	79.7 (13.1)***	29.7 (4.3)	54.4 (2.5):91.8 (9.4)						6 (6.1):19.4 (1.8):24.7 (5.8)		
Higher GI	11(2M:9F)		58.8 (13.3)	76.4 (20.4)***	29.3 (7.3)	59.3 (2.0):96.8 (11.6)						6 (6.6):17.4 (2.3):27.9 (6.3)		

Supplement Table S4d: Trial characteristics continued

Diabetes Trials														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶ ¶ ¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance§	Duration (wks)	Composition % (SD)§§	Dietary Prescription	Funding Source
Jenkins et al. 2014	141(77M:64F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum, bread supplement	Industry Association
Low GI	70 (38M:32F)		59 (10)	85 (20)	30 (5)	~51:53						~38.5:19.8:37.2		
Higher GI	71 (39M:32F)		59 (10)	84 (19)	31 (6)	~62:89						~49.2:19.8:27.4		
Visek et al. 2014	20 (12M:8F)	T2DM, OP	62.7 (5.8)	91.9 (14.1)	32 (4.2)		NR	Czech Republic	C	Neutral	12		Ad libitum	Agency
Low GI						49 (48-51)¶¶						~37.2:18.0:36.0		
Higher GI						68 (61-72)¶¶						~36.2:17.3:40.0		
Jenkins et al. 2012	121 (61M: 60F)	T2DM, OP					NR	Canada	P	Neutral	12		Ad libitum	Agency
Low GI	60		58 (10.1)	85.6 (20.1)	31.4 (7.0)	47: 80						45.4:22.8:30.5		
Higher GI	61		61 (7.8)	82.5 (17.2)	29.9 (5.5)	58: 100						48.3:21.4:28.5		
Yusof et al. 2009	100**	T2DM, OP	NR				NR	Malaysia	P	Neutral	12		Ad libitum, key foods provided to lowGI group	Agency
Low GI	51			69.12 (13.33)	27.05 (4.91)	57(6): 108(32)						52(4):18(3):30(4)		
Higher GI	49			66.83 (11.50)	26.79 (4.65)	64(5): 131(30)						54(4):17(3):28(5)		
Jenkins et al 2008	210 (125M:82F)	T2DM, OP					NR	Canada	P	Neutral	24		Ad libitum	Agency
Low GI	106 (65M:41F)		60 (10)	87.0 (20.0)	30.6 (6.0)	49.4: 91.5						44.0:21.2:33.3		
Higher GI	104 (63M:41F)		61 (9)	87.8 (19.4)	31.2 (5.8)	59.3: 117.9						47.5:20.7:30.5		
Wolever et al. 2008	103	T2DM, OW/OB, OP					NR	Canada	P	Neutral	52		Ad libitum, key foods provided	Agency
Low GI	55		60.6 (7.5)*	81.1 (18.7)*	31.6 (4.5)*	55.1 (3.0): 133 (14.8)						46.7(3.7):20.6(3.0):26.5 (5.9)		
Higher GI	48		60.4 (7.9)*	84.4(18.0)*	30.1 (4.3)*	63.2 (2.8): 135 (20.8)						46.9(3.2):20.4 (2.8):30.8 (4.8)		

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplement Table S4e: Trial characteristics continued

Diabetes Trials continued														
Study	Subjects	Sample Description	Mean Age (y) (SD)	Mean Baseline Body Weight (kg) (SD)	Mean Baseline BMI (kg/m ²) (SD)	Mean GI:GL of diet (SD) ¶¶¶	Pasta Dose (serv/wk)	Setting	Design	Energy Balance§	Duration (wks)	Dietary Composition % (SD)§§	Dietary Prescription	Funding Source
Jimenez-Cruz et al. 2003	14 (6M:8F)**	T2DM, OP	59 (34)					NR	Mexico	C	Neutral			
Low GI				91.6 (24.3)	32.4 (6.0)	44(3.4): 86(19.8)						60:21:23	Ad libitum	Industry
Higher GI				92.6 (25.4)	32.3 (6.0)	56(4.9): 139(27.3)						64:18:20		
Heilbronn et al. 2002	45 (23M:22F)**	T2DM, OW, OP			NR			3.5	Australia	P	Negative		CR to 1500kcal/d; Ad libitum, partial provision	Unknown
Low GI	24 (11M:13F)		56.0(9.4)	91.7(16.2)		43						59(9):22.2 (1.5):17.9 (3.9)		
Higher GI	21 (12M:9F)		57.5(9.6)	93.2 (13.3)		75						60(12.3):21.7 (0.9):17.1 (2.3)		
Fontvieille et al. 1992	18 (12M:6F)	T1DM/T2DM, OP	47.2(11.6)	NR	24.8(2.6)			4.7	France	C	Neutral		Ad libitum	Agency, Industry
Low GI						38.1(5.3)						59(2.2):18.0(2.5):36.2(6.8)		
Higher GI						64.2(3.1)						77(3.3):18.8(1.6):36.3(6.0)		
Fontvieille et al. 1988	8 (4M:4F)	T1DM, OP	43.5 (9.9)	NR	24.1 (6.8)			3.5	France	C	Neutral		Ad libitum, partial provision	Agency, Industry
Low GI						46.5(2.5)						59(1.5):17.4 (1.4):35.0 (2.8)		
Higher GI						60.1 (5.1)						77(2.5):16.9 (1.7):36.0 (2.8)		
CHD Trial														
Frost et al. 2004	55 (48M:7F)**	CHD, OP						NR	UK	P	Neutral §§§		Ad Libitum	Unknown
Low GI	26 (23M:3F)		63.6 (9.4)	81.2 (12.2)	26.9 (3.3)	50(4):115(39)						59 (5):18 (5):31 (5)		
Higher GI	29 (25M:4F)		61.8 (9)	81.7 (16.7)	28.7 (4.6)	57(4):106(34)						77 (10):18 (5):32 (10)		

Abbreviations: BMI=Body Mass Index; C=Crossover design; Carb=Carbohydrate; CED= carbohydrate exchange diet; CHD= Coronary Heart Disease; CR= calorie restriction; CV=cardiovascular; GI= glycemic index; GL=glycemic load; IGT= impaired glucose tolerance; IP=inpatient; IR= insulin resistant; High Carb= high carbohydrate; HI= hyperinsulinemic; M = male; mo=months; MUFA=monounsaturated fatty acids; na=not available; T1DM=type 1 diabetes mellitus; T2DM = type 2 diabetes mellitus; OB= obese; OH=otherwise healthy; OP = outpatient; OW= overweight; P=Parallel design; SD= standard deviation; F = female; UK= United Kingdom; USA= United States of America; wks=weeks; y= years

§Negative energy balance diets were designed for weight loss, Neutral energy balance diets for weight maintenance where all diets are isocaloric between test and control groups; §§Energy from carbohydrate: protein: fat for the planned diet or if not planned; §§§ Participants were advised to lose weight if they had a BMI>28kg/m².

¶ Range of values; ¶¶ median and interquartile range (IQR); ¶¶¶ Actual GI/GL or if not available, planned;

* Calculated before dropout; **Completer Analysis, as used in data analysis; *** completer analysis, data obtained from authors; **** based on less participants than in analysis;

+ approximate based on all study arms; ¥ approximate based on test meals; ‡ approximate based on n=45 from Sirtori et al. 2004, the original publication of this study;

‡‡‡ analysis includes weight loss and weight maintenance phases

Supplemental Table S5: Sensitivity analysis of the systematic removal of each trial*

	MD [95% CI], P-value I ² , P-value					
	Body Weight (kg) N=32	BMI (kg/m ²) N=18	Body Fat (%) N=10	Waist Circumference (cm) N=18	Waist-to-hip Ratio N=6	Sagittal Abdominal Diameter (cm) N=3
All studies	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.01 [-0.58, 0.56], P=0.98 65%, P=0.003	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92
Removal of:						
OW/OB						
Pereira et al. 2015	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.54	-0.24 [-0.34, -0.14], P<0.01 0.00%, P=0.93	0.19 [-0.25, 0.63], P=0.39 36%, P=0.13	-0.37 [-1.04, 0.31], P=0.28 62%, P<0.01	0.00 [-0.01, 0.01], P=0.67 0.00%, P=0.58	n/a
Karl et al. 2015	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.50	-0.25 [-0.35, -0.16], P<0.01 0.00%, P=0.91	0.10 [-0.51, 0.71], P=0.74 64%, P<0.01	n/a	n/a	n/a
Buscemi et al. 2013	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.86	-0.08 [-0.63, 0.48], P=0.79 64%, P<0.01	-0.46 [-1.06, 0.15], P=0.14 64%, P<0.01	n/a	n/a
Costa et al. 2012	-0.62 [-0.84, -0.41], P<0.01 0.00%, P=0.46	-0.26 [-0.37, -0.16], P<0.01 0.00%, P=0.86	-0.06 [-0.73, 0.61], P=0.86 66%, P<0.01	-0.33 [-0.94, 0.28], P=0.29 59%, P<0.01	n/a	n/a
Jebb et al. 2010 - LowFat	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - LowPro	-0.60 [-0.81, -0.38], P<0.01 0.00%, P=0.57	n/a	n/a	-0.45 [-1.09, 0.18], P=0.16 64%, P<0.01	n/a	-0.05 [-0.39, 0.29], P=0.79 0.00%, P=0.86
Jebb et al. 2010- HighMUFA	-0.64 [-0.85, -0.43], P<0.01 0.00%, P=0.65	n/a	n/a	n/a	n/a	n/a
Larsen et al. 2010 - HighPro	-0.62 [-0.83, -0.41], P<0.01 0.00%, P=0.48	n/a	n/a	-0.37 [-1.00, 0.26], P=0.26 62%, P<0.01	n/a	-0.13 [-0.45, 0.19], P=0.42 0.00%, P=0.92
Solomon et al. 2010	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	n/a	n/a	n/a	n/a

	0.00%, P=0.47	0.00%, P=0.88				
Philippou et al. 2009 - 6mnth	-0.65 [-0.86, -0.44], P<0.01 0.00%, P=0.59	-0.28 [-0.37, -0.18], P<0.01 0.00%, P=0.97	-0.12 [-0.71, 0.48], P=0.70 65%, P<0.01	-0.55 [-1.15, 0.04], P=0.07 61%, P<0.01	n/a	n/a
Philippou et al. 2009 - 4mnth	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.89	-0.09 [-0.72, 0.54], P=0.78 66%, P<0.01	-0.44 [-1.08, 0.19], P=0.17 64%, P<0.01	n/a	n/a
Abete et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a	n/a
Philippou et al. 2008	-0.62 [-0.83, -0.42], P<0.01 0.00%, P=0.48	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.89	-0.06 [-0.68, 0.55], P=0.84 68%, P<0.01	-0.41 [-1.02, 0.19], P=0.18 63%, P<0.01	n/a	n/a
Aston et al. 2008	-0.66 [-0.87, -0.44], P<0.01 0.00%, P=0.52	n/a	-0.00 [-0.70, 0.69], P=0.99 68%, P<0.01	-0.54 [-1.14, 0.07], P=0.08 62%, P<0.01	n/a	n/a
Jensen et al. 2008	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.27 [-0.36, -0.17], P<0.01 0.00%, P=0.88	n/a	-0.44 [-1.05, 0.18], P=0.16 64%, P<0.01	0.00 [-0.01, 0.00], P=0.24 0.00%, P=0.41	-0.09 [-0.35, 0.18], P=0.51 0.00%, P=0.69
de Rougemont et al. 2007	-0.57 [-0.80, -0.34], P<0.01 0.00%, P=0.53	-0.25 [-0.36, -0.14], P<0.01 0.00%, P=0.87	0.06 [-0.57, 0.70], P=0.84 67%, P<0.01	n/a	n/a	n/a
Sichieri et al. 2007	-0.66 [-0.88, -0.45], P<0.01 0.00%, P=0.53	n/a	n/a	n/a	n/a	n/a
Bellisle et al. 2007	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.88	n/a	-0.47 [-1.08, 0.13], P=0.13 64%, P<0.01	0.00 [-0.01, 0.00], P=0.14 0.00%, P=0.70	n/a
McMillan-Price et al. 2006 - HighCHO	-0.61 [-0.82, -0.39], P<0.01 0.00%, P=0.51	n/a	n/a	-0.38 [-1.01, 0.25], P=0.23 63%, P<0.01	n/a	n/a
McMillan-Price et al. 2006 - HighPro	-0.70 [-0.91, -0.49], P<0.01 0.00%, P=0.91	n/a	n/a	-0.62 [-1.19, -0.05], P=0.03 55%, P<0.01	n/a	n/a
Wolever et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a	n/a
Diabetes:						
Vissek et al. 2014	-0.63 [-0.84, -0.42], P<0.01	-0.26 [-0.36, -0.16], P<0.01	0.01 [-0.57, 0.60], P=0.96	n/a	n/a	n/a

	0.00%, P=0.46	0.00%, P=0.86	68%, P<0.01		
Jenkins et al. 2014	-0.66 [-0.88, -0.43], P<0.01 0.00%, P=0.47	-0.29 [-0.39, -0.18], P<0.01 0.00%, P=0.94	n/a	-0.61 [-1.18, -0.04], P=0.04 50%, P=0.01	0.00 [-0.01, 0.00], P=0.21 0.00%, P=0.43
Jenkins et al. 2012	-0.62 [-0.84, -0.40], P<0.01 0.00%, P=0.46	-0.25 [-0.35, -0.15], P<0.01 0.00%, P=0.87	n/a	-0.44 [-1.05, 0.17], P=0.16 64%, P<0.01	0.00 [-0.01, 0.01], P=0.64 0.00%, P=0.53
Yusof et al. 2009	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.87	n/a	-0.33 [-0.95, 0.28], P=0.29 58%, P<0.01	n/a
Jenkins et al. 2008	-0.61 [-0.83, -0.40], P<0.01 0.00%, P=0.48	-0.25 [-0.36, -0.15], P<0.01 0.00%, P=0.87	n/a	n/a	n/a
Wolever et al. 2008	-0.64 [-0.84, -0.43], P<0.01 0.00%, P=0.57	n/a	n/a	-0.50 [-1.10, 0.09], P=0.10 62%, P<0.01	n/a
Jimenez-Cruz et al. 2003	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	-0.26 [-0.35, -0.16], P<0.01 0.00%, P=0.86	n/a	n/a	n/a
Heilbronn et al. 2002	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	n/a	n/a	n/a	n/a
Fontvieille et al. 1992	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a
Fontvielle et al. 1988	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.46	n/a	n/a	n/a	n/a
CHD					
Frost et al. 2004	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.47	-0.26 [-0.36, -0.17], P<0.01 0.00%, P=0.89	n/a	-0.48 [-1.09, 0.12], P=0.12 63%, P<0.01	0.00 [-0.01, 0.00], P=0.25 1%, P=0.40

*Sensitivity analysis included the removal of each single study from the meta-analyses one at a time and the summary effect was recalculated. An influential outlier was considered a study whose removal changed the magnitude of the pooled effect by >10%.

BMI, body mass index; CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; MD, mean difference; mnt, month; MUFA, monounsaturated fatty acids; n/a, not applicable; OB, obese; OW, overweight; Pro, protein

Supplemental Table S6: Sensitivity analyses of the use of correlation coefficients of 0.25 and 0.75 in crossover trials

Outcome (no. crossover trials/total)	MD (95% CI), P-value I ² , P-value		
	Correlation Coefficient used in the Primary Analysis	Correlation Coefficient used in Sensitivity Analyses	
	0.5	0.25	0.75
Body Weight, kg (6*/32)	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.42], P<0.01 0.00%, P=0.51	-0.63 [-0.84, -0.43], P<0.01 0.00%, P=0.51
BMI, kg/m ² (3/18)	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.91	-0.26 [-0.36, -0.16], P<0.01 0.00%, P=0.90	-0.26 [-0.35, -0.17], P<0.01 0.00%, P=0.90
Body Fat, % (3*/10)	-0.01 [-0.58, 0.56], P=0.98 65%, P<0.01	-0.00 [-0.58, 0.58], P=0.99 64%, P<0.01	-0.02 [-0.57, 0.54], P=0.96 66%, P<0.01
Waist Circumference, cm (2*/18)	-0.46 [-1.05, 0.14], P=0.13 62%, P<0.01	-0.44 [-1.04, 0.15], P=0.14 60%, P<0.01	-0.46 [-1.07, 0.14], P=0.13 65%, P<0.01
Waist-to-hip Ratio (0/6)	-0.00 [-0.01, 0.00], P=0.27 0.00%, P=0.52	n/a	n/a
Sagittal Abdominal Diameter, cm (0/6)	-0.09 [-0.34, 0.16], P=0.48 0.00%, P=0.92	n/a	n/a

* one of these crossover trials, however, did not require the use of a correlation coefficient as complete data was available

BMI, body mass index; CI, confidence interval; MD, mean difference; no., number

Supplement Table S7. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	24.1 – 37.1 kg/m ²	32	2448	-0.04 (-0.13, 0.06)	0.00%	0.454
Follow-up	3 – 72 wks	32	2448	0.01 (-0.01, 0.02)	0.00%	0.314
Dose Pasta	1.75 – 7.0 serv/wk	11	740	0.29 (-0.91, 1.50)	0.00%	0.595
GI ³	21 - 72	31	2383	-0.01 (-0.04, 0.02)	0.45%	0.482
Difference in GI ²	4.9 - 40.2	31	2383	-0.00 (-0.04, 0.03)	2.75%	0.811
Fiber ³	8.0 - 39.4g/d	27	1851	0.02 (-0.01, 0.06)	0.00%	0.221
Change in Fiber ⁴	-7.5 - +13.8g/d	17	1571	-0.02 (-0.07, 0.04)	23.13%	0.595
Difference in Fiber ²	-8.5 - +15.3g/d	27	1851	-0.01 (-0.06, 0.04)	0.00%	0.701
Saturated Fat ³	5.1 - 12.5%	14	1396	-0.05 (-0.30, 0.20)	34.32%	0.648
Change in Saturated Fat ⁴	-8.2 - -1.2%	12	1309	-0.01 (-0.29, 0.28)	46.27%	0.965
Difference in Saturated Fat ²	-2.0 - +2.3%	14	1396	-0.34 (-1.08, 0.39)	32.80%	0.329
CHO ³	37.2 - 68.0%	30	2345	-0.01 (-0.05, 0.03)	0.00%	0.541
Change in CHO ⁴	-7.2 - +10.1%	19	2046	-0.07 (-0.12, -0.01)	0.00%	0.016
Difference in CHO ²	-11.1 - +5.4%	30	2345	0.02 (-0.05, 0.10)	0.00%	0.547
Protein ³	13.3 - 26.1%	30	2345	0.04 (-0.04, 0.12)	0.00%	0.332

Change in Protein ⁴	-0.5 - +8.6%	19	2046	0.15 (0.03, 0.27)	0.00%	0.017
Difference in Protein ²	-2.5 - +3.4%	30	2345	-0.18 (-0.38, 0.02)	0.00%	0.069
Fat ³	16.0 - 37.7%	30	2345	0.01 (-0.05, 0.06)	0.00%	0.852
Change in Fat ⁴	-12.2 - +5.4%	19	2046	0.05 (-0.02, 0.12)	17.73%	0.122
Difference in Fat ²	-4.4 - +10.6%	30	2345	0.02 (-0.05, 0.09)	0.00%	0.610

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body weight on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body weight. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplemental Table S8. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 37.1 kg/m ²	18	1038	-0.01 (-0.06, 0.03)	0.00%	0.559
Follow-up	4 – 24 wks	18	1038	0.00 (-0.01, 0.02)	0.00%	0.559
Dose Pasta*	2.33 – 7.0 serv/wk	4	143			
GI ³	39.8 - 72	17	973	0.01 (-0.01, 0.03)	0.00%	0.153
Difference in GI ²	7.0 - 40.2	17	973	-0.01 (-0.03, 0.01)	0.00%	0.275
Fiber ³	8.0 - 39.4g/d	16	935	0.01 (-0.01, 0.02)	0.00%	0.366
Change in Fiber ⁴	-1.8 - +12.4g/d	10	758	0.00 (-0.03, 0.03)	0.00%	0.846
Difference in Fiber ²	-4.9 - +13.0g/d	16	935	-0.00 (-0.02, 0.02)	0.00%	0.821
Saturated Fat* ³	7.6 - 12.5%	7				
Change in Saturated Fat* ⁴	-2.4 - -1.2%	6				
Difference in Saturated Fat* ²	-1.0 - +2.3%	7				
CHO ³	37.2 - 68.0%	16	935	-0.00 (-0.02, 0.01)	0.00%	0.594
Change in CHO ⁴	-5.6 - +3.2%	10	758	-0.02 (-0.06, 0.02)	0.00%	0.325
Difference in CHO ²	-11.1 - +2.0%	16	935	-0.01 (-0.03, 0.02)	0.00%	0.531
Protein ³	13.9 – 22.8%	16	935	0.00 (-0.04, 0.04)	0.00%	0.905

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool. by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Change in Protein ⁴	-0.2 - +3.0%	10	758	0.01 (-0.11, 0.14)	0.00%	0.786
Difference in Protein ²	-2.5 - +3.4%	16	935	-0.03 (-0.12, 0.06)	0.00%	0.441
Fat ³	16.0 - 37.7%	16	935	0.00 (-0.02, 0.02)	0.00%	0.717
Change in Fat ⁴	-4.8 - +5.4%	10	758	0.02 (-0.02, 0.06)	0.00%	0.240
Difference in Fat ²	-4.4 - +10.6%	16	935	0.01 (-0.01, 0.04)	0.00%	0.299

*For Dose, there were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in BMI on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in BMI. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplemental Table S9. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on body fat (%)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	10	285	-0.02 (-0.26, 0.23)	67.69%	0.890
Follow-up	4 – 24 wks	10	285	0.06 (-0.06, 0.17)	66.00%	0.303
Dose Pasta*	2.33 - 3.33 serv/wk	3	96			
GI ³	39.8 - 72	10	285	0.12 (-0.00, 0.25)	52.21%	0.053
Difference in GI ²	7.0 - 40.2	10	285	-0.09 (-0.15, -0.03)	19.39%	0.008
Fiber* ³	8.0 - 39.4g/d	8	228			
Change in Fiber* ⁴	-1.8 - +12.4g/d	4	87			
Difference in Fiber* ²	-4.9 - +13.0g/d	8	228			
Saturated Fat* ³	7.6 - 12.5%	3	93			
Change in Saturated Fat* ⁴	-2.4 - -1.2%	2	51			
Difference in Saturated Fat* ²	-1.0 - +2.3%	3	93			
CHO* ³	37.2 - 68.0%	9	247			
Change in CHO* ⁴	-5.6 - +3.2%	4	87			
Difference in CHO* ²	-11.1 - +2.0%	9	247			
Protein* ³	13.9 – 22.8%	9	247			

-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Change in Protein* ⁴	-0.2 - +3.0%	4	87		
Difference in Protein* ²	-2.5 - +3.4%	9	247		
Fat* ³	16.0 - 37.7%	9	247		
Change in Fat* ⁴	-4.8 - +5.4%	4	87		
Difference in Fat* ²	-4.4 - +10.6%	9	247		

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in body fat on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in body fat. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

Supplemental Table S10. Continuous meta-regression analysis for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)¹

Subgroup	Range	No. trials	N	β (95% CI)	Residual I ²	P value
Baseline BMI	26.3 – 36.0 kg/m ²	18	1380	0.10 (-0.13, 0.32)	59.02%	0.372
Follow-up	4 – 52 wks	18	1380	0.05 (-0.03, 0.13)	60.50%	0.225
Dose Pasta*	3.0 – 3.33 serv/wk	2	63			
GI ³	41.2 – 72.0	17	1315	-0.01 (-0.12, 0.10)	66.00%	0.841
Difference in GI ²	4.9 – 32.9	17	1315	-0.02 (-0.10, 0.06)	64.51%	0.596
Fiber ³	8.0 - 39.4g/d	15	1258	0.05 (-0.05, 0.14)	59.60%	0.342
Change in Fiber ⁴	-1.8 - +13.8g/d	13	1176	0.01 (-0.17, 0.20)	70.59%	0.861
Difference in Fiber ²	-4.9 - +15.3g/d	15	1258	-0.00 (-0.14, 0.13)	64.86%	0.939
Saturated Fat* ³	6.1 – 10.1%	8	604			
Change in Saturated Fat* ⁴	-7.6 - -1.2%	7	562			
Difference in Saturated Fat* ²	-2.0 - +2.3%	8	604			
CHO ³	38.5 - 58.6%	16	1277	-0.11 (-0.19, -0.04)	27.06%	0.007
Change in CHO ⁴	-7.2 - +10.1%	13	1176	-0.09 (-0.23, 0.04)	57.19%	0.148
Difference in CHO ²	-11.1 - +5.4%	16	1277	-0.01 (-0.18, 0.17)	60.81%	0.947
Protein ³	13.9 – 26.1%	16	1277	0.20 (0.01, 0.38)	43.92%	0.038

Change in Protein ⁴	-0.3 - +8.6%	13	1176	0.19 (-0.09, 0.46)	63.38%	0.165
Difference in Protein ²	-2.5 - +3.4%	16	1277	-0.14 (-0.66, 0.38)	62.79%	0.570
Fat ³	22.0 - 37.2%	16	1277	0.10 (-0.05, 0.26)	53.48%	0.185
Change in Fat ⁴	-12.2 - +5.4%	13	1176	0.08 (-0.07, 0.24)	61.15%	0.274
Difference in Fat ²	-4.4 - +10.6%	16	1277	0.09 (-0.06, 0.25)	47.17%	0.220

*There were <10 trials so subgroup analyses were not performed.

1 Data is presented as between group mean difference (95% CI) for a 1-unit change in the predictor variable. β -coefficients were estimated using continuous meta-regression analysis. A positive β -coefficient implies an increase in waist circumference on the pasta/low-GI intervention as the subgroup variable increases, and a negative β -coefficient implies a decrease in waist circumference. Residual I^2 reports inter-study heterogeneity not explained by the subgroup and was estimated using the Cochran's Q statistic.

2 Difference in diet variable between the intervention and control arms

3 Intake at the end of study in the intervention arm

4 Change in intake from end of study from baseline in intervention arm

BMI, body mass index; CHO, carbohydrate; GI, glycemic index; wk, week

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Supplementary Table S11. Post-hoc piecewise linear continuous dose-response meta-regression analyses for the effect of pasta intake in the context of low-GI dietary patterns on body weight (kg)

Dose threshold of servings of pasta consumed per week	Dose ranges of servings of pasta consumed per week	β (95% CIs) *	Residual I^2 †	p -value
3.0	≤ 3.0	-0.70 (-3.27, 1.86)	0.00%	0.890
	> 3.0	0.91 (-0.89, 2.70)		
3.33	≤ 3.33	0.05 (-1.80, 1.89)	0.00%	0.518
	> 3.33	0.44 (-1.75, 2.63)		
3.5	≤ 3.5	0.09 (-1.65, 1.82)	0.00%	0.888
	> 3.5	0.46 (-1.89, 2.81)		

* β is the slope derived from the piecewise linear meta-regression analyses and represents the treatment effect on body weight for doses above and below each dose-threshold representing servings of pasta consumption per week. † The residual I^2 value indicates heterogeneity unexplained by each dose-threshold.

-2017-019438 on 24 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Supplementary Table S12: GRADE assessment of study quality

Quality assessment*							№ of patients		Effect	Quality Importance	
№ of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)		
Body Weight (follow up: median 12 weeks)										April 2018. Downloaded from https://bmjopen.bmj.com/ on June 5, 2025 at Department of Health Services, University of California, San Francisco. All rights reserved. No reuse allowed without permission. All training, and similar technologies.	
32	randomised trials	not serious	not serious	serious ^a	not serious	none	1294	1228	MD -0.63 kg (-0.84 to -0.42)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
BMI (follow up: median 12 weeks)											
18	randomised trials	not serious	not serious	serious ^a	not serious	none	551	538	MD -0.26 kg/m² (-0.36 to -0.16)		⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Waist Circumference (follow up: median 12 weeks)											

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Quality assessment*							Nº of patients		Effect	Quality Importance
Nº of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)	
18	randomised trials	not serious	serious ^b	serious ^a	not serious	none	717	624	MD -0.46 cm (0.05 to 0.14)	⊕⊕○○ LOW ^{a,b} Due to downgrade for inconsistency and indirectness
Body Fat (follow up: median 12 weeks)										
10	randomised trials	not serious	serious ^c	serious ^a	not serious	none	176	165	MD -0.01 % (0.58 to 0.56)	⊕⊕○○ LOW ^{a,c} Due to downgrade for inconsistency and indirectness
Waist-to-hip Ratio (follow up: median 12 weeks)										

Quality assessment*							N ^o of patients		Effect	Quality Importance
N ^o of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Pasta/LGI diet	Higher GI diet	Absolute (95% CI)	
6	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	223	222	MD -0.00 (0.01 to 0.00)	⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness
Sagittal Abdominal Diameter (follow up: median 26 weeks)										
3	randomised trials	not serious	not serious	serious ^a	not serious	none ^d	237	214	MD -0.09 cm (0.34 to 0.16)	⊕⊕⊕○ MODERATE ^a Due to downgrade for indirectness

CI, Confidence interval; GI, glycemic index; LGI, low glycemic index; MD, Mean difference

*All outcomes started with high quality evidence since all studies were randomized controlled trials. Risk of Bias – We rated down for risk of bias if the majority of studies were considered to be at high risk of bias. Inconsistency – We assessed inconsistency using I^2 estimates where an I^2 of 50% or higher indicates substantial heterogeneity. I^2 is the percentage of variability in the treatment estimates that is attributable to heterogeneity between studies. We rated down for inconsistency if there was substantial heterogeneity that was unexplained by any a priori sensitivity or subgroup analyses. Indirectness – We rated down for indirectness if there were factors present relating to the population, interventions, and outcomes that limited the generalizability of the results. Imprecision – We rated down for imprecision if the 95% confidence interval (95% CI) crossed the minimally important difference (MID) for harm. MIDs used for each outcome are: 0.5kg for body weight (based on Johnston et al. JAMA 2014;312(9):923-33); 0.2kg/m² for BMI; 2.0cm for waist circumference; 2.0% for body fat; 0.02 for waist-to-hip ratio; 2.0cm for sagittal abdominal diameter.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

- a. Downgrade for serious indirectness, since most of the available trials did not quantify the amount of fibre consumed in the context of the low-GI dietary patterns
- b. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2=62\%$, P -heterogeneity<0.001), which could not be explained
- c. Downgrade for serious inconsistency, as there was evidence of substantial inter-study ($I^2= 65\%$, P -heterogeneity=0.003), which could not be explained
- d. No downgrade for publication bias, as publication bias could not be assessed due to lack of power for assessing funnel plot asymmetry and small study effects (<10 trials included in the meta-analysis)

2017-019438 on 27 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml


Supplemental Figures

	Random Sequence Generation	Allocation Concealment	Blinding of Participants, Personnel and Outcome Assessment	Incomplete Outcome Data	Selective Reporting
Abete et al. 2008	?	?	?	+	+
Aston et al. 2008	+	?	+	+	+
Bellisle et al. 2007	?	?	?	?	+
Buscemi et al. 2012	+	+	+	?	+
Cost et al. 2012	?	?	+	+	+
de Rougemont et al. 2007	?	+	+	+	+
Fontvielle et al. 1992	?	?	+	+	+
Fontvielle et al. 1988	?	?	+	+	+
Frost et al. 2004	?	?	+	+	+
Heilbronn et al. 2002	?	?	+	?	+
Jebb et al. 2010 - HighMUFA	+	?	+	?	+
Jebb et al. 2010 - LowFat	+	?	+	?	+
Jenkins et al. 2014	+	+	+	+	+
Jenkins et al. 2012	+	+	+	+	+
Jenkins et al. 2008	+	+	+	+	+
Jensen et al. 2008	?	?	+	+	+

Jimenez-Cruz et al. 2003	?	?	?	?	+
Karl et al. 2015	+	+	+	+	+
Larsen et al. 2010 -LowPro	+	+	+	+	+
Larsen et al. 2010 -HighPro	+	+	+	+	+
McMillan-Price et al. 2006- HighCHO	?	?	+	+	+
McMillan-Price et al. 2006- HighPro	?	?	+	+	+
Pereira et al. 2015	?	?	+	?	+
Philippou et al. 2008	?	?	+	?	+
Philippou et al. 2009-4mo	?	?	+	?	+
Philippou et al. 2009-6mo	?	?	?	?	+
Sicheri et al. 2007	+	?	+	?	+
Solomon et al. 2010	?	?	+	+	+
Visek et al. 2014	?	?	+	+	+
Wolever et al. 2008	+	+	?	+	+
Wolever et al. 2002	+	+	+	+	+
Yusof et al. 2009	+	?	+	+	+


Supplemental Figure S1: Cochrane risk of bias summary for all included trials

Summary of risk of bias ratings for each individual study included in the meta-analysis.



Supplemental Figure S2: Risk of bias proportion graph for all included trials


-2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S3: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on BMI (kg/m²) (n= 1038)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


BMI, body mass index; CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S4: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n= 285)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of P<0.10 and quantified with the use of the I² statistic where I²≥50% is considered to be evidence of substantial heterogeneity and ≥75% considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S5: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S6: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on waist-to-hip ratio (n = 445)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot’s bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran’s Q statistic at a significance level of $P<0.10$ and quantified with the use of the I^2 statistic where $I^2\geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet



Supplemental Figure S7: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on sagittal abdominal diameter (cm) (n = 478)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.


CHD, coronary heart disease; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; Pro, protein



Supplemental Figure S8: Forest plot of randomized controlled trials investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) using fixed effects models (n = 2448)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance fixed effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CHD, coronary heart disease; CHO, carbohydrate; CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids; Pro, protein.




Supplemental Figure S9: Forest plot of randomized controlled trials which contain data for approximating pasta intake which are investigating the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 740)

Data are expressed as mean differences represented by a square and 95% CIs by the line through the square. 95% CIs exceeding the plot's bounds are represented by an arrowhead. Pooled effect estimates are represented by diamonds and were estimated with the use of generic inverse variance random effects models. Between-study heterogeneity was detected with the use of the Cochran's Q statistic at a significance level of $P < 0.10$ and quantified with the use of the I^2 statistic where $I^2 \geq 50\%$ is considered to be evidence of substantial heterogeneity and $\geq 75\%$ considerable heterogeneity.

CI, confidence interval; HGI, higher glycemic index diet; GI, glycemic index; LGI, low glycemic index diet; MUFA, monounsaturated fatty acids

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S10: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p = 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.

BMI, body mass index; btw, between; CHD, coronary heart disease; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; serv, serving; trt, treatment; wk, week.


*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.00 kg (-0.08, 0.65) (1 vs. 2) to 1.32 kg (-4.28, 6.91) (1 vs. 3) to -1.29 kg (-6.91, 4.32) (2 vs. 3).

† data available on 31 studies

‡ data available on 17 studies

§ data available on 27 studies

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 5, 2025 at Department GEZ-LTA Erasmus Hogeschool .
by copyright, including for uses related to text and data mining, AI training, and similar technologies.



Supplemental Figure S11: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body weight (kg) continued (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $p < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes.


BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); trt, treatment.

§ data available on 30 studies

¥ data available on 19 studies

¶ data available on 14 studies

Ψ data available on 12 studies




Supplemental Figure S12: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body weight (kg) (n = 2448)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body weight. Within subgroup mean differences are the pooled effect estimates represented by a diamond and 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup.

LRB=Low Risk of Bias, URB= Unclear Risk of Bias, HRB=High Risk of Bias. *Within and between subgroup analysis could not be performed against HRB since no values for HRB subgroup. ** no values for URB subgroup. § Blinding of Participants, Personnel, and Outcome Assessors.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S13: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I² was estimated with the use of the Cochran’s Q statistic and indicates the between study

heterogeneity unexplained by the variability in response because of the specific between studies factor. $I^2 = 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (no trials), therefore analyses were not performed.

*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.00kg/m² (-0.156, 0.274) (1 vs. 2) to 0.584kg/m² (-1.009, 2.178) (1 vs. 3) to -0.525kg/m² (-2.122, 1.072) (2 vs. 3).


† data available on 17 studies

‡ data available on 16 studies

§ data available on 10 studies

BMI, body mass index; btw, between; CI, confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47




Supplemental Figure S14: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) continued (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=7 trials), saturated fat within treatment (n=6 trials) and saturated fat between treatments (n=7 trials), therefore analyses were not performed.

§ data available on 16 studies

¥ data available on 10 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.



Supplemental Figure S15: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body mass index, BMI (kg/m²) (n = 1038)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on BMI. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup.


*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.

BMI, body mass index; CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47



Supplemental Figure S16: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P=0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=6), test carbohydrate, protein and fat (n=9), carbohydrate, protein and fat within treatment (n=4), carbohydrate, protein and fat between treatment (n=9), test fibre (n=8), fibre within treatment (n=4), fibre between treatment (n=8), test saturated fat (n=3 trials), saturated fat within treatment (n=2 trials) and saturated fat between treatments (n=3 trials), therefore analyses were not performed.

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.



Supplemental Figure S17: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on body fat (%) (n = 285)


The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on body fat. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P = 0.05$ indicated that the effect size differed between levels of the subgroup.

*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.

** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.

CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias



Supplemental Figure S18: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for dose (n=2 trials), therefore analyses were not performed.


*Pairwise between-subgroup mean differences (95% CIs) for Metabolic Phenotype were as follows: 0.15cm (-1.492, 1.818) (1 vs. 2) to 1.608cm (-3.432, 6.648) (1 vs. 3) to -1.445cm (-6.630, 3.740) (2 vs. 3).

† data available on 17 studies

‡ data available on 15 studies

BMI, body mass index; btw, between; CI = confidence interval; GI, glycemic index (glucose scale); OB, obese; OW, overweight; trt, treatment.

2017-019438 on 2 April 2018. Downloaded from <http://bmjopen.bmj.com/> on June 9, 2025 at Department GEZ-LTA Erasmus Hogeschool. All rights reserved. No reuse allowed without permission. For copyright, including for uses related to text and data mining, AI training, and similar technologies.




Supplemental Figure S19: Subgroup analysis for mean differences (95% CIs) of the effects of pasta in the context of low-GI dietary patterns on waist circumference (cm) continued (n = 1380)

The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's Q statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup. Within and between-treatment changes were categorically assessed based on the median intakes. There were less than 10 trials available for categorical subgroup assessment for test saturated fat (n=8 trials), saturated fat within treatment (n=7 trials) and saturated fat between treatments (n=8 trials), therefore analyses were not performed.

§ data available on 16 studies

¥ data available on 13 studies

BMI, body mass index; btw, between; CHO, carbohydrate; CI = confidence interval; GI, glycemic index (glucose scale); trt, treatment.



Supplemental Figure S20: A priori subgroup analyses on risk of bias for mean differences (95% CI) of the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm) (n = 1380)


The dashed line represents the pooled estimate for the overall primary analysis of pasta in the context of low-GI dietary patterns on waist circumference. Within subgroup mean differences are the pooled effect estimates represented by a diamond. 95% CIs are represented by the line through the square. The residual I^2 was estimated with the use of the Cochran's statistic and indicates the between study heterogeneity unexplained by the variability in response because of the specific between studies factor. $P < 0.05$ indicated that the effect size differed between levels of the subgroup.

*Within and between subgroup analyses could not be performed against HRB since no values for HRB subgroup.


** no values for URB subgroup.

§ Blinding of Participants, Personnel, and Outcome Assessors.


CI, confidence interval; HRB, High Risk of Bias; GI, glycemic index; LRB, Low Risk of Bias; URB, Unclear Risk of Bias




Supplemental Figure S21: Continuous meta-regression of change in carbohydrate intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)
CHO, carbohydrate; MD, mean difference



Supplemental Figure S22: Continuous meta-regression of change in protein intake in the low-GI dietary pattern intervention arms with change in body weight (n=19)
MD, mean difference




Supplemental Figure S23: Pasta intake dose-response analyses by spline curve modeling (MKSPLINE procedure)




Supplemental Figure S24: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body weight (kg)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.




Supplemental Figure S25: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on BMI (kg/m²)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S26: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on body fat (%)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.



Supplemental Figure S27: Funnel plot for the effect of pasta in the context of low-GI dietary patterns on waist circumference (cm)

The solid line represents the overall pooled estimate for all studies included in the meta-analysis expressed as a weighted mean difference. Dashed lines represent 95% CIs. P values are derived from the quantitative assessment of publication bias by Egger's and Begg's tests.

Supplemental Table S1: PRISMA Checklist



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5-6, Supplemental Table S2
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5, Supplemental Table S1
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplemental Table S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5-6, Figure 1
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and	6, Supplemental

		simplifications made.	Table S2
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6-7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7-8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	8

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7-10
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10-11, Table 1, Supplemental Tables S2-S3
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	11, Supplemental Figures S1-S2
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	11-13, Figures 2-3, Supplemental Figures S3-S7
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	11-13, Figures 2-3, Supplemental Figures S3-S7
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see item 15).	11,15 Supplemental

			Table S1, S2, S12, S15, S17, S20
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	12-16, Supplemental Tables S4-S10, Supplemental Figures S8-S26
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	16-22
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	19-21
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	22-23
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	26-31

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.