













BMJ Open EUROLinkCAT protocol for a European population-based data linkage study investigating the survival, morbidity and education of children with congenital anomalies

Joan K Morris ¹, Ester Garne ², Maria Loane ³, Ingeborg Barisic ⁴, James Densem⁵, Anna Latos-Bieleńska ⁶, Amanda Neville⁷, Anna Pierini ⁸, Judith Rankin ⁹, Anke Rissmann ¹⁰, Hermien de Walle ¹¹, Joachim Tan ¹, Joanne Emma Given ³, Hugh Claridge ¹, EUROLinkCAT Consortium

To cite: Morris JK, Garne E, Loane M, *et al.* EUROLinkCAT protocol for a European population-based data linkage study investigating the survival, morbidity and education of children with congenital anomalies. *BMJ Open* 2021;**11**:e047859. doi:10.1136/bmjopen-2020-047859

► Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-047859>).

Received 14 December 2020
Accepted 07 May 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Professor Joan K Morris;
jmorris@sgul.ac.uk

ABSTRACT

Introduction Congenital anomalies (CAs) are a major cause of infant mortality, childhood morbidity and long-term disability. Over 130 000 children born in Europe every year will have a CA. This paper describes the EUROLinkCAT study, which is investigating the health and educational outcomes of children with CAs for the first 10 years of their lives.

Methods and analysis EUROCAT is a European network of population-based registries for the epidemiological surveillance of CAs. EUROLinkCAT is using the EUROCAT infrastructure to support 22 EUROCAT registries in 14 countries to link their data on births with CAs to mortality, hospital discharge, prescription and educational databases. Once linked, each registry transforms their case data into a common data model (CDM) format and they are then supplied with common STATA syntax scripts to analyse their data. The resulting aggregate tables and analysis results are submitted to a central results repository (CRR) and meta-analyses are performed to summarise the results across all registries. The CRR currently contains data on 155 594 children with a CA followed up to age 10 from a population of 6 million births from 1995 to 2014.

Ethics The CA registries have the required ethics permissions for routine surveillance and transmission of anonymised data to the EUROCAT central database. Each registry is responsible for applying for and obtaining additional ethics and other permissions required for their participation in EUROLinkCAT.

Dissemination The CDM and associated documentation, including linkage and standardisation procedures, will be available post-EUROLinkCAT thus facilitating future local, national and European-level analyses to improve healthcare. Recommendations to improve the accuracy of routinely collected data will be made.

Findings will provide evidence to inform parents, health professionals, public health authorities and national treatment guidelines to optimise diagnosis, prevention and treatment for these children with a view to reducing health inequalities in Europe.

Strengths and limitations of this study

- The implementation of a common data model enables the same centrally developed syntax script to be run in all registries which is efficient and ensures standardisation of analysis.
- The use of a reference population allows country differences to be adjusted for and enables more accurate comparisons of the burden of disease attributable to congenital anomalies (CAs) across countries to be made.
- Merging births with CAs to their records in routine healthcare data enables a detailed evaluation of the accuracy of the data and enables improvements to be suggested.
- Being unable to share individual case data or aggregate data that might be disclosive reduces the power of the analysis that can be performed, particularly for rare anomalies.
- Only specific areas in Europe are represented, with a lack of data in particular from Eastern Europe.

INTRODUCTION

Over 130 000 children born in Europe every year will have a major congenital anomaly (CA), equivalent to 2.5% of all European births. CAs include structural defects, chromosomal anomalies and genetic syndromes. CAs are a leading cause of perinatal and infant mortality, especially in developed countries.¹ From 2003 to 2012, CAs were associated with about 40% of all infant deaths in Sweden and England.² There is a large variation in child death rates across Europe; in 2013 the child death rates (age 0–14 years) were 60% higher in the UK and Belgium compared with Sweden, with an additional 10 countries being 30% higher than Sweden.³

To identify potentially preventable and remedial causes it is important to investigate the health inequalities in survival in children with CA across Europe.

It has been shown that relying on death certificates as a source of information on mortality due to CAs does not provide an accurate assessment of the survival for children with specific CAs. Death certificates state the direct or primary cause of death which may be infection, seizures or others and therefore may not mention the CA.⁴ Copeland and Kirby⁴ concluded that the only way to accurately study mortality and survival in children with rare CAs is to pool data across CA registries and link these to death registries.

Advances in fetal, neonatal and paediatric care have improved outcomes for individuals with some CAs, for example, Down syndrome (DS)^{5–7} and cardiac anomalies. Several studies have shown that children with CAs account for a very high proportion of all hospital admissions.^{8,9} However, there is a lack of information on the length of hospital stays for children with specific CAs, with most studies concerning children with DS, orofacial clefts or congenital heart defects (CHD).¹⁰ Often hospital stays are investigated for the first 2 or 3 years of a child's life.^{11–14} However, Wehby *et al* showed that hospital admissions for those born with orofacial clefts were increased at all ages up to 60 years of age.¹⁵ Rarely has length of hospital stay been related to other factors, such as social class. Two studies (Derrington 2013 in the USA to 3 years of age and Hung 2011 in Taiwan for all ages) both identified other factors such as ethnicity and socioeconomic factors as important influences on the length of inpatient stays in children and adults with DS.^{12,16}

The proportion of children born with a CA surviving beyond infancy is increasing.^{5,6} How these children are performing in school and their additional educational needs is therefore becoming increasingly important as there may be a growing population of children and young people requiring additional support and resources in the future. However, apart from the more common genetic syndromes, there is a paucity of information about this.^{17,18} The American Heart Association reviewed the literature on children with CHD and concluded that they are at an increased risk of developmental delay, even once the frequent occurrence of genetic syndromes has been taken into account, particularly for neonates or infants requiring open heart surgery.¹⁹ Wehby *et al* also showed that children with isolated orofacial clefts were at a much greater risk of low achievement at school than their classmates.²⁰ A systematic review of neurocognitive outcomes following general anaesthesia and surgery in children concluded that exposure to general anaesthesia in young children did affect their development in some neurocognitive domains.²¹ However, the authors recommended that the effects of surgery should be considered separately for each specific anomaly.

EUROCAT (<https://eu-rd-platform.jrc.ec.europa.eu/eurocat>) is a European network of population-based CA registries, which started in 1979 and has expanded to

include 39 registries in 21 countries covering more than 29% of European births (1.7 million) per year.^{22–24} The main objectives of EUROCAT are to provide essential epidemiological information and surveillance on CAs in Europe, to evaluate the effectiveness of primary prevention and to assess the impact of developments in prenatal screening.^{25,26} Hence, the emphasis is on information collected up to a baby's first year of life.

The aim of EUROLINKCAT is to investigate the survival, morbidity and educational outcomes of children with specific CAs for the first 10 years of their lives by linking births with CAs in EUROCAT registries to electronic healthcare and education databases. The availability of population-based data on births with CAs across the EUROCAT network will enable survival, morbidity and education to be investigated for specific CAs as well as differences in these outcomes across Europe according to specific risk factors and social inequalities to be explored.

Electronic healthcare data are increasingly being used by researchers to investigate the epidemiology of CAs, rather than using information from CA registries. Such healthcare data have often been found to be incomplete.^{27–31} A small number of registries will analyse the maternal pregnancy records for women registered as having had a termination of pregnancy for a fetal anomaly (TOPFA) in EUROCAT. This will enable the accuracy of routine information on TOPFAs to be evaluated. The accuracy of CA coding in live births will be evaluated by comparing the EUROCAT data for live births with the CA diagnosis from the electronic healthcare databases covering both inpatient and outpatient visits. The information on death certificates will be compared with the anomalies recorded in EUROCAT. Recommendations will be developed to enable the maximum information from electronic healthcare data to be extracted for research purposes and to quantify the amount of data that cannot be obtained.

This paper describes the design of the study, the methods used to obtain and analyse the linked data and evaluates the first 3 years' progress of EUROLINKCAT.

METHODS AND ANALYSIS

Design and setting

In 2017, all EUROCAT registries were invited to participate in the EUROLINKCAT study. Twenty-two registries from 14 countries agreed to participate and to link all live births with a CA registered in their registries and born from 1 January 1995 to 31 December 2014. Almost all EUROCAT registries send anonymised data on CAs occurring in all live births, fetal deaths from 20 weeks gestation and TOPFAs to the EUROCAT central database. Comprehensive coding instructions³² and the use of the EUROCAT Data Management Programme (EDMP) to import data into the central database ensure that standard variables, definitions and coding are used by all registries in the network. CAs are coded locally using the WHO International Statistical Classification of Diseases

and Related Health Problems (ICD) 9th or 10th Revision with the British Paediatric Association code extension offering more specificity (table 1). Cases with minor anomalies only are excluded (see EUROCAT Guide 1.4, Minor Anomalies for Exclusion (V.14.10.14)). Registries can code up to nine anomalies for each case and provide additional information in the specified text fields. Based on the ICD-9 or ICD-10 codes present, cases are automatically assigned by EDMP to defined major CA subgroups in accordance with the EUROCAT Guide 1.4. A case with more than one major anomaly may be assigned to more than one subgroup. Since 2015, the central database has been hosted by the European Commission Joint Research Centre in Ispra (Italy).

Tables 1 and 2 provide a list of the 81 EUROLINKCAT CA subgroups which include structural anomalies, genetic syndromes and chromosomal anomalies that will be investigated. There are 60 EUROCAT subgroups (table 1) and an additional 21 new CA subgroups not defined in EUROCAT (table 2). The subgroups have been identified as being reasonably homogeneous to provide meaningful information and also to be prevalent enough to enable sufficiently precise estimates to be obtained from the analyses. For example, the EUROCAT subgroup 'Chromosomal' was not included as it includes all genetic syndromes, but specific syndromes such as DS (a EUROCAT subgroup) and Di George syndrome (a new subgroup) were included. For some analysis, such as mortality, DS children will be analysed according to whether they have a cardiac anomaly and/or a gastrointestinal anomaly, as these are common and are likely to influence their survival. As outcomes are expected to be more severe for children with multiple and more complex CAs, analyses are also performed separately for children with isolated anomalies or with multiple anomalies defined according to the methodology by Garne *et al.*³³ Isolated anomalies are defined as a CA in one organ system only or with a known sequence where multiple CAs cascade as a consequence of a single primary anomaly. Multiple anomalies are defined as two or more major structural CAs in different organ systems, where the pattern of anomalies has not been recognised as part of a syndrome or sequence.

Linkage

Table 3 provides details of the linkages originally planned by the EUROCAT registries and the current linkages occurring (as of August 2020). The reasons why some registries could not link their data are explored in detail in another paper submitted for publication—they include not being able to obtain the necessary permissions, relevant outcomes not being recorded in specific data sources and the time scale for the data supply being after the end of the study's funding. Currently, 19 registries are linking their data to mortality records, 15 plan to link to hospital in-patient records and 7 to prescription records for the work package that will consider morbidity for children born with a CA. At the time of writing, nine registries plan

to link their information on children with CAs to education records. To evaluate the accuracy of the routine healthcare data, five registries are additionally linking to outpatient data and four will also link to pregnancy information recorded in the mother's health records about TOPFAs. The 19 registries survey over 6 million births in the population.

For the evaluation of survival and morbidity, each child will be followed up for a maximum of 10 years. This age cut-off has been chosen to enable enough children to be identified and followed up; a longer follow-up would mean fewer children would be eligible as currently national or local electronic healthcare record sources often do not go back more than 10 years. For education the maximum follow-up is until the end of compulsory school age (typically 16 across participating countries), although for some registers data are only available for a shorter period of follow-up. The longer follow-up was chosen, because in Finland there are no national education tests and national data on education attainment are available at the age of 15–16 (9th grade).

Reference population

Where possible, each registry obtained information from electronic healthcare records, prescription records and education records on children without a CA. The definition of these 'control' cohorts will vary according to the registry, ranging from all children in the same population covered by the registry to a 10% random sample of children stratified by birth year and child's sex. The use of such a reference population is essential in interpreting differences across countries, as it will provide information on key outcomes, such as duration of hospital stays and medication prescribing, on children without reported anomalies, which is expected to vary by country. Table 4 provides information on the reference populations being identified.

Standardisation and common data models

EUROCAT registries submit 96 core and non-core variables to the EUROCAT central database providing pseudonymised information on the baby and mother, diagnosis, karyotype (if known), exposure, family history and sociodemographic details. These have already been standardised and table 5 lists the 52 variables of which 34 are core variables and their common coding scheme that are used in the EUROLINKCAT study. In contrast, all the data obtained from linkage have to be standardised to a common format, as the healthcare and educational systems across Europe use different native languages and coding classification schemes. To do this, each registry provided their data dictionaries describing the variables in their local databases including the variable names, format, definitions and coding schemes.

Table 6 shows how the variable identifying the sex (male and female) of the child is coded in the different registries with different variable names, different formats and different coding schemes. For each substudy in

Table 1 EUROCAT congenital anomaly subgroups in EUROLinkCAT

EUROCAT subgroups	ICD-10-BPA	ICD-9-BPA
All anomalies*	Q-chapter, D215, D821, D1810†, P350, P351, P371	74, 75, 27910, 2281†, 76076, 76280, 7710, 7711, 77121
Structural anomalies		
Spina bifida	Q05	741
Hydrocephalus	Q03	7423
Severe microcephaly	Q02	7421
Congenital cataract	Q120	74332
Congenital heart defects (CHD)	Q20–Q26	745, 746, 7470–7474
Severe CHD	Q200, Q201, Q203, Q204, Q212, Q213, Q220, Q224, Q225, Q226, Q230, Q232, Q233, Q234, Q251, Q252, Q262	74500, 74510, 7452, 7453, 7456, 7461, 7462, 74600, 7463, 7465, 7466, 7467, 7471, 74720, 74742
Transposition of great vessels	Q203	74510
Ventricular septal defect	Q210	7454
Atrial septal defect (ASD)	Q211	7455
Atrialventricular septal defect (AVSD)	Q212	7456
Tetralogy of Fallot	Q213	7452
Pulmonary valve stenosis	Q221	74601
Aortic valve atresia/stenosis	Q230	7463
Mitral valve anomalies	Q232, Q233	7465, 7466
Hypoplastic left heart	Q234	7467
Coarctation of aorta	Q251	7471
Patent ductus arteriosus (PDA) as only CHD in term infants (GA+37 weeks)	Q250	7470
Cystic adenomatous malformation of lung	Q3380	No code
Cleft lip with or without cleft palate	Q36, Q37	7491, 7492
Cleft palate	Q35	7490
Oesophageal atresia with/without trachea-oesophageal fistula	Q390–Q391	75030–75031
Duodenal atresia or stenosis	Q410	75110
Atresia or stenosis of other parts of small intestine	Q411–Q418	75111–75112
Ano-rectal atresia and stenosis	Q420–Q423	7521–75124
Diaphragmatic hernia	Q790	75661
Gastroschisis	Q793	75671
Omphalocele	Q792	75670
Multicystic renal dysplasia	Q6140, Q6141	75316
Congenital hydronephrosis	Q620	75320
Hypospadias	Q54	75260
Limb reduction defects	Q71–Q73	7552–7554
Craniosynostosis	Q750	75600
Chromosomal anomalies		
Down syndrome	Q90	7580
Trisomy 13	Q914–Q917	7581

Continued

Table 1 Continued

EUROCAT subgroups	ICD-10-BPA	ICD-9-BPA
Trisomy 18	Q910-Q913	7582
Turner syndrome	Q96	75860, 75861, 75862, 75869
Klinefelter syndrome	Q980-Q984	7587
Rare structural anomalies with a EUROCAT subgroup		
Encephalocele	Q01	7420
Arhinencephaly/ holoprosencephaly	Q041, Q042	74226
Anophthalmos/microphthalmos	Q110, Q111, Q112	7430, 7431
Anophthalmos	Q110, Q111	7430
Congenital glaucoma	Q150	74320
Anotia	Q160	74401
Common arterial truncus	Q200	74500
Double outlet right ventricle	Q201	No code
Single ventricle	Q204	7453
Tricuspid atresia and stenosis	Q224	7461
Ebstein's anomaly	Q225	7462
Pulmonary valve atresia	Q220	74600
Hypoplastic right heart	Q226	No code
Aortic atresia/interrupted aortic arch	Q252	74720
Total anomalous pulmonary venous return	Q262	74742
Choanal atresia	Q300	7480
Hirschsprung's disease	Q431	75130-75133
Atresia of bile ducts	Q442	75165
Annular pancreas	Q451	75172
Indeterminate sex	Q56	7527
Situs inversus	Q893	7593
VATER/VACTERL	Q8726	759895

*All anomalies=all cases of congenital anomaly, excluding cases with only minor anomalies as defined in Section 3.2 in EUROCAT Guide 1.4 for cases born post-2005. Cases with more than one anomaly are only counted once in the 'all Anomalies' subgroup.

†ICD10 code D1810 (ICD9 code 2281) is the code for cystic hygroma.

GA, gestational age; ICD-9-BPA, International Statistical Classification of Diseases and Related Health Problems 9th Revision with the British Paediatric Association; ICD-10-BPA, International Statistical Classification of Diseases and Related Health Problems 10th Revision with the British Paediatric Association.

EUROlinkCAT, a common data model (CDM) containing all variables required for its analyses were developed. All the EUROlinkCAT CDMs contain the variable L_CH_SEX, defined as 'sex of child' and with a coding scheme in integer format of 1=male, 2=female, 3=indeterminate, 9=not known, '.'=not recorded or not available for study. Ulster University (UU) used the information in each registry's data dictionary to create the new EUROlinkCAT 'standardised' L_CH_SEX variable. UU, in collaboration with the registries, created registry-specific syntax scripts to standardise all the variables in the EUROlinkCAT CDMs. Online supplemental appendices 1-2 list all the variables included in the substudies (mortality, electronic healthcare records and prescription records).

The CDMs also specify how the data are stored. For mortality, all the relevant variables occur only once for each EUROCAT case and are stored in the same data file (or table) (figure 1). However, when analysing hospital admissions, each child may have more than one admission and for each admission may receive more than one diagnosis. Therefore, the hospital admissions data are stored in a separate data file (or table) from the diagnoses data and separately from the EUROCAT data on the child; each data file contains a reference key which serves to link all records belonging to one person for analysis (see figure 2). The standardisation syntax scripts from UU specify the separate data files (or tables) for each linkage containing all the variables in the CDMs.

Table 2 New congenital anomaly subgroups in EUROLinkCAT

New subgroups for EUROLinkCAT	ICD-10-BPA	ICD-9-BPA
Structural anomalies		
Anomalies of corpus callosum	Q040	74221
Anomalies of intestinal fixation	Q433	7514
Unilateral renal agenesis	Q600	No code
Accessory kidney	Q630	75330
Bladder exstrophy	Q641	7535
Epispadia	Q640	75261
Posterior urethral valves	Q6420	75360
Prune Belly	Q794	75672
Arthrogryposis multiplex congenita	Q743	75580
Genetic syndromes		
Di George syndrome	D821	27910
Goldenhar syndrome	Q8704	75606
Cornelia de Lange syndrome	Q8712	759821
Noonan syndrome	Q8714	759896
Prader-Willi	Q8715	759872
Beckwith-Wiedemann syndrome	Q8730	759874
Williams syndrome	Q8784	No code
Angelman syndrome	Q8785	No code
Chromosomal anomalies		
Wolff-Hirschhorn syndrome	Q933	75832
Cri-du chat syndrome	Q934	75831
Karyotype XXX	Q970	75885
Sequences		
Pierre-Robin sequence	Q8708	75603

ICD-9-BPA, International Statistical Classification of Diseases and Related Health Problems 9th Revision with the British Paediatric Association; ICD-10-BPA, International Statistical Classification of Diseases and Related Health Problems 10th Revision with the British Paediatric Association.

UU included validation routines in the syntax scripts to determine if the data have been correctly transformed. For example, it is checked that a date of death does not occur prior to the date of birth; while the primary purpose is to ensure the data have been standardised correctly, it can also reveal any errors in the linked data.

A CDM is not being defined for the linked education data as there is limited scope for comparison and pooling of data across countries. This is due in part to inherent differences in the educational stages and systems, the variability in data available, and fewer registries being able to participate (5 of the registries are from England, and 1 from Wales, Denmark, Italy and Finland).

Work of the Standardisation Committee

In addition to defining the CDM and its structure, the EUROLinkCAT Standardisation Committee was responsible for taking other decisions, usually in consideration of local data characteristics, to ensure that data were meaningful and comparable across registries. The most important issues are listed below:

1. Inclusion according to gestational age (GA) at birth: for the mortality study all live births with a GA below 24 weeks were excluded, as these cases could have been miscoded terminations of pregnancy with signs of life at birth. However, after running the mortality analysis, it was noticed locally that there were survivors in EUROCAT registries at GA 23 weeks. For the morbidity studies, the exclusion criterion was lowered to be below 23 weeks.
2. Strength of linkage: the success of data linkage depended on the matching method and type of personal identifiers used. Where a national unique identifier (ID) was available (eg, Denmark, Finland) over 99% of cases were matched, but success rates were generally lower when intermediary databases and a combination of other identifiers (eg, names, postcodes) were required to establish a match, particularly if these were incomplete or incorrect. A standard way of evaluating confidence in a match had to be developed so that decisions on inclusion for analyses could be consistently made, in order to avoid bias.
3. GA groups: the GA at birth was categorised into <28 weeks, 28–31 weeks, 32–36 weeks and ≥37 weeks when analysing survival, but due to small numbers of survivors at under 28 weeks gestation, the two lowest GA categories were combined to <32 weeks.
4. Birth weight: birth weight was categorised into very low birth weight '<1500 g', low birth weight '≥1500 g to <2500 g', normal birth weight '≥2500 g to <4000 g' and high birth weight '≥4000 g'. Births <1000 g were not distinguished from those between 1000 and 1499 g as there were too few cases for the data to be analysed accurately.
5. Singletons versus multiples: there is uncertainty about whether the survival in twins with CAs is lower than that in singletons.^{34–37} Hence, all survival analyses were performed on singletons alone and then multiples and singletons combined (multiples were not analysed alone as for many registries small numbers would limit the analyses that could be performed). This enables the survival of singletons and multiples in children with CAs to be analysed in detail. When examining morbidity, multiplicity was treated as one of the risk factors for increased risk of hospitalisations and lengths of stay to enable any association to be analysed, but with less detail than for survival. The majority of analyses included singletons and multiples combined.
6. Prenatal diagnoses: the GA at prenatal diagnoses was categorised into <22 weeks, 22–31 weeks, ≥32 weeks, GA not known and no prenatal diagnosis. For Finland,

Table 3 EUROCAT congenital anomaly registries in EUROLINKCAT: start year, births in the population up to 2014, live births with an anomaly in the study period and ability to link to mortality, healthcare, prescription and education data

Congenital anomaly registry	Planned start year	Actual start year	Planned to be linked	Linked to mortality data by 8/2020	Linkages occurring (Y), Not occurring (N) and Planned but no longer occurring (Y)							
					Number of live births with an anomaly				Hospital data for child			
					Mortality	Hospital data for child		Mortality	Hospital data for mother		Prescriptions	Education
						In-patient	Out-patient		In-patient	Out-patient		
Belgium: Antwerp	1995	1997	8083	7865	Y	N	N	N	N	N	N	N
Croatia: Zagreb	1995	2011	2232	441†	Y	Y	Y	N	N	N	N	N
Denmark: Funen	1995	1995	2418	2425	Y	Y	Y	Y	Y	Y	Y	Y
Finland	1995	1995	44869	42861	Y	Y	Y	Y	Y	Y	Y	Y
France: Île de la Réunion	2002	NC	3855	NC	Y	Y	N	N	N	N	N	N
France: Paris	1997	1995	13335	11623	Y	N	N	N	N	N	N	N
Germany: Saxony-Anhalt	1995	2005	8821	8698	Y	N	N	N	N	N	N	N
Italy: Emilia Romagna	1995	2008	11447	7327	Y	Y	N	N	N	N	Y	Y
Italy: Tuscany	1995	2005	9827	5187	Y	Y	Y	Y	N	N	Y	Y
Malta	1995	2005	2470	2718	Y	N	N	N	N	N	N	N
Netherlands: Northern	1995	2005	8567	8325	Y	Y	Y	Y	Y	Y	Y	N
Norway	1999	1999	26938	27201	Y	N	N	N	N	N	N	N
Portugal: South	1995	2000*	3425*	2447*	N	Y	N	N	N	N	N	N
Spain: Basque Country	1995	1995	4883	5904	Y	Y	N	N	N	N	Y	N
Spain: Valencian Region	2007	2007	7438	7389	Y	Y	N	N	N	N	Y	N
UK: East Midlands and South Yorkshire	1998	AL	18549	AL	Y	Y	N	N	N	N	Y	Y
UK: Northern England	2000	AL	8617	AL	Y	Y	N	N	N	N	Y	Y
UK: South West England	2005	AL	11671	AL	Y	Y	N	N	N	N	Y	Y
UK: Thames Valley	1995	AL	5142	AL	Y	Y	N	N	N	N	Y	Y
UK: Wales	1998	1998	18239	18128	Y	Y	Y	Y	Y	Y	Y	Y
UK: Wessex	1995	AL	7771	AL	Y	Y	N	N	N	N	Y	Y
Ukraine: West	2005	2006	6166	5835	Y	Y	N	N	N	N	N	N
Total			234 763	155 594†	19	15	5	4	4	4	7	9

The registry in Basque was unable to complete the planned linkages to hospital data due to COVID-19.

*Values for WP4 morbidity linkage have been provided, as mortality linkage was never planned.

†The 441 cases in Zagreb were not included in analyses due to poor quality of the mortality linkage.

AL: awaiting linkage as of August 2020; NC: linkages could not be completed; Y: linkages are no longer planned as of August 2020.

Table 4 Use of a reference population in morbidity and education analyses

Congenital anomaly registry	Reference population
Croatia: Zagreb	Sample of children
Denmark: Funen	Whole population
Finland	Whole population
France: Île de la Réunion	Not provided
Italy: Emilia Romagna	Whole population
Italy: Tuscany	10% of population
Netherlands: Northern	10% of population
Portugal: South	Sample of children
Spain: Basque	Not provided
Spain: Valencian Region	Whole population
UK: East Midlands and South Yorkshire	Aggregate data from population for morbidity and population sample for education
UK: Northern England	Aggregate data from population for morbidity and population sample for education
UK: South West England	Aggregate data from population for morbidity and population sample for education
UK: Thames Valley	Aggregate data from population for morbidity and population sample for education
UK: Wales	Whole population
UK: Wessex	Aggregate data from population for morbidity and population sample for education
Ukraine: West	No longer in morbidity study

the GA was often not recorded, only the trimester of diagnoses. Finland's first trimester diagnoses (week 0 to week 12) mapped exactly to the EUROLINKCAT <22 weeks category. It was decided that Finland's second trimester (week 13 to week 27) diagnoses were assumed to occur at 22–31 weeks and third trimester (week 28 onwards) diagnoses occurred at ≥32 weeks. These assumptions were also checked based on the distribution of those cases in Finland with a 'known' age at discovery and the assumptions held.

- Length of stay (LOS): the LOS of the child in hospital was calculated after excluding the stay associated with the birth. Methods of identifying the birth stay varied in different countries. For hospital admissions where admission and discharge occurred on the same day, the LOS was considered to be 0.5 days. If an admission record was missing a discharge date, then discharge date = date of admission + 2 × (date of latest

Table 5 Standardised variables from the EUROCAT database

EDMP variables used (core variables are shaded in blue)		
Baby and mother		
1	CENTRE	Centre number
2	NUMLOC	Local ID of case
3	BIRTH_DATE	Date of birth
4	SEX	Sex
5	NBRBABY	Number of babies delivered
6	SP_TWIN	Specify twin type of birth, like or unlike, zygosity
7	NBRMALF	Number of malformed in multiple set
8	TYPE	Type of birth
9	CIVREG	Civil registration status
10	WEIGHT	Birth weight
11	GESTLENGTH	Length of gestation in completed weeks
12	SURVIVAL	Survival beyond 1 week of age
13	DEATH_DATE	Date of death
14	DATMO	Date of birth of mother
15	AGEMO	Age of mother at delivery
16	BMI	Maternal body mass index
17	RESIDMO	Mother's residence code
Diagnosis		
19	WHENDISC	When discovered
20	CONDISC	Condition at discovery
21	AGEDISC	If prenatally diagnosed, gestational age at discovery
22	FIRST PRE	First positive prenatal test
24	KARYO	Karyotype of infant/fetus
25	SP_KARYO	Specify karyotype
26*	GENTEST	Genetic test
27*	SP_GENTEST	Specify genetic test
28	PM	Postmortem examination
29	SURGERY	First surgery for malformation performed or planned
30	SYNDROME	Syndrome
31	SP_SYNDROME	Specify syndrome
32	MALFO1	Malformation
33	SP_MALFO1	Specify malformation
34	MALFO2	As MALFO1
35	SP_MALFO2	Specify malformation
36	MALFO3	As MALFO1
37	SP_MALFO3	Specify malformation
38	MALFO4	As MALFO1
39	SP_MALFO4	Specify malformation
40	MALFO5	As MALFO1
41	SP_MALFO5	Specify malformation
42	MALFO6	As MALFO1
43	SP_MALFO6	Specify malformation

Continued

Table 5 Continued

EDMP variables used (core variables are shaded in blue)		
44	MALFO7	As MALFO1
45	SP_MALFO7	Specify malformation
46	MALFO8	As MALFO1
47	SP_MALFO8	Specify malformation
57	OMIM	OMIM code/type of Mendelian inheritance
Exposure and family history		
58	ASSCONCEPT	Assisted conception (where available)
59†	OCCUPMO	Mother's occupation at time of conception
Sociodemographic		
91	MATEDU	Maternal education
92*	SOCM	Socioeconomic status of mother
93 *	SOCF	Socioeconomic status of father
94	MIGRANT	Migrant status
Derived variables		
	Byear	Year of birth
	birth_type	Live birth, stillbirth, spontaneous abortion, TOPFA, not known definitions of stillbirths and spontaneous abortions vary between regions. This variable recodes birth type according to EUROCAT's specifications: cases with gestational age ≥20 weeks are recoded as 'stillbirths' (irrespective of the local definition of stillbirth/spontaneous abortion).
	casestatus	Only cases with casestatus=1 or 2
	al1-al114	EUROCAT subgroups: (0=No, 1=Yes). Based on EUROCAT coding in Guide 1.4
	mult_malf	Algorithm for case classification into isolated and multiple anomalies

*See work of the Standardisation Committee (viii).

†EUROCAT Guide 1.4 use ISCO-08 classifications.

EDMP, EUROCAT Data Management Programme; TOOPFA, termination of pregnancy for a fetal anomaly.

procedure–date of admission). The date of discharge was set to the date of the child's 10th birthday or the end of the study period if it was after either of these two dates.

- Socioeconomic status (SES): all registries had different variables that could be considered to be a measure of the mother's SES. The variables included maternal occupation, maternal education and index of multiple deprivation derived from residential codes at birth. Registries were asked to select the variable they believed was the most relevant and to recode their selected SES proxy variable into three groups of approximately equal proportion to enable comparing

Table 6 Coding of male or female of the live births in different linked databases in EUROLINKCAT

Centre	Variable name	Code	
		Male	Female
UK: Wales	DEC_SEX_CD	1	2
Germany: Saxony-Anhalt	EF306	1	2
Finland	SUKUP	1	2
Italy: Tuscany	SESSO	1	2
France: Île de la Réunion	SexeDefunt	1	2
France: Paris	SexeDefunt	1	2
Netherlands: Northern	geslacht	1	2
Croatia: Zagreb	GENDER	M or 1	F or 2
Ukraine: West	CH_SEX	1	2
Belgium: Antwerp	SEX	1	2
Norway	KJONN	1	2
UK: England	sex	1	2
Spain: Basque Country	SEXO	1	6
Spain: Valencian Region	SEXO	1	6
Malta	gender	M	F
Italy: Emilia Romagna	SEX	M	F
Denmark: Funen	C_SEX	M	K

between, for example, mothers in the highest group to mothers in the lowest group. The effect of SES on survival would be analysed using Cox proportional hazard models within each registry. However, only seven registries were able to provide a proxy SES variable that was reasonably complete for some or all of the time period of the study. It was also planned to investigate the association between risk factors such as birth weight after adjusting for SES, but due to the lack of information on SES this was not included in further multivariable analyses.

- Maternal country of birth: it was determined that the maternal country of birth variable would be used as a proxy for non-European ethnic origin, as we were aware that ethnic origin is poorly recorded. However, for those registries with reasonably complete data on this, almost 100% of children were reported as being of European ethnic origin. Therefore, this variable was not included in subsequent analyses as the number of children considered as 'non-European ethnic origin' was too small to analyse.
- Cause of death: cause of death based on the death certificates was classified for deaths <1 year and for 1–9 years separately. Death related to preterm birth is very common in the first year after birth, but not as relevant to children at 1–9 years of age. Injuries and poisoning are more common after the first year. The main causes of deaths were classified into six groups for deaths <1 year and 11 groups for children aged 1–9 years. When working with the results tables it was clear that many of these classification groups included many small numbers and data could not be

Patient data - 1 row per EUROCAT case							
Child ID	Child date of Birth	Child date of death	Type Matching	Variables relating to linkage	Variables relating to child	Variables relating to mortality	Variables relating to mother

Figure 1 Structure of mortality and EUROCAT data used for analysing children's survival.

extracted from the databases. For some registries it was only possible to give cause of death as either 'congenital anomaly' or else 'any other cause of death'.

11. Surgery: a number of different coding systems were used across registries to code surgeries and other procedures (eg, NCSP by NOMESCO, ICD-9-CM, OPCS-4). Frequency lists for all codes describing surgeries and other procedures were obtained from the linked datasets. Two paediatricians then independently determined if a code was a surgery or for another procedure and then a consensus between the two clinicians was reached over codes classified as codes for surgeries. Further subdivision into anomaly-specific surgeries was carried out for anomalies for which specific

surgeries could be identified that would be expected to be performed on these children.

12. Intensive care: it was planned to analyse the number of days in intensive care, however, only five registries could provide this. Therefore, only whether a child had ever been admitted to intensive care was analysed rather than their LOS.
13. Ventilation: it was planned to analyse the number of days on ventilation. However, as it was decided that the LOS in intensive care was not going to be analysed, the same decision was made for ventilation and only whether a child had ever been on ventilation was analysed.




Figure 2 Structure of hospital admissions, prescription data and EUROCAT data used for analysing children's morbidity. ATC code, Anatomical Therapeutic Chemical Classification System code; ICU, intensive care unit.

Assessment of quality of linkage and quality of linked data

Many registries linked their data to National Vital Statistics, which are databases that record all live births with follow-up until the child dies or emigrates outside the country/region of interest. Therefore, for these registries for the survival analysis, any child whose record was not in the National Vital Statistics Database was judged to be a non-match and overall linkage could be assessed. Some registries were only able to link to death certificates which meant that a non-match, that is, no death certificate found, was assumed to indicate that the child was still alive. The data from these registries were only included in the survival analyses if there was additional information about the quality of the linkage. For example, in Malta, due to the small well-defined population, there was confidence that all deaths had been identified.

In some countries all national databases use the same unique ID number (eg, Finland). So, identifying a child in the National Vital Statistics meant that there was confidence that any hospital stays up to 10 years of age would also be identified. For other registries, as not all children were likely to be admitted to hospital, each case was searched for in, not only the in-patient hospital database (which included the mother's visit for the birth), but also any other healthcare databases (such as outpatient, primary-care or prescription databases) for longer than the 10 years of follow-up in the study. The lack of information in any healthcare database was judged to mean a non-match. Sensitivity analysis was performed to assess if there were differences in results if the non-matches were included.

For the education data, all children known to be alive were assumed to be included in the National Education Databases and therefore any case not identified was assumed to be a non-match.

Syntax scripts were developed centrally by St George's, University of London (SGUL) to evaluate the accuracy of the linkage and identify any factors leading to missed links (eg, deaths within the first week of life). For each registry, the proportion of births in any single year of data that have not been linked is calculated and the data from any year with less than 85% of cases linked will be excluded from further analyses. Second, the quality of linked data items was also evaluated: a variable that was >20% missing in a year would be excluded from any analysis in which it featured; and variables that were recorded by both the CA registry and the linked database would be compared for agreement, by year. In general, it was found that data quality was poorer in the earlier years and tended to improve over time; however, if data quality fluctuated across the years, then only the longest consecutive period where quality was above the threshold would be analysed.

Statistical analysis

Protocols and syntax scripts are developed centrally to create aggregate data and perform specific analyses on the individual cases in each standardised data set in

STATA (V.13 and upwards). This allows each register to submit aggregated data and analytical results (eg, Kaplan-Meier estimates, HRs and CIs), rather than individual case data, to the EUROLINKCAT central results repository (CRR) at UU, UK using a secure web platform. UU collates the aggregate data and results and provides these data to the researchers responsible for the different analyses and publications. Multi-centre European analyses will be performed by combining the individual registries' aggregated data and analytic results, using meta-analytic techniques. Additional work is required to develop suitable models for combining survival data from several registries when the sample sizes are very small as observed in many registries.

Small number restrictions (statistical disclosure control)

Four countries have limitations on the release of aggregate data and analytic results if the numbers of births involved are very small (generally under eight births). This situation arises in many analyses involving specific CAs, as CAs are rare, with some affecting less than 1 in 10 000 live births. Solutions to enable the maximum amount of data to be included in all multi-centre European analyses varied according to country. The Northern Netherlands released data if all exported results were rounded to the nearest 5. Rounding all frequencies ensures that original numbers cannot be inferred. For Denmark, a few named researchers at SGUL and UU were allowed access to the aggregate data for the purpose of collating and including in pooled-analysis, on condition that it was securely stored and processed; that any individual results involving fewer than five people were not released; and that personal identification was not possible from any released results. The SAIL databank (Wales) provided data to the CRR with the requirement that aggregate data on fewer than five people were not released and could not be calculated from any information in the public domain. The registry from Antwerp, Belgium could not release any information on three or fewer cases.

Patient and public involvement

A series of focus groups has been held in different European countries involving parents with a child with one of four predefined CAs with different health problems covering learning difficulties, physical disabilities, visible defects and non-visible defects with higher mortality. The four anomalies selected were: CHD requiring surgery (referred to as severe CHD—a usually non-visible defect with high mortality), cleft lip (a visible defect often with speech problems), spina bifida (a physical disability with associated incontinence problems) and DS (Trisomy 21; a visible defect with learning difficulties and often associated with CHD). The focus groups have investigated parental experiences of having a child with one of the above anomalies and assessed parental research priorities and a paper will be published in due course.

In addition, a European survey concerning the diagnosis, medical care, education and everyday life will be

distributed to parents across Europe with children with the same four CAs as described above. Registries will ensure the questions in the survey are appropriate for their country (eg, the provision of health services, given how this differs in various European countries) and will translate the survey into their native language with back translation to confirm the accuracy of the translation. The aim is for the survey to be distributed via social media by parent support groups across Europe to engage with a wide spectrum of parents.

DISCUSSION

The aggregate data and results from the CRR in EUROLINKCAT will provide important information on the survival, morbidity and education of children born with a CA in Europe. Researchers in each CA registry will be encouraged to also perform specific local analysis, in order to fully exploit the research potential of linked datasets. The establishment of a method of standardising data from each registry linkage into a CDM provides valuable infrastructure enabling future multi-national studies to be performed in an efficient manner and new registries to become involved.

The strength of this study is that the researchers are a multidisciplinary group, many of whom have collaborated successfully for many years through being members of EUROCAT. In addition, EUROLINKCAT is able to build on all the standardisation procedures already established in EUROCAT. The implementation of a CDM enables the same centrally developed syntax script to be run in all the different registries which is efficient and also ensures standardisation of analysis across the registries. The use of a reference population when analysing healthcare data will aid in the identification of the source of differences between registries (eg, average LOS in hospitals will differ) and therefore enable us to better quantify the burden of disease attributable to CAs in each country. Comparisons of the accuracy of healthcare databases with respect to recording CA cases will be informative and enable improvements in those areas with less accurate data. We will also be developing recommendations on how to use the available healthcare data in an optimal way to provide information on children with CAs in areas without active CA registries.

One of the challenges of EUROLINKCAT is the ability of the CA registries to link their data to external data sources due to different local data information governance issues and the availability of suitable electronic healthcare databases. This requires flexibility in including registries in only specific subprojects and acceptance that not all registries may be able to perform the linkages planned. In addition, some registries require support from other partners in all aspects of the project, including applying for ethics permissions, adapting protocols, standardising data and running statistical syntax scripts. The restriction of not being able to share individual case data and also aggregate data that might be disclosive or identifiable

means that all analyses must be performed locally using a generic modelling strategy. This does limit the use of iterative procedures to explore data in detail. The major limitation to the study is that only specific areas in Europe are represented, with a lack of data in particular from Eastern Europe. Interpretation of differences across Europe is challenging as it will be essential to interpret results in the light of knowledge about the differences in healthcare and education practices across Europe.

The EUROLINKCAT project will enable important hypotheses concerning the survival, health and education of children with CAs in Europe to be investigated. The standardised methods and CDMs will all be available freely on the EUROLINKCAT website and will be available for use in future research projects to benefit from and build on this work, so as to enable other multi-centre European projects to exploit routine healthcare data available in Europe.

ETHICS AND DISSEMINATION

The CA registries have the required ethics permissions and procedures for routine surveillance, data collection and transmission of anonymised data to the EUROCAT central database, according to national guidelines and they were required to submit evidence of these permissions to the EUROLINKCAT ethics portfolio. Local registries follow national legislation as to whether parental consent is needed for registration of babies with anomalies. Each registry was responsible for applying for and obtaining the additional ethics and other permissions (eg, data sharing agreements) required to link and analyse their data for EUROLINKCAT. This was an extremely lengthy process in some countries as the original data collection did not include expectation or consent for the data to be used in research, and a new legal basis had to be established. Additional assurances and procedures were adopted by registries (eg, publication of privacy notices) to ensure compliance with the General Data Protection Regulation which came into force on 25 April 2018 in the European Union. A checklist of minimum specifications for data storage/backup was completed by each registry. Three registries took over 3 years to get the ethics, legal basis, data protection, information governance and data sharing agreements in place. UU obtained ethics permission for the CRR.

Each registry participating in the focus groups with parents was responsible for ensuring the correct ethics approvals were in place. Similarly, the registries participating in the dissemination of the parents' survey will be responsible for ensuring the necessary ethics permissions are obtained.

An Ethics and Data Protection Advisory Board consisting of three independent advisors with the relevant expertise monitor all ethical considerations in this project.

The CRR will be used for multiple studies and the results from these will be disseminated in peer reviewed papers and conference presentations. It is hoped that the

experience gained with distributing the parents' survey using parent support groups and social media will also lead to development of a framework to enable dissemination of results to be made more directly to parents. In addition, a series of reports will be written including recommendations for improving the collection and analysis of data on CAs in routinely collected data in the healthcare databases.

Author affiliations

¹Population Health Research Institute, St George's University of London, London, UK

²Paediatric Department, Hospital Lillebaelt, Kolding, Denmark

³Faculty of Life and Health Sciences, Ulster University, Coleraine, UK

⁴Children's Hospital Zagreb, Centre of Excellence for Reproductive and Regenerative Medicine, Medical School University of Zagreb, Zagreb, Croatia

⁵Biomedical Computing Limited, Battle, UK

⁶Polish Registry of Congenital Malformations, Chair and Department of Medical Genetics, Poznan University of Medical Sciences, Poznan, Poland

⁷IMER Registry (Emilia Romagna Registry of Birth Defects), University Hospital of Ferrara, Emilia-Romagna, Italy

⁸Istituto di Fisiologia Clinica, Consiglio Nazionale delle Ricerche, Pisa, Italy

⁹Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

¹⁰Malformation Monitoring Centre Saxony-Anhalt, Otto von Guericke University Medical Faculty, Magdeburg, Germany

¹¹Department of Genetics, University Medical Centre Groningen, Groningen, The Netherlands

Collaborators All below named members of the EUROlinkCAT Consortium have contributed to the development of the protocol with their knowledge and experience of the surveillance of congenital anomalies. Names are given in Consortium organisation numerical order: Population Health Research Institute, St George's, University of London, UK: Abigail Reid, Liz Limb, Gillian Briggs, Nicholas Connor; Hospital Lillebaelt, Region Syddanmark, Denmark and Section of Epidemiology, University of Copenhagen, Denmark: Stine Kjaer Urhoj; Hospital Lillebaelt, Region Syddanmark, Denmark: Mads Damkjær, Christina Neergaard Pedersen; Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK: Svetlana V. Glinianaia; Emilia Romagna Registry of Birth Defects (IMER), University of Ferrara, Italy: Gianni Astolfi, Annarita Armaroli; Neonatal Intensive Care Unit, IMER Registry, Department of Medical Sciences, University Hospital of Ferrara, Italy: Elisa Ballardini; Territorial Care Service - Emilia Romagna Health Authority, Italy: Aurora Puccini; Childrens Hospital Zagreb, Croatia: Ljubica Boban; Institute of Clinical Physiology, National Research Council (Consiglio Nazionale delle Ricerche-Istituto di Fisiologia Clinica), Pisa, Italy: Silvia Baldacci, Alessio Coi, Lorena Mezzasalma, Michele Santoro; University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands: L. Renée Lutke, Nicole H.A. Siemensma-Mühlenberg; Public Health Wales National Health Service Trust, Wales, UK: David Tucker; Institut National de la Santé et de la Recherche Médicale, Paris, France: Babak Khoshnood, Nathalie Bertille, Nathalie Lelong, Makan Rahshenas; Rare Diseases Research Unit, Foundation for the Promotion of Health and Biomedical Research in the Valencian Region, Valencia, Spain: Clara Cervero Carbonell, Laia Barrachina Bonet, Laura Garcia Villodre; Rare Diseases Research Unit, Foundation for the Promotion of Health and Biomedical Research in the Valencian Region, Valencia, Spain and Public Health and Preventive Medicine Department, University of València, Spain and Public Health Regional Health Administration (DG Salud Pública y Adicciones), Generalitat Valenciana, Valencia, Spain: Óscar Zurriaga; Polish Registry of Congenital Malformations, Department of Medical Genetics, Poznan University of Medical Sciences, Poznan, Poland: Anna Jamry-Dziurla; Finnish Institute for Health and Welfare, Finland: Sonja Kiuru-Kuhlefelt, Mika Gissler, Anna Heino; OMNI-Net for Children International Charitable Fund, Rivne, Ukraine: Wladimir Wertenlecker; OMNI-Net for Children International Charitable Fund, Rivne, Ukraine and Khmelnytsky City Children's Hospital, Khmelnytsky, Ukraine: Diana Akhmedzhanova, Nataliia Zymak-Zakutnia; OMNI-Net for Children International Charitable Fund, Rivne, Ukraine and Rivne Regional Medical Diagnostic Center, Rivne, Ukraine: Lyubov Yevtushok, Lyubov Ostapchuk, Oksana Tsih; OMNI-Net for Children International Charitable Fund, Rivne, Ukraine and Volyn Regional Children's Territorial Medical Center, Lutsk, Ukraine: Serhii Lapchenko; Epidemiology Department, National Registry of Congenital Anomalies, National Institute of Health Doctor Ricardo Jorge

(Instituto Nacional de Saúde Doutor Ricardo Jorge), Lisbon, Portugal: Carlos Matias Dias, Ausenda Machado, Ana João Santos, Lilianna Antunes, Paula Braz; Register of Congenital Malformations, Centre Hospitalier Universitaire de La Réunion, Île de la Réunion, France: Hanitra Randrianaivo-Ranjatoelina, Benedicte Bertaut Nativel; Provinciaal Instituut voor Hygiëne (PIH), Antwerp, Belgium: Vera Nelen, Guy Thys, Elly Den Hond; Departamento de Salud Gobierno Vasco, Basque Country, Spain: Olatz Mokoroa Carollo; Swansea University, Wales, UK: Daniel Thayer, Ieuan Scanlon, Sue Jordan; Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway and Division of Mental and Physical Health Norwegian Institute of Public Health, Bergen, Norway: Kari Klungsøyr; Directorate for Health Information and Research, G'Mangia, Malta: Miriam Gatt.

Contributors JKM is the principal author and the project's scientific coordinator. ML is the data coordinator. EG is the clinical coordinator. JD, AN, AP, JR and HdW are Steering Committee and Standardisation Committee members with JKM, ML and EG. IB and AL-B are Steering Committee members. AR is a Standardisation Committee member. JT is the senior statistician. JEG the research fellow. HC is the project manager. All authors contributed to, read and approved the final manuscript.

Funding This project has received funding from the European Union's Horizon 2020 Research and Innovation programme under grant agreement No 733001 for 5 years: 1 January 2017 to 31 December 2021. The views presented here are those of the authors only, and the European Commission is not responsible for any use that may be made of the information presented here.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Joan K Morris <http://orcid.org/0000-0002-7164-612X>
Ester Garne <http://orcid.org/0000-0003-0430-2594>
Maria Loane <http://orcid.org/0000-0002-1206-3637>
Ingeborg Barisic <http://orcid.org/0000-0002-9085-6747>
Anna Latos-Bieleńska <http://orcid.org/0000-0002-9152-1674>
Anna Pierini <http://orcid.org/0000-0003-3321-9343>
Judith Rankin <http://orcid.org/0000-0001-5355-454X>
Anke Rissmann <http://orcid.org/0000-0002-9437-2790>
Hermien de Walle <http://orcid.org/0000-0002-1418-8281>
Joachim Tan <http://orcid.org/0000-0003-0462-4761>
Joanne Emma Given <http://orcid.org/0000-0003-4921-1944>
Hugh Claridge <http://orcid.org/0000-0001-5998-2860>

REFERENCES

- Boyle B, Addor M-C, Arriola L, *et al*. Estimating global burden of disease due to congenital anomaly: an analysis of European data. *Arch Dis Child Fetal Neonatal Ed* 2018;103:F22-8.
- Zylbersztejn A, Gilbert R, Hjern A, *et al*. Child mortality in England compared with Sweden: a birth cohort study. *Lancet* 2018;391:2008-18.
- Wang H, Liddell CA, Coates MM, *et al*. Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990-2013: a systematic analysis for the global burden of disease study 2013. *Lancet* 2014;384:957-79.
- Copeland GE, Kirby RS. Using birth defects registry data to evaluate infant and childhood mortality associated with birth defects: an alternative to traditional mortality assessment using underlying

- cause of death statistics. *Birth Defects Res A Clin Mol Teratol* 2007;79:792–7.
- 5 Wu J, Morris JK. The population prevalence of Down's syndrome in England and Wales in 2011. *Eur J Hum Genet* 2013;21:1016–9.
 - 6 Tennant PWG, Pearce MS, Bythell M, *et al.* 20-Year survival of children born with congenital anomalies: a population-based study. *Lancet* 2010;375:649–56.
 - 7 Brodwall K, Greve G, Leirgul E, *et al.* The five-year survival of children with Down syndrome in Norway 1994–2009 differed by associated congenital heart defects and extracardiac malformations. *Acta Paediatr* 2018;107:845–53.
 - 8 Muranjan M, Vijayalakshmi P. The unforeseen toll of birth defects and their economic burden at a tertiary care public Institute in Mumbai. *Indian J Pediatr* 2014;81:1005–9.
 - 9 Simeone RM, Oster ME, Hobbs CA, *et al.* Population-Based study of hospital costs for hospitalizations of infants, children, and adults with a congenital heart defect, Arkansas 2006 to 2011. *Birth Defects Res A Clin Mol Teratol* 2015;103:814–20.
 - 10 Shetty S, Kennea N, Desai P, *et al.* Length of stay and cost analysis of neonates undergoing surgery at a tertiary neonatal unit in England. *Ann R Coll Surg Engl* 2016;98:56–60.
 - 11 Fitzsimons KJ, Copley LP, Deacon SA, *et al.* Hospital care of children with a cleft in England. *Arch Dis Child* 2013;98:970–4.
 - 12 Derrington TM, Kotelchuck M, Plummer K, *et al.* Racial/Ethnic differences in hospital use and cost among a statewide population of children with Down syndrome. *Res Dev Disabil* 2013;34:3276–87.
 - 13 Dawson AL, Cassell CH, Oster ME, *et al.* Hospitalizations and associated costs in a population-based study of children with Down syndrome born in Florida. *Birth Defects Res A Clin Mol Teratol* 2014;100:826–36.
 - 14 Hook-Dufresne DM, Yu X, Bandla V, *et al.* The economic burden of gastroschisis: costs of a birth defect. *J Surg Res* 2015;195:16–20.
 - 15 Wehby GL, Pedersen DA, Murray JC, *et al.* The effects of oral clefts on hospital use throughout the lifespan. *BMC Health Serv Res* 2012;12:58.
 - 16 Hung W-J, Lin L-P, Wu C-L, *et al.* Cost of hospitalization and length of stay in people with Down syndrome: evidence from a national hospital discharge claims database. *Res Dev Disabil* 2011;32:1709–13.
 - 17 O'Toole C, Lee AS-Y, Gibbon FE, *et al.* Parent-mediated interventions for promoting communication and language development in young children with Down syndrome. *Cochrane Database Syst Rev* 2018;10:CD012089.
 - 18 Iyer NP, Tucker DF, Roberts SH, *et al.* Outcome of fetuses with Turner syndrome: a 10-year congenital anomaly register based study. *J Matern Fetal Neonatal Med* 2012;25:68–73.
 - 19 Marino BS, Lipkin PH, Newburger JW, *et al.* Neurodevelopmental outcomes in children with congenital heart disease: evaluation and management: a scientific statement from the American heart association. *Circulation* 2012;126:1143–72.
 - 20 Wehby GL, Collett BR, Barron S, *et al.* Children with oral clefts are at greater risk for persistent low achievement in school than classmates. *Arch Dis Child* 2015;100:1148–54.
 - 21 Clausen NG, Kähler S, Hansen TG. Systematic review of the neurocognitive outcomes used in studies of paediatric anaesthesia neurotoxicity. *Br J Anaesth* 2018;120:1255–73.
 - 22 Boyd PA, Haeusler M, Barisic I, *et al.* Paper 1: The EUROCAT network--organization and processes. *Birth Defects Res A Clin Mol Teratol* 2011;91 Suppl 1:S2–15.
 - 23 Greenlees R, Neville A, Addor M-C, *et al.* Paper 6: EUROCAT member registries: organization and activities. *Birth Defects Res A Clin Mol Teratol* 2011;91 Suppl 1:S51–100.
 - 24 Kinsner-Ovaskainen A, Lanzoni M, Garne E, *et al.* A sustainable solution for the activities of the European network for surveillance of congenital anomalies: EUROCAT as part of the EU platform on rare diseases registration. *Eur J Med Genet* 2018;61:513–7.
 - 25 Dolk H, Loane M, Teljeur C, *et al.* Detection and investigation of temporal clusters of congenital anomaly in Europe: seven years of experience of the EUROCAT surveillance system. *Eur J Epidemiol* 2015;30:1153–64.
 - 26 Loane M, Dolk H, Kelly A, *et al.* Paper 4: EUROCAT statistical monitoring: identification and investigation of ten year trends of congenital anomalies in Europe. *Birth Defects Res A Clin Mol Teratol* 2011;91 Suppl 1:S31–43.
 - 27 Salemi JL, Tanner JP, Sampat D, *et al.* The accuracy of hospital discharge diagnosis codes for major birth defects: evaluation of a statewide registry with passive case ascertainment. *J Public Health Manag Pract* 2016;22:E9–19.
 - 28 Wang Y, Cross PK, Druschel CM. Hospital discharge data: can it serve as the sole source of case ascertainment for population-based birth defects surveillance programs? *J Public Health Manag Pract* 2010;16:245–51.
 - 29 Andrade SE, Scott PE, Davis RL, *et al.* Validity of health plan and birth certificate data for pregnancy research. *Pharmacoepidemiol Drug Saf* 2013;22:7–15.
 - 30 Blais L, Bérard A, Kettani F-Z, *et al.* Validity of congenital malformation diagnostic codes recorded in Québec's administrative databases. *Pharmacoepidemiol Drug Saf* 2013;22:881–9.
 - 31 Steele A, Johnson J, Nance A, *et al.* A quality assessment of reporting sources for microcephaly in Utah, 2003 to 2013. *Birth Defects Res A Clin Mol Teratol* 2016;106:983–8.
 - 32 EUROCAT. Guide 1.4. Available: http://www.eurocat-network.eu/aboutus/datacollection/guidelinesforregistration/guide1_4 [Accessed 14 Feb 2018].
 - 33 Garne E, Dolk H, Loane M, *et al.* Paper 5: surveillance of multiple congenital anomalies: implementation of a computer algorithm in European registers for classification of cases. *Birth Defects Res A Clin Mol Teratol* 2011;91 Suppl 1:S44–50.
 - 34 Herskind AM, Larsen LA, Pedersen DA, *et al.* Comparison of late mortality among twins versus singletons with congenital heart defects. *Am J Cardiol* 2017;119:1680–6.
 - 35 Gnanendran L, Bajuk B, Oei J, *et al.* Neurodevelopmental outcomes of preterm singletons, twins and higher-order gestations: a population-based cohort study. *Arch Dis Child Fetal Neonatal Ed* 2015;100:F106–14.
 - 36 Wang W, Pan W, Chen J, *et al.* Outcomes of congenital diaphragmatic hernia in one of the twins. *Am J Perinatol* 2019;36:1304–9.
 - 37 Martinka D, Barrett J, Mei-Dan E, *et al.* Respiratory morbidity in late preterm twin infants. *Arch Gynecol Obstet* 2019;300:337–45.

Variables in EUROLinkCAT Common Data Model: Mortality

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
Core variable required for linkage to vital statistics or mortality databases and for calculating age at death			
L_CH_ID	Unique identifier of child A unique ID that links child to another database	As recorded locally	
L_CH_DATE_B	Child's date of birth Needed to calculate age at death.	DDMonYYYY	
L_CH_DATE_D	Child's date of death Needed to calculate age at death.	DDMonYYYY	
Variables relating to linkage			
L_MATCH_TYPE	Match with vital statistics database (or local health care databases)	Numeric	1 = Linked to national/vital statistics database – match 2 = Linked to national/vital statistics database – non-match 3 = Linked to mortality database only – match 4 = Linked to mortality database only – non-match 5 = EUROCAT death only

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
L_CONFIDENCE	Strength of match with vital statistics or health care database. Use local data provider's codes for assessing confidence that the case is correctly matched. If local code unavailable, use suggested coding (see Appendix for full details)	Numeric	1=Excellent 2=Good 3=Fair 4=Poor 9=Not Matched
L_DATE_LOST	Date lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)	DDMonYYYY	.=Not recorded by registry or not available for study
L_YEAR_LOST	Year lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)	YYYY	.=Not recorded by registry or not available for study
L_AGE_L_D	Age lost to follow-up/ linkage <u>in complete days</u>	Numeric (1-4 digits)	.=Not recorded by registry or not available for study
Standardised variables relating to child			
L_CH_YEAR_B	Child's year of birth	YYYY	
L_CH_SEX	Child's sex	Numeric	1 = Male 2 = Female 3=Indeterminate 9 = Not known .=Not recorded by registry or not available for study
L_CH_REG_TYPE	Type of birth/ civil registration of baby	Numeric	1 = Livebirth 2 = Stillbirth 9 = Not known .=Not recorded by registry or not available for study
L_CH_NON_EUR	Citizenship/ Nationality of infant /country of origin	Numeric	1= National 2= Other European 3 = Non-European

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
			4= Non-national (exact nationality not specified) 9 = Not known . = Not recorded by registry or not available for study
L_CH_BW	Child's birthweight (grams)	Numeric	9999=Not known .=Not recorded by registry or not available for study
L_CH_GA_B	Child's gestational age at birth (completed weeks)	Numeric	99=Not known .=Not recorded by registry or not available for study
Variables relating to mortality			
L_CH_STATUS	<p>Outcome status</p> <p>-Died = child is known to have died before 10th birthday or 31 Dec 2015 (whichever earlier)</p> <p>-Alive at 10th birthday = child was born on or before the 31st Dec 2005 and:</p> <p>(i) is definitively known to be alive on 10th birthday; or (ii) there is no information on death or lost to follow-up</p> <p>- Censored on 31st Dec 2015 = child was born on or after 1st Jan 2006 and:</p> <p>(i) is definitively known to be alive on 31st Dec 2015; or (ii) there is no information on death or lost to follow-up</p> <p>-Lost to follow up = child is lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)</p>	Numeric	<p>1 = Died</p> <p>2 = Alive at 10th birthday</p> <p>3 = Censored on 31st Dec 2015</p> <p>4 = Lost to follow up</p>

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
L_EXIT_DATE	<p>Date of last day in study (censored/lost/died/alive)</p> <ul style="list-style-type: none"> - Date lost - if child was lost to follow up - Date of death - if child has died - Date of birth plus 3652 days (approx. 10th birthday) if child was born on or before the 31st Dec 2005 and: <ul style="list-style-type: none"> (i) is definitively known to be alive on 10th birthday; or (ii) there is no information on death or lost to follow-up - Date of last day of the study (31st Dec 2015) if child was born on or after 1st Jan 2006 and: <ul style="list-style-type: none"> (i) is definitively known to be alive on 31st Dec 2015; or (ii) there is no information on death or lost to follow-up 	DDMonYYYY	
L_EXIT_DAYS	<p>Number of days child is in study</p> <p>This is calculated as the last date child was in the study (L_EXIT_DATE) minus the child's birth date (L_CH_DATE_B).</p>	Numeric (1-4 digits)	
L_CH_YEAR_D	Year of child's death as recorded on the death certificate	YYYY	
L_CH_AGED_H	<p>Age at death in complete hours for day 0 (first 24 hours) = applies to infants who died within the first 24 hours</p> <p>This variable can be used to check values recorded under variable L_CH_AGED_D.</p> <p>A child who died within 24 hours but across two dates is coded "0" on L_CH_AGED_D below.</p>	Numeric (1-2 digits)	<p>0 = Died <1 hour after birth</p> <p>1 = Died 1 complete hour after birth</p> <p>2 = Died 2 complete hours after birth</p> <p>Etc.</p> <p>23 = Died 23 complete hours after birth</p> <p>88 = Alive at 24 hours</p> <p>99 = Died within first 24 hours, but exact time unknown</p>

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
L_CH_AGED_D	Age at death <u>in complete days</u> (up to 10 th birthday). A calculated field using the difference between date of death and death of birth i.e. subtract child's date of birth from child's date of death.	Numeric (1-4 digits)	0 = died <24 hours after birth 1 = died 1 complete day after birth 2 = died 2 complete days after birth Etc. 8888 = Alive on 10th birthday or by end of study period, whichever is sooner 9999 = Died before 10th birthday, but exact time unknown
L_CAUSE_D_U	Underlying cause of death - diagnosis There can only be one underlying cause of death	String	ICD9 or ICD10 code
L_CAUSE_D_P	Primary/ immediate cause of death - diagnosis	String	ICD9 or ICD10 code
L_CAUSE_D_C	Contributing cause of death - diagnosis	String	ICD9 or ICD10 code
L_CAUSE_D_O1	Other causes of death- diagnosis 1	String	ICD9 or ICD10 code
L_CAUSE_D_O2	Other causes of death- diagnosis 2	String	ICD9 or ICD10 code
L_CAUSE_D_O3	Other causes of death- diagnosis 3	String	ICD9 or ICD10 code
L_CAUSE_D_O4	Other causes of death- diagnosis 4	String	ICD9 or ICD10 code
L_CAUSE_D_O5	Other causes of death- diagnosis 5 Additional "Other causes of death" can be recorded in variables L_CAUSE_D_O6 to L_CAUSE_D_15 (not listed)	String	ICD9 or ICD10 code
L_CH_PLACE_D	Place of death	Numeric	1 = Home 2 = Hospital 3 = Other 9 = Not known .=Not recorded by registry or not available for study

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
Variables relating to mother			
L_MAT_YEAR_B	Maternal year of birth	YYYY	99 = Not known . = Not recorded by registry or not available for study
L_MATAGE_B	Maternal age at infant's birth <u>in completed years</u> May be used to link information held on the child in local health care databases	Numeric	99=Not known . =Not recorded by registry or not available for study
L_MULT_BIRTH	Singleton or multiple birth	Numeric	1=Singleton 2=Twins 3=Triplets or higher 4= Multiple birth, number unknown 9= Not known . =Not recorded by registry or not available for study
L_MAT_CTRY_B	Maternal country of birth/ place of birth/ country of origin	Numeric	1= National 2= Other European 3 = Non-European 4= Non-national (exact nationality not specified) 9 = Not known . =Not recorded by registry or not available for study
L_MAT_BMI	Maternal Body Mass Index (BMI) at first antenatal visit/at booking Expected range 15 – 50	Numeric (Whole number only)	Exact BMI value 97 = <30 98 = >=30 99 = Not known . =Not recorded by registry or not available for study
L_MAT_EDUC	Maternal education	Numeric	1 = Pre-primary /Primary 2 = Any secondary 3 = Postsecondary (non-tertiary)

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
	(UNESCO's International Standard Classification of Education (ISCED))		4 = Tertiary 5=No education 9 = Not known . = Not recorded by registry or not available for study
L_MAT_OCC	Maternal occupation (Based on Paris (INSERM) coding classification)	Numeric	1= Farmer 2= Artisan (ex: baker) / shop-owner 3= Professional 4= Intermediate 5= Administrative/public service 6= Business employees: shop assistants/ salesperson 7= Household and personal service 8= Skilled manual worker 9=Unskilled manual worker 0= No occupation declared / student 99= Not known . =Not recorded by registry or not available for study
L_MATDEPR_IND	Deprivation index at maternal residence Multiple deprivations scores are ranked into quintiles where 1= Least deprived and 5 = Most deprived (coding scheme used in Wales & Basque Country)	Numeric	1 = First quintile (Least deprived) 2 = Second quintile 3 = Third quintile 4 = Fourth quintile 5 = Fifth quintile (Most deprived) 9 = Not known . =Not recorded by registry or not available for study
L_PROXY_SES	Proxy variable for SES This is registry-specific. Use the agreed proxy variable for each registry. Maternal education: <ul style="list-style-type: none"> • Tertiary/ post-secondary=High • Any secondary = Middle 	Numeric	1 = High 2 = Middle 3 = Low 9=Not known . = Not recorded by registry or not available for study

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
	<ul style="list-style-type: none"> Primary/ pre-primary/ No education = Low <p>Maternal occupation:</p> <ul style="list-style-type: none"> Professional = High Intermediate= Middle No occupation = Low <p>Multiple Deprivation Index</p> <ul style="list-style-type: none"> Quintile 1 (Least deprived)=High Quintiles 2-4= Middle Quintile 5 (Most deprived)= Low 		
L_MATMAR_STA	Maternal marital status	Numeric	1 = Single 2 = Married/ Living together 3 = Widow 4 = Divorced/ Separated 9 = Not known . =Not recorded by registry or not available for study
Recoded variables			
Yeargp	Grouped year of birth 1995/2004=1 2005/2014 =2	Numeric	1 = 1995-2004 2 = 2005-2014
BMI_gp	Grouped BMI BMI <10, code as unknown BMI >60, code as unknown Blank or missing, code as unknown	Numeric	1 = <30 2 = 30+ 9 = unknown
BW_gp	Grouped Birthweight BW <400g, code as unknown BW >7000g, code as unknown Blank or missing, code as unknown	Numeric	1= <1000g 2 = 1000-1499g 3 = 1500-2499g 4 = 2500-3999g 5 = 4000+ g 9 = unknown
GA_gp	Grouped Gestational age	Numeric	1 = 24-27weeks 2 = 28-31 weeks

Variable Name	Variable Definition and Instructions	Variable Format	Variable Values
	GA <24 weeks, excluded from study GA >45 weeks, code as unknown Blank or missing, code as unknown		3= 32-36 weeks 4 = 37+ weeks 9 = unknown
GA_disc_gp	Grouped Gestational age at discovery GA at discovery <8 weeks, code as unknown GA at discovery >42 weeks, code as unknown Blank or missing, code as unknown	Numeric	1= <22 weeks 2 = 22-31 weeks 3 = 32+ weeks 9 = unknown
matage_gp	Grouped Maternal age at infant's birth Maternal age range 10-19 years, code =1 20-34 years, code=2 35-59 years, code=3 All other values, blanks or missing, code=9	Numeric	1= <20 years 2= 20-34 years 3= 35+ years 9=Not known

Variables in EUROLINKCAT Common Data Model: Health Care

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
Table: Patient level data One row per child Core variables for linkage to hospital discharge records and for calculating age at event e.g. hospitalisation				
L_CH_ID	Unique identifier of child A unique ID that links child to another database	As recorded locally		- Patient - EUROCAC case data
L_CH_REC	Unique child record number Generated by syntax script for all children (cases and controls), irrespective of whether the child has been in hospital For children with a hospital admission, this is used to link admission, diagnoses, procedures & surgeries and prescriptions to child	Numeric		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
PROVIDER_ID	Unique reference number as present in the provider's linked data	String		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_CASECON	Case or control status (apply status AFTER linkage is complete) Exclusions: - child with gestational age <23 weeks Control with ICD10 Q code or ICD9 (-CM) 740-759 codes recorded. This is a child with a congenital anomaly code (ICD10 Q code, or ICD9-CM 740-759) in hospital discharge records that is not recorded as a EUROCAC case. Children coded as 1 or 2 will be included in WP4 analysis. Children coded as 3 are required for WP6	Numeric	1= Case 2= Control with no ICD10 Q code or ICD9(-CM) 740-759 codes recorded 3=Control with ICD10 Q code or ICD9(-CM) 740-759 codes recorded	- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_HOSP_ADM	Admitted to hospital (up to 10 th birthday) Exclude obstetric stays i.e. hospital stays relating to the birth and Accident & Emergency/ Emergency room stays	Binary	0= No hospital admissions 1= One or more hospital admission (s)	- Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
L_PRESCRIPT	Prescription for one of the 5 specified conditions: Asthma, Cardiac, Diabetes, Epilepsy, and Infections. N.B. UK prescriptions are <u>issued</u> by a GP, while in continental Europe, prescriptions are <u>dispensed</u> by the pharmacy (either community or hospital outpatient pharmacy)	Numeric	0= No prescription 1= One or more prescription (s)	- Patient
L_CH_DATE_B	Child's date of birth Used to calculate age at hospitalisation, diagnoses, procedures, & prescriptions	DDMonYYYY		- Patient
L_CH_YEAR_B	Child's year of birth	YYYY		- Patient
L_CH_DATE_D	Child's date of death	DDMonYYYY		- Patient
L_CH_YEAR_D	Child's year of death	YYYY		- Patient
L_CH_AGED_D	Age at death in complete days (up to 10th birthday). A calculated field using the difference between date of death and date of birth i.e. subtract child's date of birth from child's date of death.	Numeric (1-4 digits)	0 = died <24 hours after birth 1 = died 1 complete day after birth 2 = died 2 complete days after birth Etc. 8888 = Alive on 10th birthday or by end of study period, whichever is sooner 9999 = Died before 10th birthday, but exact time unknown	- Patient
Information on risk factors – needed for all children				
L_MATAGE_B	Maternal age at infant's birth <u>in completed years</u>	Numeric	99 = Not known . = Not recorded or not available for study	- Patient
L_MATMAR_STA	Maternal marital status at delivery	Numeric	1 = Single 2 = Married/ Living together 3 = Widow 4 = Divorced/ Separated 9 = Not known . = Not recorded or not available for study	- Patient
L_MAT_CTRY_B	Maternal country of birth/ place of birth/ country of origin	Numeric	1= National 2= Other European 3 = Non-European 4= Non-national (exact nationality not specified) 9 = Not known . = Not recorded or not available for study	- Patient
L_MAT_BMI	Maternal Body Mass Index (BMI) at first antenatal visit/at booking Expected range 15 – 50	Numeric (Whole number only)	Exact BMI value 97 = <30 98 = >=30 99 = Not known . = Not recorded or not available for study	- Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables¹
L_MAT_EDUC	Maternal education (UNESCO's International Standard Classification of Education (ISCED))	Numeric	1 = Pre-primary /Primary 2 = Any secondary 3 = Postsecondary (non tertiary) 4 = Tertiary 5=No education 9 = Not known . = Not recorded or not available for study	- Patient
L_MATDEPR_IND	Deprivation index at maternal residence Multiple deprivations scores are ranked into quintiles where 1= Least deprived and 5 = Most deprived (coding scheme used in Wales & Basque Country)	Numeric	1 = First quintile (Least deprived) 2 = Second quintile 3 = Third quintile 4 = Fourth quintile 5 = Fifth quintile (Most deprived) 9 = Not known . = Not recorded or not available for study	- Patient
L_PROXY_SES	Proxy variable for Socio-Economic Status (SES) This is registry-specific. Use the agreed proxy variable for each registry. <u>Maternal education:</u> <ul style="list-style-type: none"> • Tertiary/ post-secondary=High • Any secondary = Middle • Primary/ pre-primary/ No education = Low <u>Maternal occupation:</u> <ul style="list-style-type: none"> • Professional = High • Intermediate= Middle • No occupation = Low <u>Multiple Deprivation Index</u> <ul style="list-style-type: none"> • Quintile 1 (Least deprived) =High • Quintiles 2-4= Middle • Quintile 5 (Most deprived) = Low 	Numeric	1 = High 2 = Middle 3 = Low 9=Not known . = Not recorded or not available for study	- Patient
L_MULT_BIRTH	Singleton or multiple birth	Numeric	1=Singleton 2=Twins 3=Triplets or higher 4= Multiple birth, number unknown 9= Not known . = Not recorded or not available for study	- Patient
L_CH_SEX	Child's sex	Numeric	1 = Male 2 = Female 3 = Indeterminate 9 = Not known . = Not recorded or not available for study	- Patient
L_CH_BW	Child's birth weight (grams)	Numeric	9999 = Not known . = Not recorded or not available for study	- Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
L_CH_GA_B	Child's gestational age at birth (in completed weeks).	Numeric	99 = Not known .= Not recorded or not available for study	- Patient
L_PRENATAL	Prenatal diagnosis of congenital anomaly [EUROCAT case only, i.e. L_CASECON=1 All control children in the linked dataset (i.e. L_CASECON=2 or L_CASECON=3) should be coded as .] [Derived from the EUROCAT variables when discovered (WHENDISC) and age at prenatal diagnosis (AGEDISC)]	Numeric	0 = Not prenatally diagnosed 1 = Prenatally diagnosed, <22 weeks 2 = Prenatally diagnosed, 22-31 weeks 3 = Prenatally diagnosed, 32+ weeks 4 = Prenatally diagnosed, GA unknown 9 = Not known if prenatally diagnosed .= Not recorded or not available for study	-Patient
Recoded variables				
Yeargp_WP4	Grouped year of birth 1995/2004=1 2005/2009 =2 2010-2014 = 3	Numeric	1 = 1995-2004 2 = 2005-2009 3 = 2010 - 2014	-Patient
Yeargp_Px	Grouped year of birth for the prescription study 2000/2004=1 2005/2009 =2 2010-2014 = 3	Numeric	1 = 2000-2004 2 = 2005-2009 3 = 2010 - 2014	-Patient
BMI_gp	Grouped BMI BMI <10, code as unknown BMI >60, code as unknown Blank or missing, code as unknown	Numeric	1 = <30 2 = 30+ 9 = unknown	-Patient
BW_WP4_gp	Grouped Birth weight (BW) BW <400g, code as unknown BW >7000g, code as unknown Blank or missing, code as unknown	Numeric	1= <1500g 2 = 1500-2499g 3 = 2500-3999g 4 = 4000+ g 9 = unknown	-Patient
GA_WP4_gp	Grouped Gestational age GA <23 weeks, excluded from study GA >44 weeks, code as unknown Blank or missing, code as unknown	Numeric	1 = 23-27 weeks 2 = 28-31 weeks 3 = 32-36 weeks 4 = 37+ weeks 9 = unknown	-Patient
GA_disc_gp	Grouped Gestational age at discovery GA at discovery <8 weeks, code as unknown GA at discovery >42 weeks, code as unknown Blank or missing, code as unknown	Numeric	1= <22 weeks 2 = 22-31 weeks 3 = 32+ weeks 9 = unknown	-Patient
Matage_gp	Grouped Maternal age at infant's birth Maternal age range 10-19 years, code =1 20-34 years, code=2 35-59 years, code=3 All other values, blanks or missing, code=9	Numeric	1= <20 years 2= 20-34 years 3= 35+ years 9=Not known	-Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
Derived variables relating to linkage				
L_MATCH_TYPE_V	Match with national/ vital statistics database	Numeric	1 = Linkage to national/vital statistics database - match 2 = Linkage to national/vital statistics database - non-match 3 = EUROCAT death only . = No linkage to national/ vital statistics	-Patient
L_MATCH_TYPE_H	<p>Match with hospital database</p> <p>The value 4 "Only matched to hospital discharge database outside the study period or matched to hospital outpatient records in the study period" is included for matching purposes only i.e. the hospital stay for this child is not included in analysis.</p> <p>The rationale is that if a child is not matched to hospital discharge records during the study period, but is found in hospital records AFTER the study period, then we can be more confident that the child did not have a hospital admission during our study period.</p> <p>Similarly, if a child is found in outpatient records during the study period but is not found in the hospital discharge records, we can be more confident that the child did not have a hospital admission during our study period.</p>	Numeric	1 = Linkage to hospital database - match 2 = Linkage to hospital database - non-match 3 = EUROCAT death only 4 = Only matched to hospital discharge database outside the study period or matched to hospital outpatient records in the study period	-Patient
L_CONFID_HDR	<p>Strength of match with hospital database.</p> <p>Use local data provider's codes for assessing confidence that the case is correctly matched. If local code unavailable, use suggested coding</p> <p>The value 0 represents a child:</p> <ul style="list-style-type: none"> not matched to hospital discharge records in the study period, but found in other databases within national statistics (such as outpatient records) found only in hospital discharge records AFTER the study period. 	Numeric	0=Found in other database or found only in hospital database outside the study period 1=Excellent 2=Good 3=Fair 4=Poor 9=Not Matched	- Patient
L_MORB_INCLUDE	<p>Include child in WP4 analysis</p> <p>Used in sensitivity analysis to assess if a child should be included in the WP4 analysis or not i.e. it will assess confidence that a child definitely did not go to hospital if hospital admission (variable L_HOSP_ADM = 0).</p> <p>Please see appendix 1 for algorithm)</p>	Binary	0 = Do not include in analysis 1 = Include in analysis	- Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
L_MATCH_PX	Match with GP / community pharmacy prescription database / hospital outpatient pharmacy records	Numeric	1 = Linked prescription database – match 2 = Linked to prescription database – non-match 3 = Linkage to prescription data not available	- Patient
L_CONFID_PX	Strength of match with prescription database. The value 9 represents a child not matched/ found in the local GP/ pharmacy prescription records (i.e. L_MATCH_PX=2); or a child whose GP/ pharmacy do not contribute data to this study (i.e. L_MATCH_PX=3). Children coded as L_MATCH_PX=3 will be excluded from analysis.	Numeric	1=Excellent 2=Good 3=Fair 4=Poor 9=Not Matched	- Patient
L_DATE_LOST	Date lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)	DDMonYYYY	. = Not recorded or not available for study	- Patient
L_YEAR_LOST	Year lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)	YYYY	. = Not recorded or not available for study	- Patient
L_AGE_L_D	Age lost to follow-up/ linkage <u>in complete days</u>	Numeric (1-4 digits)	. = Not recorded or not available for study	- Patient
L_EXIT_DATE	Date of last day in study (censored/lost/died/alive) - Date lost - if child was lost to follow up - Date of death - if child has died - Date of birth plus 3652 days (approx. 10th birthday) if child was born on or before the 31st Dec 2005 and: (i) is definitively known to be alive on 10th birthday; or (ii) there is no information on death or lost to follow-up - Date of last day of the study (31st Dec 2015) if child was born on or after 1st Jan 2006 and: (i) is definitively known to be alive on 31st Dec 2015; or (ii) there is no information on death or lost to follow-up	DDMonYYYY		- Patient
L_EXIT_DAYS	Number of days child is in study. This is calculated as the last date child was in the study (L_EXIT_DATE) minus the child's birth date (L_CH_DATE_B).	Numeric (1-4 digits)		- Patient
L_CH_STATUS	Outcome status -Died = child is known to have died before 10th birthday or 31 Dec 2015 (whichever earlier)	Numeric	1 = Died 2 = Alive at 10 th birthday 3 = Censored on 31 st Dec 2015 4 = Lost to follow up	- Patient

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
	<p>-Alive at 10th birthday = child was born on or before the 31st Dec 2005 and: (i) is definitively known to be alive on 10th birthday; or (ii) there is no information on death or lost to follow-up</p> <p>- Censored on 31st Dec 2015 = child was born on or after 1st Jan 2006 and: (i) is definitively known to be alive on 31st Dec 2015; or (ii) there is no information on death or lost to follow-up</p> <p>-Lost to follow up = child is lost to follow-up/ linkage (i.e. due to emigration, adoption or other reason)</p>			
Table: Hospital admission variables (excluding Obstetric stays and Accident & Emergency/ Emergency room stays) One row for each hospital admission. If multiple hospital admissions, please complete all hospital admission variables for each admission				
L_CH_REC	<p>Unique child record number</p> <p>Generated by syntax script for all children (cases and controls), irrespective of whether the child has been in hospital</p> <p>For children with a hospital admission, this is used to link admission, diagnoses, procedures & surgeries and prescriptions to child</p>	Numeric		<ul style="list-style-type: none"> - Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
PROVIDER_ID	<p>Unique reference number as present in the provider's linked data</p>	String		<ul style="list-style-type: none"> - Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_ADM_REC	<p>Unique admission record number</p> <p>Used to link the child's admission records to the child's diagnoses and procedures & surgeries during that admission</p> <p>Generated by syntax script</p>	Numeric		<ul style="list-style-type: none"> - Hospital admission - - Diagnoses - Procedures & surgeries
L_CASECON	<p>Case or control status (apply status AFTER linkage is complete)</p> <p>Exclusions: - child with gestational age <23 weeks</p> <p>Control with ICD10 Q code or ICD9 (-CM) 740-759 codes recorded. This is a child with a congenital anomaly code (ICD10 Q code, or ICD9-CM 740-759) in hospital discharge records that is not recorded as a EUROCAT case.</p> <p>Children coded as 1 or 2 will be included in WP4 analysis. Children coded as 3 are required for WP6</p>	Numeric	<p>1= Case</p> <p>2= Control with no ICD10 Q code or ICD9(-CM) 740-759 codes recorded</p> <p>3=Control with ICD10 Q code or ICD9(-CM) 740-759 codes recorded</p>	<ul style="list-style-type: none"> - Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
L_HOSP_SEQ	<p>This variable is only required for the registries: registry numbers 13, 28</p> <p>Sequential number allocated to each hospital in registry area e.g. Hospital A=1, Hospital B=2 etc.</p> <p>Purpose of this variable is to identify the hospital for each record. It is known that some hospitals in some registries are missing >20% hospital admissions in specific years. This variable will be used in sensitivity analysis, to include/ exclude the hospital (s) with >20% missing information to see its effect on the analysis.</p>	Numeric	. = Not recorded or not available for study	- Hospital admission
L_DATE_ADM	Date of admission to hospital	DDMonYYYY		- Hospital admission
L_YEAR_ADM	Year of admission to hospital	YYYY		- Hospital admission
L_DATE_DIS	Date of discharge from hospital	DDMonYYYY		- Hospital admission
L_YEAR_DIS	Year of discharge from hospital	YYYY		- Hospital admission
L_HOSP_DAYS	<p>Length of stay in hospital in days</p> <p>Subtract discharge date from hospital admission date.</p> <p>If value =0 for children admitted and discharged on the same day, replace value 0 with value 0.5</p>	Numeric	<p>0.5 = <1 day</p> <p>1=1 day/ over-night stay</p> <p>2=2 days etc</p> <p>999= Not known</p>	- Hospital admission
L_AGE_ADM	<p>Child's age at hospital admission <u>in days</u> (up to 10th birthday).</p> <p>Subtract child's date of birth from date of admission to hospital</p>	Numeric	<p>0= <1 day old</p> <p>1= 1 day old</p> <p>2= 2 days old etc</p> <p>9999= exact age not known</p>	- Hospital admission
L_ICU_ADM	Admission to Neonatal, Paediatric or other Intensive Care Unit (ICU) during this hospital admission	Numeric	<p>0= No ICU admission</p> <p>1=ICU admission</p> <p>9 = Not known</p> <p>. = Not recorded or not available for study</p>	- Hospital admission
L_VENT	<p>Mechanical ventilator use during this hospital admission</p> <p>Non-invasive ventilation is excluded</p>	Numeric	<p>0= No</p> <p>1= Yes</p> <p>9= Not known</p> <p>. = Not recorded or not available for study</p>	- Hospital admission
<p>Table: Diagnosis variables (up to 10th birthday)</p> <p>Each diagnosis is a separate observation (row) in the table</p> <p><u>0, 1 or more rows for each hospital admission</u></p> <p><i>If multiple diagnoses, please provide date of each diagnosis</i></p>				
L_CH_REC	Unique child record number	Numeric		- Patient - Hospital admission - Diagnoses

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
	Generated by syntax script for all children (cases and controls), irrespective of whether the child has been in hospital For children with a hospital admission, this is used to link admission, diagnoses, procedures & surgeries and prescriptions to child			- Procedures & surgeries - Prescription
PROVIDER_ID	Unique reference number as present in the provider's linked data	String		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_ADM_REC	Unique admission record number Used to link the child's admission records to the child's diagnoses and procedures during that admission Generated by syntax script	Numeric		- Hospital admission - Diagnoses - Procedures & surgeries
L_CASECON	Case or control status (apply status AFTER linkage is complete) Exclusions: - child with gestational age <23 weeks Control with ICD10 Q code or ICD9 (-CM) 740-759 codes recorded. This is a child with a congenital anomaly code (ICD10 Q code, or ICD9-CM 740-759) in hospital discharge records that is not recorded as a EUROCAT case. Children coded as 1 or 2 will be included in WP4 analysis. Children coded as 3 are required for WP6	Numeric	1= Case 2= Control with no ICD10 Q code or ICD9(-CM) 740-759 codes recorded 3=Control with ICD10 Q code or ICD9(-CM) 740-759 codes recorded	- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_DIAG	Diagnosis in ICD9 or ICD10 for the hospital stay	String As recorded in the hospital database		- Diagnoses
L_DATE_DIAG	Date of diagnosis Use date of discharge as a proxy for date of diagnosis, if date of diagnosis is not available	DDMonYYYY		- Diagnoses
L_AGE_DIAG	Child's age at diagnosis in <u>days</u> (up to 10 th birthday). Subtract child's date of birth from date of diagnosis to calculate the child's age at diagnosis. If date of diagnosis is not available, then use date of discharge	Numeric	0= <1 day old 1= 1 day old 2= 2 days old etc 9999= exact age not known	- Diagnoses

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
L_CANCER	A diagnosis of cancer (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_CPALSY	A diagnosis of cerebral palsy (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_EPILEPSY	A diagnosis of epilepsy or seizures (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_RENAL	A diagnosis of renal failure (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_HEARING	A diagnosis of hearing loss (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_VISION	A diagnosis of visual impairment or blindness (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_INJURYPOISON	A diagnosis of injury or poisoning (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
L_BATTERED	A diagnosis of physical abuse (battered child) (ICD9 or ICD10)	Binary	0 = No 1 = Yes	- Diagnoses
<p>Table: Surgery and Procedures (including admission to intensive care unit) Each surgery/ procedure /ICU admission is a separate observation (row) in the table 0,1 or more rows for each hospital admission If multiple procedures, please provide procedure date and age of child at each procedure</p>				
L_CH_REC	Unique child record number Generated by syntax script for all children (cases and controls), irrespective of whether the child has been in hospital For children with a hospital admission, this is used to link admission, diagnoses, procedures & surgeries and prescriptions to child	Numeric		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
PROVIDER_ID	Unique reference number as present in the provider's linked data	String		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_ADM_REC	Unique admission record number Used to link the child's admission records to the child's diagnoses and procedures during that admission Generated by syntax script			- Hospital admission - Diagnoses - Procedures & surgeries
L_CASECON	Case or control status (apply status AFTER linkage is complete) Exclusions: - child with gestational age <23 weeks Control with ICD10 Q code or ICD9 (-CM) 740-759 codes recorded. This is a child with a congenital anomaly code (ICD10 Q code, or ICD9-CM 740-759) in hospital	Numeric	1= Case 2= Control with no ICD10 Q code or ICD9(-CM) 740-759 codes recorded 3=Control with ICD10 Q code or ICD9(-CM) 740-759 codes recorded	- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription

Variable name	Variable definition/ description	Variable format	Variable values	Tables¹
	discharge records that is not recorded as a EUROCAT case. Children coded as 1 or 2 will be included in WP4 analysis. Children coded as 3 are required for WP6			
L_SURG_DATE	Date of surgery / procedure If date of surgery is missing, use the date of ICU admission as a proxy. If date of ICU is also missing, use date of discharge	DDMonYYYY		- Procedures & surgeries
L_SURG_YEAR	Year of surgery / procedure	YYYY		- Procedures & surgeries
L_SURG_AGE	Child's age at surgery <u>in days</u> (up to 10 th birthday). Subtract child's date of birth from date of surgery / procedure	Numeric	0= <1 day old 1= 1 day old 2= 2 days old etc 9999=surgery, but exact age not known	- Procedures & surgeries
L_SURG_CODE	Code for surgery / procedure performed during this hospital stay As recorded in the hospital database	String		- Procedures & surgeries
SU_ANY	Any Surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_FEEDING	Surgical Feeding Tube inserted As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_SB	Surgery for spina bifida As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_REVSHUNT	Revisions of shunt surgery – relevant to both spina bifida and hydrocephaly As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_EYE	Eye surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_CARDIAC	Any Cardiac surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_RASHKIND	Rashkind procedure As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TGA_ARTERIAL	Arterial switch As recorded in the hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TGA_ATRIAL	Atrial switch As recorded in the hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_VSD	Surgery for ventricular septal defect As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_ASD	Surgery for atrial septal defect As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries

Variable name	Variable definition/ description	Variable format	Variable values	Tables¹
SU_AVSD	Surgery for atrioventricular septal defect As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_COARCT	Surgery for coarctation of aorta As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_PDA	Surgery for Patent ductus arteriosus (PDA) or catheter closure As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_BLALOCK	Blalock shunt As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_HEMIFON	Hemi-Fontan Surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_FONTAN	Complete Fontan surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_CLEFT	Cleft surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_OESOPH	Surgery for oesophageal atresia As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_DILATION	Oesophageal atresia dilation As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_INTESTINE	Intestinal surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_STOMY	Stoma Surgery As recorded in the hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_KASAI	Liver/bile surgery (Kasai) As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_CDH	Surgery for diaphragmatic hernia As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_GASTRO	Surgery for gastroschisis As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_RENAL	Renal surgery As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_HYPOSPA	Surgery for hypospadias As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TRANY	Transplantation (heart, kidney or liver) As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TRHEART	Transplantation (heart) As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TRKIDNEY	Transplantation (kidney) As recorded in hospital database	Binary	0 = No 1 = Yes	- Procedures & surgeries
SU_TRLIVER	Transplantation (liver)	Binary	0 = No	- Procedures & surgeries

Variable name	Variable definition/ description	Variable format	Variable values	Tables ¹
	As recorded in hospital database		1 = Yes	
L_ICU_DATE_ADM	Date of admission to ICU	DDMonYYYY	. = Not recorded or not available for study	- Procedures & surgeries
L_ICU_DATE_DIS	Date moved from ICU to other ward	DDMonYYYY	. = Not recorded or not available for study	- Procedures & surgeries
L_ICU_YEAR	Year of admission to ICU	YYYY	. = Not recorded or not available for study	- Procedures & surgeries
L_ICU_AGE	Child's age at admission to ICU <u>in days</u> (up to 10 th birthday). Subtract child's date of birth from date of ICU admission	Numeric	0= <1 day old 1= 1 day old 2= 2 days old etc 9999= exact age not known . = Not recorded by registry or not available for study	- Procedures & surgeries
Table: Prescriptions (for 5 conditions only: Asthma, Cardiac, Diabetes, Epilepsy and Infections) <u>One row for each prescription.</u> <i>If multiple prescriptions, please provide prescription date and age of child at each prescription</i>				
L_CH_REC	Unique child record number Generated by syntax script for all children (cases and controls), irrespective of whether the child has been in hospital For children with a hospital admission, this is used to link admission, diagnoses, procedures & surgeries and prescriptions to child	Numeric		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescriptions
PROVIDER_ID	Unique reference number as present in the provider's linked data	String		- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_CASECON	Case or control status (apply status AFTER linkage is complete) Exclusions: - child with gestational age <23 weeks Control with ICD10 Q code or ICD9 (-CM) 740-759 codes recorded. This is a child with a congenital anomaly code (ICD10 Q code, or ICD9-CM 740-759) in hospital discharge records that is not recorded as a EUROCAT case. Children coded as 1 or 2 will be included in WP4 analysis. Children coded as 3 are required for WP6	Numeric	1= Case 2= Control with no ICD10 Q code or ICD9(-CM) 740-759 codes recorded 3=Control with ICD10 Q code or ICD9(-CM) 740-759 codes recorded	- Patient - Hospital admission - Diagnoses - Procedures & surgeries - Prescription
L_AGE_PX	Child's age at prescription <u>in complete days</u> (up to 10 th birthday). Subtract child's date of birth from the date prescription was issued/ dispensed	Numeric	0= <1 day old 1= 1 day old 2= 2 days old etc 9999= exact age not known . = Not recorded or not available for study	- Prescriptions

Variable name	Variable definition/ description	Variable format	Variable values	Tables¹
L_DATE_PX	Date prescription was issued/ dispensed	DDMonYYYY	. = Not recorded or not available for study	- Prescriptions
L_DRUG_CODE	Codes for medication: Anatomical Therapeutic Chemical classification (ATC), or Read codes	As recorded in local prescription database		- Prescriptions
DIA_ANY	Any prescribed antidiabetic medication (ATC codes beginning A10)	Binary	0 = No 1 = Yes	- Prescriptions
DIA_INS	Any prescribed Insulin or insulin analogues (ATC codes beginning A10A)	Binary	0 = No 1 = Yes	- Prescriptions
DIA_GLU_LOW	Any prescribed blood glucose lowering drugs, excluding insulins (ATC codes beginning A10B)	Binary	0 = No 1 = Yes	- Prescriptions
INF_ANY	Any prescribed anti-infective medication (ATC codes beginning J01-J05)	Binary	0 = No 1 = Yes	- Prescriptions
INF_ANTIBAC_SYS	Any prescribed antibacterials for systemic use (ATC codes beginning J01)	Binary	0 = No 1 = Yes	- Prescriptions
INF_PEN	Any prescribed beta-lactam antibacterials, penicillins (ATC codes beginning J01C)	Binary	0 = No 1 = Yes	- Prescriptions
INF_MAC	Any prescribed macrolides, lincosamides and streptogramins (ATC codes beginning J01F)	Binary	0 = No 1 = Yes	- Prescriptions
INF_ANTIVIR_SYS	Any prescribed antivirals for systemic use (ATC codes beginning J05)	Binary	0 = No 1 = Yes	- Prescriptions
EPI_ANY	Any prescribed antiepileptic drugs (AEDs) (ATC codes beginning N03)	Binary	0 = No 1 = Yes	- Prescriptions
EPI_OLD	Any prescribed older AEDs (ATC codes beginning N03AA phenobarbital, N03AB phenytoin, N03AE clonazepam, N03AF carbamazepine)	Binary	0 = No 1 = Yes	- Prescriptions
EPI_VPA	Any prescribed fatty acid derivatives (ATC codes beginning N03AG, valproic acid)	Binary	0 = No 1 = Yes	- Prescriptions
EPI_NEW	Any prescribed newer AEDs (ATC codes N03AX lamotrigine, topiramate, gabapentin, levetiracetam)	Binary	0 = No 1 = Yes	- Prescriptions
AST_ANY	Any prescribed antiasthmatic medications (ATC codes beginning R03)	Binary	0 = No 1 = Yes	- Prescriptions

Variable name	Variable definition/ description	Variable format	Variable values	Tables¹
AST_IN_B2	Any prescribed Inhaled β 2- agonists (ATC codes beginning R03AC)	Binary	0 = No 1 = Yes	- Prescriptions
AST_IN_CORT	Any prescribed Inhaled corticosteroids (ATC codes beginning R03BA)	Binary	0 = No 1 = Yes	- Prescriptions
AST_ANTICHO	Any prescribed Anticholinergic inhaled medications (ATC codes beginning R03BB)	Binary	0 = No 1 = Yes	- Prescriptions
AST_ANTIALL	Any prescribed Antiallergic agents, excl. corticosteroids (ATC codes beginning R03BC)	Binary	0 = No 1 = Yes	- Prescriptions
AST_LEU	Any prescribed Leukotriene receptor antagonists (ATC codes beginning R03DC)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_ANY	Any prescribed cardiac medication, excluding quinidine and adrenalin pen (ATC codes beginning C01-C03, C07-C09, excluding C01BA51, C01BA71, C01CA24)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_VWC1	Any prescribed sodium channel blockers Vaughan Williams class (VWC) 1 (ATC codes beginning C01BA, C01BB and C01BC, excluding C01BA51, C01BA71 quinidine)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_VWC2	Any prescribed betablockers VWC 2 (ATC codes beginning C07A)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_VWC3	Any prescribed VWC 3 medications (amiodarone) (ATC codes beginning C01BD)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_VWC5	Any prescribed VWC 5 medications (digoxine) (ATC codes beginning C01AA05)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_HYP	Any prescribed medications for hypertension (calcium antagonists and agents acting on renin-angiotensin system) (ATC codes beginning C08, C09)	Binary	0 = No 1 = Yes	- Prescriptions
CAR_DIU	Any prescribed diuretics (ATC codes beginning C03)	Binary	0 = No 1 = Yes	- Prescriptions
L_GP_HOSP	Indicates if community pharmacy or hospital outpatient pharmacy prescription (for Italian registries only)	Numeric	1= Community pharmacy prescription 2= Hospital outpatient pharmacy prescription 3= Not available (for non-Italian registries)	- Prescriptions