






BMJ Open Prevalence of skin pressure injury in critical care patients in the UK: results of a single-day point prevalence evaluation in adult critically ill patients

Francesca Rubulotta ^{1,2,3}, Stephen Brett ³, Carole Boulanger,⁴ Bronagh Blackwood ⁵, Mieke Deschepper,⁶ Sonia O Labeau ⁷, Stijn Blot ⁸, UK Collaborating Site Investigators, On behalf of the DecubICUs study team and the European Society of Intensive Care Medicine Trials' Group UK Collaborators

To cite: Rubulotta F, Brett S, Boulanger C, *et al.* Prevalence of skin pressure injury in critical care patients in the UK: results of a single-day point prevalence evaluation in adult critically ill patients. *BMJ Open* 2022;**12**:e057010. doi:10.1136/bmjopen-2021-057010

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057010>).

FR and SB are joint first authors.

Received 25 September 2021
Accepted 30 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Francesca Rubulotta;
frubulot@ic.ac.uk

ABSTRACT

Objectives Hospital-acquired pressure injuries (PIs) are a source of morbidity and mortality, and many are potentially preventable.

Design This study prospectively evaluated the prevalence and the associated factors of PIs in adult critical care patients admitted to intensive care units (ICU) in the UK.

Setting This service evaluation was part of a larger, international, single-day point prevalence study of PIs in adult ICU patients. Training was provided to healthcare givers using an electronic platform to ensure standardised recognition and staging of PIs across all sites.

Participants The characteristics of the ICUs were recorded before the survey; deidentified patient data were collected using a case report form and uploaded onto a secure online platform.

Primary and secondary outcome measures Factors associated with ICU-acquired PIs in the UK were analysed descriptively and using mixed multiple logistic regression analysis.

Results Data from 1312 adult patients admitted to 94 UK ICUs were collected. The proportion of individuals with at least one PI was 16% (211 out of 1312 patients), of whom 8.8% (n=115/1312) acquired one or more PIs in the ICU and 7.3% (n=96/1312) prior to ICU admission. The total number of PIs was 311, of which 148 (47.6%) were acquired in the ICU. The location of majority of these PIs was the sacral area, followed by the heels. Braden score and prior length of ICU stay were associated with PI development.

Conclusions The prevalence and the stage of severity of PIs were generally low in adult critically ill patients admitted to participating UK ICUs during the study period. However, PIs are a problem in an important minority of patients. Lower Braden score and longer length of ICU stay were associated with the development of injuries; most ICUs assess risk using tools which do not account for this.

Trial registration number NCT03270345.

INTRODUCTION

In 2014, the UK National Institute for Health and Care Excellence (NICE) published a document for prevention and management of pressure injuries (PIs) in primary and

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is the first study ever completed to evaluate the impact of pressure injuries (PIs) in the adult population.
- ⇒ It is a robustly planned and executed study.
- ⇒ Limitations are mainly due to the low incidence and severity of PIs.
- ⇒ The inability to link preventive measures used in patients to the prevalence of PIs is also a limitation.
- ⇒ The study was conducted in the pre-COVID-19 era and majority of the patients were nursed in the supine position.

secondary care in the UK.^{1 2} This document was revised in 2019 and states that 'pressure ulcers are serious and distressing adverse events that can represent a failure of care'.³ PIs result when an area of the skin and/or the underlying tissues are damaged due to being placed under sufficient pressure for a period long enough to impair blood supply or when shearing forces generate friction on the skin during manual handling.^{1 2} An international classification categorises the injuries into stages 1–4, unstageable and suspected deep tissue injury according to the extent of tissue damage.^{3 4}

Pressure ulcers caused by both moisture and shear stress are directly associated with the quality of care provided as well as patient outcomes.^{1–4} NICE guidelines² state that PIs consume significant resources and that these should be preventable in the National Health Service (NHS), which has therefore employed tissue viability nurses (TVN) in the majority of hospitals. TVNs provide expert advice in the prevention and treatment of wounds including PIs.

Unfortunately, critically ill patients in intensive care units (ICUs) are extremely vulnerable due to the severity of their illness, immobility, sedation, poor tissue perfusion, hypoxia and frequent haemodynamic instability. The current prevalence of PIs in the UK ranges widely, with estimates from 4.7% to 32.1% in ward areas, while in the ICUs the prevalence is not known.^{2 5 6} The aim of this service evaluation was to provide up-to-date information on PIs within UK adult ICUs. The specific objectives were (1) to provide a picture of the prevalence of PIs in adult critically ill patients, specifically in relation to the presence of PIs on admission and PIs acquired within the ICU; (2) to provide the characteristics of PIs in terms of severity and anatomical distribution; and (3) to contribute data to the international Decubitus in Intensive Care Units (DecubICUs) study, of which this project formed a part.¹

METHODS

Study design and participants

This service evaluation was conducted using a single-day (midnight to midnight) point prevalence survey of all adult ICU patients. A full description of the methods can also be found in the DecubICUs global study report.¹ The study was registered at ClinicalTrials.gov (NCT03270345).

Patient and public involvement

There was no patient and public involvement in this study.

Data collection

The study website provides the protocol and an electronic case report form (CRF; online supplemental appendix A and <https://www.esicm.org/trials-group-2/decubicus/>). Data were collected from all participating centres before the study day. Anonymous data were collected in the CRF on all adult patients present in the participating ICUs on 15 May 2018 (online supplemental appendix A). Admission data included patients' demographic, type of ICU admission (ie, medical, elective or emergency surgical, or trauma/burns), principal diagnosis leading to ICU admission, mechanical ventilation on admission and whether the patient already had PIs at the time of ICU admission. Data included PI assessment, which included site and stage (generally referred to as 'grade' in the UK),³⁻⁶ and whether injuries were present at ICU admission (specifically these were areas exhibiting PIs, rather than discrete injuries; referred to as PIs for simplicity). Injury risk was evaluated using the Braden Scale.⁷ This scale combines six subscales (mobility, activity, sensory perception, skin moisture, nutritional state and friction/shear) and scores range from 6 to 23, with lower scores reflecting higher risk. Follow-up data gathered were survival status and length of ICU and hospital stay until hospital discharge, maximally at 12 weeks following the study day (7 August 2018). An online training module was developed for all clinical data collectors and was published on the study website (protocol available in online supplemental appendix A or online at <https://www.esicm.org/trials-group-2/decubicus/>;

training module available online on the European Society of Intensive Care Medicine website at <https://www.esicm.org/wp-content/uploads/2018/04/Module-DecubICUs-LR.pdf>) prior to initiation in order to assist with consistency of data collection. Hospital and follow-up data were collected by clinical and research nurses and the site study coordinator. Individual patient data were collected by the bedside nurse by direct skin observation according to international staging definitions.³⁻⁶

Statistics

Data were summarised as mean (SD) or median (IQR) as appropriate; categorical data were summarised as proportions (%). Overall PI prevalence was calculated as the proportion of the sample who had at least one PI on the study day. ICU-acquired prevalence was calculated as the proportion of the sample who had at least one PI determined to be acquired in the ICU present on the study day. Prevalence is reported as percentage with 95% CI.

Associations with ICU-acquired PI were explored using a mixed multiple logistic regression analysis with the logit link function and including a random intercept for site.

This method was chosen to balance potential effects resulting from variability in care processes across the participating sites. All demographic variables as well as those related to acute illness and chronic conditions were included. The variable 'length of ICU stay before study day' was included based on both clinical judgement and the literature on PI risk factors.⁸⁻¹⁰ As such, all variables were included following an exploratory approach, irrespective of their relationship with PI in univariate analysis.¹¹ Results are reported as OR with 95% CI. Statistical analysis was performed using IBM SPSS for Windows V.26.0 and R statistical software V.3.6.1.¹²

RESULTS

Ninety-four adult ICUs distributed across the whole of the UK participated in this study. In 2018, the Scottish Intensive Care Society Audit Group collected data from 72 adult ICUs,¹³ while the Intensive Care National Audit and Research Centre report was based on data from 263 NHS adult ICUs in the rest of the UK.¹⁴ The definition of an ICU can be broad; however, for the purpose of this study, the authors assumed that 28% (94 out of 335) of ICUs participated in this study in the UK; 51% declared themselves to be university-affiliated. The majority of ICUs were mixed medical and surgical (75; 80%), and the remainder were surgical of one type or another, with one specialist burns unit. Four participating units did not submit descriptive data. The ICUs had a median of 14 beds (IQR 10–20) and 68% declared themselves to be 'closed'. There were six very large units with more than 40 beds.

Local investigators collected data from 1312 critically ill patients, the majority of whom were cared for in university-affiliated centres (67.3% of patients). The median nurse

Table 1 Characteristics of patients

Characteristics	All patients (N=1312, 100%)*	Patients without pressure injuries (n=1101, 83.9%)*	Patients with ICU-acquired pressure injuries and/or pressure injuries acquired prior to ICU admission (n=211, 16%)*	Patients with ICU-acquired pressure injuries only (n=115, 8.8%)*
Age (years), M (IQR)	62 (50–72)	62 (49–72)	65 (54–74)	62 (52–73)
Sex (male), n (%)	783 (59.7)	656 (59.6)	127 (60.2)	68 (59.1)
Body mass index class, n (%)				
Underweight (<18.5 kg/m ²)	57 (4.3)	42 (3.8)	15 (7.1)	10 (8.7)
Normal weight (18.5–24.9 kg/m ²)	464 (35.4)	381 (34.6)	83 (39.3)	49 (42.6)
Preobesity (25–29.9 kg/m ²)	391 (29.8)	341 (31.0)	50 (23.7)	27 (23.5)
Obesity class I (30–34.9 kg/m ²)	227 (17.3)	195 (17.7)	32 (15.2)	12 (10.4)
Obesity class II (35–39.9 kg/m ²)	90 (6.9)	75 (6.8)	15 (7.1)	11 (9.6)
Obesity class III (≥40 kg/m ²)	83 (6.3)	67 (6.1)	16 (7.6)	6 (5.2)
Mechanical ventilation on ICU admission, n (%)	669 (51.0)	555 (50.4)	114 (54.0)	77 (67.0)
Type of admission, n (%)				
Medical	617 (47.0)	495 (45.0)	122 (57.8)	63 (54.8)
Elective surgery	279 (21.3)	260 (23.6)	19 (9.0)	11 (9.6)
Emergency surgery	309 (23.6)	255 (23.2)	54 (25.6)	31 (27.0)
Trauma and burns	107 (8.2)	91 (8.3)	16 (7.6)	10 (8.7)
Comorbidities, n (%)				
AIDS	6 (0.5)	5 (0.5)	1 (0.5)	1 (0.9)
Chronic obstructive pulmonary disease	173 (13.2)	143 (13.0)	30 (14.2)	17 (14.8)
Malignancy	164 (12.5)	137 (12.4)	27 (12.8)	14 (12.2)
Cancer, solid tumour	118 (9.0)	102 (9.3)	16 (7.6)	8 (7.0)
Metastatic cancer	32 (2.4)	27 (2.5)	5 (2.4)	1 (0.9)
Haematological cancer	27 (2.1)	19 (1.7)	8 (3.8)	5 (4.3)
Immunocompromised	104 (7.9)	79 (7.2)	25 (11.8)	15 (13.0)
Corticosteroid therapy	57 (4.3)	39 (3.5)	18 (8.5)	10 (8.7)
Immunosuppression	33 (2.5)	24 (2.2)	9 (4.3)	5 (4.3)
Chemotherapy	37 (2.8)	34 (3.1)	3 (1.4)	2 (1.7)
Cirrhosis	39 (3.0)	27 (2.5)	12 (5.7)	7 (6.1)
Diabetes	239 (18.2)	193 (17.5)	46 (21.8)	25 (21.7)
Heart failure	130 (9.9)	102 (9.3)	28 (13.3)	15 (13.0)
Impaired mobility	120 (9.1)	95 (8.6)	25 (11.8)	10 (8.7)
Malnutrition	45 (3.4)	27 (2.5)	18 (8.5)	12 (10.4)
Peripheral vascular disease	41 (3.1)	27 (2.5)	14 (6.6)	5 (4.3)
Renal failure	106 (8.1)	93 (8.4)	13 (6.2)	5 (4.3)
SAPS II category, n (%)†				
≤23	385 (29.3)	353 (32.1)	32 (15.2)	16 (13.9)
24–33	327 (24.9)	283 (25.7)	44 (20.9)	26 (22.6)
34–44	290 (22.1)	220 (20.0)	70 (33.2)	37 (32.2)
≥45	310 (23.6)	245 (22.3)	65 (30.8)	36 (31.3)
Braden score category‡, n (%)				
Very high risk (≤9)	110 (8.4)	79 (7.2)	31 (14.7)	18 (15.7)

Continued

Table 1 Continued

Characteristics	All patients (N=1312, 100%)*	Patients without pressure injuries (n=1101, 83.9%)*	Patients with ICU-acquired pressure injuries and/or pressure injuries acquired prior to ICU admission (n=211, 16%)*	Patients with ICU-acquired pressure injuries only (n=115, 8.8%)*
High risk (10–12)	370 (28.2)	284 (25.8)	86 (40.8)	51 (44.3)
Moderate risk (13–14)	236 (18.0)	194 (17.6)	42 (19.9)	27 (23.5)
Mild risk (15–18)	396 (30.2)	349 (31.7)	47 (22.3)	19 (16.5)
No risk (19–23)	193 (14.7)	189 (17.2)	4 (1.9)	0 (0.0)
Length of stay in ICU prior to study day (days), M (IQR)	4 (1–10)	3 (1–9)	8 (3–20)	12 (6–26)
Length of stay in ICU (days), M (IQR)	9 (4–24)	8 (3–20)	19.5 (8–45.75)	26.5 (12–53)
Length of stay from ICU admission to hospital discharge (days), M (IQR)	18 (8–40)	16 (8–35)	33 (14.25–61.75)	42 (18–63.75)
Length of stay in hospital after study day (days), M (IQR)	11 (6–28)	10 (5–24)	21 (9–41)	22 (10–42)
Patients still in ICU 3 months after study day	2 (0.2)	1 (0.1)	1 (0.5)	0 (0.0)
Patients still in non-ICU ward 3 months after study day	93 (7.1)	59 (5.4)	34 (16.1)	22 (19.1)
Deceased during hospital stay	271 (20.7)	210 (19.1)	61 (28.9)	29 (25.2)
28-day mortality	196 (14.9)	157 (14.3)	39 (18.5)	15 (13.0)

*Total may not sum to 1312, 1101, 211 and 115, respectively, owing to missing values.

†The range of possible scores is 0–163; higher SAPS II score indicates higher severity of disease and acute illness; scores are categorised according to the sample's quartiles.

‡The range of possible scores is 6–23.

ICU, intensive care unit; M, median; SAPS, Simplified Acute Physiology Score.

to patient ratio for the night shift on the survey day was 1:1 (IQR 0.8–1), reflecting the UK national standard of 1:1 nursing for level 3 patients (these are individuals requiring ventilatory and multiple organ support) and 1:2 for level 2 'high dependency' patients. Only one hospital reported that neither physiotherapy nor dietetic support was available on the study day. Four units did not report on this specification. A list of the preventive measures used is provided in online supplemental appendix B tables 1 and 2. Four units reported that they did not have a section on the patient record for describing PI, and four units entered no data. Of the units, 59 (63%) employed the Waterlow Scale, 11 (12%) used the Braden Scale, 19 (20%) reported using a different PI risk assessment scale, and 1 unit reported not using a scale (4 units entered no data).

The characteristics of the patients included in the study are shown in table 1; the majority were men (59.7%) and the median age was 62 years (IQR 50–72). The median number of days in the ICU before the study day was 4 (IQR 1–10), and 51% of these patients were requiring invasive mechanical ventilation on ICU admission. The source of admission to the ICU was the operating theatre in 34.2% of cases, followed by the emergency room (30.5%) and

the general ward (23.3%). Of the patients, 992 (76%) were known to be alive at 84 days, with 271 (20.7%) known to have died (missing data on 49 patients).

The total number of PIs was 311, of which 148 (47.6%) were acquired in the ICU and 154 (52.4%) prior to ICU admission. Thus, 54.5% of patients with PIs acquired their injuries following their ICU admission. Majority of the PIs were classified as grade 1 or 2 (figure 1A). Of the patients, 115 had ICU-acquired PIs and 96 had injuries that were pre-existing at the time of admission. The number of PIs per patient varied from one to nine sites. Anatomical site of 311 injuries (47.6% present prior to ICU admission) were acquired in the ICU and 163 (52.4%) were prior to ICU admission. The sacral area, along with the heels, mouth and nose, appeared the most vulnerable, with the sacral area and heels accounting for most of the higher grade injuries; of 97 areas with injuries classified as grade 3 or above, 33 (34%) were sacral areas and 28 (29%) were heels (figure 1B).

The mixed multiple logistic regression analysis identified the following factors as independently associated with ICU-acquired PIs: decreasing Braden scores (OR=0.77 (95% CI 0.69 to 0.85), $p<0.001$) and increasing prior duration of ICU stay (0–3 days as reference; 4–6 days




Figure 1 (A) Grade of injury and (B) anatomical site of injury. ICU, intensive care unit.

in ICU before study day, OR=2.29 (95% CI 1.06 to 4.95), $p=0.03$; 7–9 days in ICU before study day, OR=2.30 (95% CI 0.91 to 5.84), $p=0.08$; 10–12 days in ICU before study day, OR=7.77 (95% CI 3.42 to 17.69), $p<0.001$; >12 days in ICU before study day, OR=7.73 (95% CI 3.94 to 15.15), $p<0.001$; table 2).

DISCUSSION

This service evaluation identified that 16% of 1312 patients in a large sample of UK ICUs had an area exhibiting pressure damage on the study day. Although 96 patients already had injuries on admission, 115 had acquired them during their ICU stay. Generally, these injuries were not severe; however, there were some that were, and these injuries come with a human and institutional cost. The impact of PIs is not easy to measure in terms of patient outcome and total cost. In 2004, the estimated annual cost paid by the NHS for treatment of PIs was between £1.4 billion and £2.1 billion a year. A

more recent estimate suggests that the cost of treating a PI varies from £1214 (stage 1) to £14108 (stage 4 more severe).²

Recently, Labeau *et al*¹ conducted a worldwide, prospective, point prevalence study comprising 1117 ICUs in 90 countries and found 6747 PIs in 3526 patients. The proportion of ICU-acquired PIs was 59.2%. They identified several factors associated with ICU-acquired PIs, including older age, presence of organ support and high severity of illness scores. This analysis of the UK data identified a low prevalence of generally low severity of PIs. However, having a lower Braden score and a longer length of prior ICU stay were associated with a greater likelihood of acquiring a PI in the ICU. The global study (including data from many significantly lower resource settings) identified numerous risk factor associations, but the overall prevalence was greater (26.6%), as was the total sample ($n=13\,254$).¹ Importantly, the global cohort was able to identify that local factors, case-mix and especially the type of ICU admission (ie, medical, elective or emergency surgery) were associated with ICU-acquired PI risk.¹⁵

The locations of PIs were mainly the sacral area and the heels, and the more severe injuries tended to be in these areas. This clearly needs continuous focus on equipment and practice development and education, along with communication among the multidisciplinary team. It is a reasonable assumption that the overwhelming majority of the patients were nursed in the supine position on the study day; nevertheless, there were reported injuries to the nose, mouth and ears. The results might have been different had the study been conducted during the COVID-19 pandemic, during which there was a widespread need for nursing patients in the prone position.^{16–24} The prevalence and location of PIs likely would reflect staff experience and training, positioning of patients, and workload.²¹ The mouth and the nose may be damaged when using non invasive ventilation with limited options for interface or suctioning.²³ The nurse to patient ratio was 1:1 for level 3 patients and this is a reflection of UK standards of good care. Due to limitations of the data collected, we cannot comment on the detailed acuity which contributing units experienced in the period running up to the study day. Similarly, we cannot infer that staffing has an impact on the prevalence of PIs; the impact of a period of inadequate staffing will be reflected in PIs sometime later.

The study sought information on preventive measures used.^{2–4} The NICE guidance does not cover the specific issues relating to the critically ill patient^{2–4}; however, a list of preventive measures used in the ICUs is provided in online supplemental appendix B table 1. Data in this table are reflective of current NICE guidelines, but the lack of longitudinal data and the low prevalence of PIs precluded exploration of the relative effectiveness of such measures.

In the UK, the presence of ICU PIs generates a mandatory investigation generally initiated via adverse event

Table 2 Mixed multiple logistic regression model

Variable	OR (95% CI)	P value
Days in ICU before study day		
0–3 days	Reference	
4–6 days in ICU before study day	2.29 (1.06 to 4.95)	0.03
7–9 days in ICU before study day	2.30 (0.91 to 5.84)	0.08
10–12 days in ICU before study day	7.77 (3.42 to 17.69)	<0.001
>12 days in ICU before study day	7.73 (3.94 to 15.15)	<0.001
Age, years	1.01 (0.99 to 1.03)	0.27
Male sex	0.98 (0.61 to 1.56)	0.92
Body mass index, kg/m ²		
18.5–25 (normal weight)	Reference	
<18.5 (underweight)	1.97 (0.80 to 4.85)	0.14
25–30 (preobesity)	0.56 (0.32 to 1.08)	0.053
≥30 (obesity)	0.60 (0.34 to 1.07)	0.08
Braden score	0.77 (0.69 to 0.85)	<0.001
Admission type: medical	1.11 (0.47 to 2.60)	0.82
Admission type: elective surgery	0.83 (0.28 to 2.47)	0.74
Admission type: emergency surgery	1.03 (0.41 to 2.54)	0.96
Chronic obstructive pulmonary disease	0.96 (0.50 to 1.88)	0.92
AIDS	3.67 (0.27 to 50.17)	0.33
Heart failure	1.07 (0.51 to 2.22)	0.86
Peripheral vascular disease	0.97 (0.30 to 3.06)	0.95
Diabetes	1.22 (0.69 to 2.15)	0.49
Cirrhosis	2.25 (0.83 to 6.17)	0.11
Malignancy	0.90 (0.42 to 1.93)	0.80
Immunocompromised	1.96 (0.91 to 4.20)	0.09
Vasopressor use	0.64 (0.34 to 1.21)	0.17
Sedation	0.77 (0.40 to 1.47)	0.43
Muscle relaxant use	1.13 (0.31 to 4.12)	0.86
Mechanical ventilation on admission	1.34 (0.79 to 2.27)	0.28
Renal replacement	1.63 (0.83 to 3.22)	0.16
Simplified Acute Physiology Score II score	0.99 (0.97 to 1.01)	0.46
ICU, intensive care unit.		

reporting systems such as ‘Datix’ (<https://rldatix.com>). This triggers investigation and challenge but may not account the specific issues relevant to the critically ill. The ability to differentiate which PIs were preventable with appropriate measures and those which were not, due to acuity, nutrition, vasopressors, hydration status, etc, would ensure appropriate attention but cannot be evaluated with this study methodology. It has been acknowledged that some PIs, particularly in a critical care setting, are unavoidable.²³

The mixed multiple logistic regression analysis identified decreasing Braden scores as associated with ICU-acquired PIs. The Braden Scale includes largely static factors, and a dynamic system which adjusts risk as time goes on may be worth evaluating. This is even more

relevant given that prior length of stay has a significant association with the incidence of PIs demonstrated in both this and other studies.²⁴ The development of such a scale including elapsed time as a variable would require an extensive longitudinal study; it is not currently clear whether the effort would be justified. Of note, the majority of sites reported using the Braden Scale or the Waterlow Scale; such scales would be the primary trigger for additional measures (online supplemental appendix B table 3); neither Braden or Waterlow has a notion of prior length of stay. The importance of length of stay has been highlighted previously, but this seems not to have translated into current risk assessment.²⁵ Although better risk prediction can be valuable for comparative audit as part

of quality improvement, we do not know if an improved risk prediction score would translate into fewer PIs.

One potential limitation of the study is the sample of ICUs contributing data; we estimate 28% of UK services contribute and we cannot assume this to be a representative sample. However, the site data submitted and the patient diagnostic data are in line with broader UK critical care (online supplemental appendix B table 2).²⁶ Conceivably, participating sites may have had greater interest in PI or evaluative practice, which may be different from non-contributing sites, and be reflected in the quality of care. Finally, bedside nurses may perhaps have been inhibited from reporting injuries over anxieties that this would be regarded badly by managers. An important mitigation of this is that PI reporting is mandatory in the UK and has arguably become routine.

CONCLUSION

The prevalence and the stage of severity of PIs, both ICU-acquired and non-ICU-acquired, were low in adult critically ill patients admitted to UK ICUs. Nevertheless, 16% of patients had evidence of PI on the study day, and this clearly represents an opportunity for improvement. Decreasing Braden scores and increasing ICU stay were identified as risk factors associated with the prevalence of ICU-acquired PIs. The sacral area and the heels are clearly very vulnerable areas with greater numbers and are the sites of more severe injuries.

Author affiliations

¹Intensive Care Medicine, McGill University, Montreal, Quebec, Canada

²Montreal General Hospital, Montreal, Quebec, Canada

³Anaesthetics and Intensive Care, Imperial College London, London, UK

⁴Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

⁵Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast, UK

⁶Universitair Ziekenhuis Gent, Gent, Belgium

⁷Ghent University, Gent, Belgium

⁸Ghent University Hospital, Ghent, Belgium

Twitter Francesca Rubulotta @frubulotta and Mieke Deschepper @MiekeDeschepper

Acknowledgements This work was supported by the European Society of Intensive Care Medicine.

Collaborators UK Collaborating Site Investigators: Aberdeen Royal Infirmary (Aberdeen): Faye Morren; Buckinghamshire Healthcare Trust (Aylesbury): Charlotte Young; Ysbyty Gwynedd (Bangor): Nicola Vaughan-Jones; North Devon District Hospital (Barnstaple): Stephanie Harris; Furness General Hospital (Barrow-in-Furness): Karen Burns; Basildon Hospital (Basildon): Carmel Georgiev, Rosina Shayamano; Royal United Hospital Bath NHSFT (Bath): Ian Kerslake, Peter Creber; Bedford Hospital NHS Trust (Bedford): Ana Vochin, Catherine O'Brien; Belfast Trust (Belfast): Paul Caddell; Ulster Hospital (Belfast): Samantha Hagan, Mandy Hughes; Queen Elizabeth Hospital Birmingham (Birmingham): Tomasz Torlinski, James Sherwin; City Hospital (Birmingham): Santhana Kannan, Amber Markham; Blackpool Teaching Hospitals NHS Foundation Trust (Blackpool): Richard Lebon, Jason Cupitt; Royal Bournemouth Hospital (Bournemouth): Julius Cranshaw, Nigel White; Bradford Teaching Hospital (Bradford): Victoria Marriott, Wendy Milner; Brighton and Sussex University Hospitals (Brighton): Casiano Barrera Groba; Cambridge University Hospitals NHS Foundation Trust (Cambridge): Joao Azoia, Petra Polgarova; Kent and Canterbury Hospital (Canterbury): Shaly George, Ritoo Kapoor; Royal Glamorgan Hospital (Cardiff): Ceri Lynch; Broomfield Hospital GICU (Chelmsford): Nathalie Fox, Karen Cranmer; Broomfield Hospital St Andrews Burns Unit (Chelmsford):

Natalie Fox; Cheltenham General Hospital (Cheltenham): Thomas Llewellyn, Kelly Matthews; St Peter's Hospital (Chertsey): Louise Maltby, Jowenalbao; Countess of Chester Hospital NHS Foundation Trust (Chester): Karen Boulton, Rachel Jarman; Colchester Hospital (Colchester): Karen Baxter; Croydon University Hospital (Croydon): Ashok Sundai Raj, Arif Moghal; Ysbyty Glan Clwyd (Denbighshire): Joanne White; Doncaster Royal Infirmary (Doncaster): Suzanne Barrowcliffe; Dorset County Hospital (Dorchester): Mark Pullett, Vaarisan Ganeshalingam; Western General Hospital (Edinburgh): Rosaleen Baruah; Royal Devon and Exeter NHS Foundation Trust (Exeter): Carole Boulanger, Helen Baker; Frimley Park Hospital (Frimley): Justin Woods, Poe Poe Ei; Medway Maritime Hospital (Gillingham): Vongayi Ogbeide, Paul Hayden; Royal Gloucestershire Hospital (Gloucester): Kelly Matthews; Bolton Royal Hospital (Greater Manchester): Jennifer Hughes, Madhu Balasubramanian; Royal Surrey County Hospital (Guildford): Armored Salberg; The Princess Alexandra Hospital NHS Trust (Harlow): Rajnish Saha, Dagmar Holmquist; Wycombe Hospital (High Wycombe): Charlotte Young; Calderdale and Huddersfield NHS Foundation Trust (Huddersfield): Claire Derbyshire; Hull Royal Infirmary (Hull): Neil Smith, Elizabeth Stones; Ipswich Hospital NHS Trust (Ipswich): Jane Ademokun; West Middlesex University Hospital (Isleworth): Monica Popescu, Maria Schofield Legorburo; Jersey General Hospital (Jersey): Samantha North, Carole Brett; Queen Elizabeth The Queen Mother Hospital (Kent): Helen Jaundoo; Royal Lancaster Infirmary (Lancaster): Jayne Craig; St James University Hospital (Leeds): Simon Whiteley; Leeds General Infirmary (Leeds): Clare Howcroft, Liz Wilby; Royal Liverpool University Hospital POCU (Liverpool): Peter Delve, David Shaw; Royal Liverpool University Hospital ICU (Liverpool): Karen Williams, Ingeborg D Welters; Liverpool Heart and Chest Hospital (Liverpool): Jane McMullen; Hammersmith Hospital GICU (London): Stephen Brett, Leah Flores; Hammersmith Hospital CICU (London): Treiza Trueman-Dawkins; Charing Cross Hospital (London): Francesca Rubulotta, Mae Templeton; St Mary's Hospital (London): John Adams; The Royal London Hospital (London): Catherine Smith, John Prowle; St Bartholomew's Hospital (London): Heather Byers, Andrea McDonnell; University Hospital Lewisham (London): Bernd Oliver Rose, Rosie Reece-Anthony; King's College Hospital NHS Foundation Trust (London): Luis Mendes; Chelsea and Westminster Hospital (London): Marcela Vizcaychipi, Rhian Bull; St George's University Hospitals NHS Foundation Trust (GICU, London): Grace Lacaden, Eleanor Santiago; St George's University Hospitals NHS Foundation Trust (CTICU, NICU, London): Carlos Castro Delgado, Sarah Farnell-Ward; University College Hospital (London): Elaine Thorpe; Macclesfield District General Hospital (Macclesfield): Justine Somerville, Anne Williams; Manchester Royal Infirmary (Manchester): Donna Cummings, Helen Derrick; Freeman Hospital (Newcastle): Sarah Brumwell, Claire Randall; Royal Victoria Infirmary Ward 18 (Newcastle upon Tyne): Nicola McCann, Emma Aves; Royal Victoria Infirmary Ward 38 ITU (Newcastle upon Tyne): Gillian Berry; Royal Gwent Hospital (Newport): Tamas Szakmany, Una Gunter; King's Mill Hospital (Nottingham): Paul Pulak; Nottingham University Hospitals NHS Trust (Nottingham): Nikki Sarkar, Kerry Wright; The Manor Hospital (Oxford): Vitor Gomes; Tunbridge Wells Hospital (Pembury): Jo Jones, Ruth Palfrey; Poole Hospital NHS FT (Poole): Julie Camsooksai; Queen Alexandra Hospital (Portsmouth): Abby Lewis, Antony Eneas; St Helens and Knowsley Teaching Hospitals NHS Trust (Prescot): Ascanio Tridente, Louise Barr; Queen's Hospital (Romford): Tomas Jovaisa, Beverley Thomas; Salford Royal NHS Foundation Trust (Salford): Emma Parkin, Daniel Horner; South Tyneside District Hospital (South Shields): Christian Frey; Royal National Orthopaedic Hospital (Stanmore): Suzanne Bench, Rachel Baumber; Stepping Hill Hospital (Stockport): Phil Broadhurst, Matthew Jackson; North Tees and Hartlepool NHS Foundation Trust (Stockton-on-Tees): Lynne Williams, Michele Clark; Royal Cornwall Hospital Trust (Truro): Jonathan Paddle, Sarah Bean; Mid Yorkshire Hospital (Wakefield): Sarah Buckley, Christopher Palfreeman; Warrington Hospital (Warrington): Sophie Liu, Nicola Allison; Warwick Hospital (Warwick): Ben Attwood, Penny Parsons; West Hertfordshire Hospitals Trust (Watford): Victoria Houghton, Sarah Jane Turner; Southend University Hospital (Westcliff-on-Sea): David Higgins, Egidija Bielskute; Royal Albert Edward Infirmary (Wigan): Nicola Horrigan; Wirral University Teaching Hospital (Arrow Park Hospital) (Wirral): Reni Jacob, Karen Habgood; Queen Elizabeth Hospital (Woolwich): Ahmed Zaki, Amy Collins; Worthing Hospital (Worthing): Jenny Lord, Charalice Ramiro; Yeovil District Hospital (Yeovil): Agnieszka Kubisz-Pudelko, Michelle Kotze; Wrexham Maelor Hospital (Wrexham): Helen Williams.

Contributors FR, SBr, CB and BB were responsible for coordinating the study in the UKFR. SBr drafted the manuscript. MD, SOL and SBI produced a UK extract of data from the global study database. FR, SBr, BB, MD and SOL conducted the analysis. SOL and SBI were the lead investigators for the global study. FR is responsible for the overall content as the guarantor. All authors revised the manuscript for important intellectual content.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Obtained.

Ethics approval The study protocol was reviewed by the Joint Research Compliance Office at Imperial College Healthcare NHS Trust London and by the Health Research Authority. These bodies deemed the survey a service evaluation. All participating trusts/hospitals confirmed registration as a service evaluation according to their local protocols and non-objection from the relevant local Caldicott Guardian. A full description of the methods can also be found in the DecubICUs global study report.¹ Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. An online training module was developed for all clinical data collectors and was published on the study website (<https://www.esicm.org/trials-group-2/decubicus/> online supplemental appendix A). See also Labeau S O, Afonso E, Benbenishty J, et al. Prevalence, associated factors and outcomes of pressure injuries in adult intensive care unit patients: the DecubICUs study. *Intensive Care Medicine* 2021;47:160-9. <https://doi.org/10.1007/s00134-020-06234-9>.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Francesca Rubulotta <http://orcid.org/0000-0001-8644-1728>

Stephen Brett <http://orcid.org/0000-0003-4545-8413>

Bronagh Blackwood <http://orcid.org/0000-0002-4583-5381>

Sonia O Labeau <http://orcid.org/0000-0003-3863-612X>

Stijn Blot <http://orcid.org/0000-0003-2145-0345>

REFERENCES

- 1 Labeau SO, Afonso E, Benbenishty J, et al. Prevalence, associated factors and outcomes of pressure injuries in adult intensive care unit patients: the DecubICUs study. *Intensive Care Med* 2021;47:160-9.
- 2 NICE. Pressure ulcer prevention. The prevention and management of pressure ulcers in primary and secondary care. Clinical guidelines 179. Methods, evidence and recommendations April 2014, 2014. Available: <https://www.nice.org.uk/guidance/cg179/evidence/full-guideline-prevention-pdf-547610509> [Accessed Aug 2021].
- 3 National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel, Pan Pacific Pressure Injury Alliance. *Prevention and treatment of pressure ulcers: clinical practice guideline*. Emily Haesler ed. Osborne Park, Australia: Cambridge Media, 2014.
- 4 National Pressure Injury Advisory Panel, European Pressure Ulcer Advisory Panel, and Pan Pacific Pressure Injury Alliance. *Prevention and treatment of pressure ulcers: clinical practice guideline*. Emily Haesler ed. Osborne Park, Australia: Cambridge Media, 2019.
- 5 Edsberg LE, Black JM, Goldberg M, et al. Revised national pressure ulcer Advisory panel pressure injury staging system: revised pressure injury staging system. *J Wound Ostomy Continence Nurs* 2016;43:585-97.
- 6 National Pressure Injury Advisory Panel. Position statement on staging – 2017 clarifications. Available: <https://npupap.org/page/PositionStatements> [Accessed Aug 2021].
- 7 Bergstrom N, Braden B, Laquazza A. The Braden scale for predicting pressure sore risk - reliability studies. *Nurs Res* 1985;36:205-10.
- 8 El-Marsi J, Zein-El-Dine S, Zein B, et al. Predictors of pressure injuries in a critical care unit in Lebanon: prevalence, characteristics, and associated factors. *J Wound Ostomy Continence Nurs* 2018;45:131-6.
- 9 Cox J. Predictors of pressure ulcers in adult critical care patients. *Am J Crit Care* 2011;20:364-75.
- 10 Society of Critical Care Medicine. Critical care statistics. Available: <https://www.sccm.org/Communications/Critical-Care-Statistics> [Accessed Aug 2021].
- 11 R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria, 2018. <https://www.R-project.org/>
- 12 The Scottish Intensive Care Society audit group. Audit of critical care in Scotland 2018, reporting on 2017, 2017. Available: https://www.sicsag.scot.nhs.uk/publications/_docs/2018/SICSAG-report-2018-2609-final.pdf?55 [Accessed Aug 2021].
- 13 Intensive Care national audit & research centre. Annual Quality Report 2018/19 for adult critical care. Available: <https://online.narc.org/Reports/2019/12/annual-quality-report-201819-for-adult-critical-care> [Accessed Aug 2021].
- 14 Deschepper M, Labeau SO, Waegeman W, et al. Heterogeneity hampers the identification of general pressure injury risk factors in intensive care populations: a predictive modelling analysis. *Intensive Crit Care Nurs* 2022;68:103117.
- 15 Shearer SC, Parsa KM, Newark A, et al. Facial pressure injuries from prone positioning in the COVID-19 era. *Laryngoscope* 2021;131:E2139-42.
- 16 Perrillat A, Foletti J-M, Lacagne A-S, et al. Facial pressure ulcers in COVID-19 patients undergoing prone positioning: how to prevent an underestimated epidemic? *J Stomatol Oral Maxillofac Surg* 2020;121:442-4.
- 17 Kim RS, Mullins K. Preventing facial pressure ulcers in acute respiratory distress syndrome (ARDS). *J Wound Ostomy Continence Nurs* 2016;43:427-9.
- 18 Lucchini A, Bambi S, Mattiussi E, et al. Prone position in acute respiratory distress syndrome patients: a retrospective analysis of complications. *Dimens Crit Care Nurs* 2020;39:39-46.
- 19 Nazerali RS, Song KR, Wong MS. Facial pressure ulcer following prone positioning. *J Plast Reconstr Aesthet Surg* 2010;63:e413-4.
- 20 Azoulay E, Timsit J-F, Sprung CL, et al. Prevalence and factors of intensive care unit conflicts: the conflictus study. *Am J Respir Crit Care Med* 2009;180:853-60.
- 21 Alqahtani JS, AlAhmari MD. Evidence based synthesis for prevention of noninvasive ventilation related facial pressure ulcers. *Saudi Med J* 2018;39:443-52.
- 22 Edsberg LE, Langemo D, Baharestani MM, et al. Unavoidable pressure injury: state of the science and consensus outcomes. *J Wound Ostomy Continence Nurs* 2014;41:313-34.
- 23 Zhang Y, Zhuang Y, Shen J, et al. Value of pressure injury assessment scales for patients in the intensive care unit: systematic review and diagnostic test accuracy meta-analysis. *Intensive Crit Care Nurs* 2021;64:103009.
- 24 Cox J. Risk factors for pressure injury development among critical care patients. *Crit Care Nurs Clin North Am* 2020;32:473-88.
- 25 Coyer F, Cook J-L, Doubrovsky A, et al. Exploring medical device-related pressure injuries in a single intensive care setting: a longitudinal point prevalence study. *Intensive Crit Care Nurs* 2022;68:103155.
- 26 Coyer F, Labeau S, Blot S. Preventing pressure injuries among patients in the intensive care unit: insights gained. *Intensive Care Med* 2022;22:1-3.