



# BMJ Open Association of sarcopenia with ideal cardiovascular health metrics among US adults: a cross-sectional study of NHANES data from 2011 to 2018

Weihua Chen <sup>1</sup>, Shanshan Shi,<sup>1</sup> Yizhou Jiang,<sup>2</sup> Kaihong Chen,<sup>1</sup> Ying Liao,<sup>1</sup> Rongchong Huang,<sup>3</sup> Kun Huang <sup>4</sup>

**To cite:** Chen W, Shi S, Jiang Y, *et al.* Association of sarcopenia with ideal cardiovascular health metrics among US adults: a cross-sectional study of NHANES data from 2011 to 2018. *BMJ Open* 2022;**12**:e061789. doi:10.1136/bmjopen-2022-061789

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-061789>).

WC, SS and YJ contributed equally.

Received 08 February 2022  
Accepted 09 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Kun Huang;  
k-huang18@mails.tsinghua.edu.cn and  
Rongchong Huang;  
rhuang@ccmu.edu.cn

## ABSTRACT

**Objective** The American Heart Association (AHA) proposed the concept of ideal cardiovascular health (CVH) to reduce the risk of cardiovascular mortality. We attempted to broaden the impact of CVH and further contribute to AHA 2030 goals by identifying the relationship between CVH and non-cardiovascular diseases such as sarcopenia.

**Design** Cross-sectional survey

**Setting** National Health and Nutrition Examination Survey conducted in the USA from 2011 to 2018.

**Participants** This study included participants with reliable first 24-hour dietary recall and ≥20 years of age and excluded those who could not diagnose sarcopenia or insufficient data to calculate the CVH scores.

**Primary and secondary outcome measures** The prevalence of sarcopenia as measured by dual-energy X-ray absorptiometry.

**Results** This cohort study involving 9326 adults ≥20 years comprised 4733 females (50.0%). The number of intermediate or ideal and poor CVH participants was 5654 and 3672 with mean CVH score of  $9.70 \pm 0.03$  and  $5.66 \pm 0.04$ , respectively. After adjusting for related confounding factors, intermediate or ideal CVH was associated with an odds reduction of sarcopenia than poor CVH (adjusted OR (aOR): 0.36, 95% CI 0.26 to 0.50,  $p < 0.001$ ) and the odds of sarcopenia was significantly lower for each incremental increase of 1 in CVH metrics (aOR: 0.75, 95% CI 0.71 to 0.79,  $p < 0.001$ ). Moreover, if the number of ideal CVH metrics was >5, the odds of sarcopenia decreased by up to 84% (aOR: 0.16, 95% CI 0.08 to 0.30).

**Conclusions** Our findings suggest a relationship between the CVH and the prevalence of sarcopenia in adults. The results of our study can contribute to achieving the 2030 public health goal of achieving CVH for all, which may be supported by efforts to reduce the prevalence of sarcopenia.

## INTRODUCTION

Life expectancy in the USA has been stagnant since 2010 which has been attributed to a lack of progress in cardiovascular disease (CVD) mortality.<sup>1</sup> Indeed, CVD remains the primary cause of mortality globally and a huge burden on public health expenditure.<sup>2</sup> Previous

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The main strength of this study is the large sample representative of the adult population of the USA.
- ⇒ Use of a validated survey instrument and standardised data collection methods allows for comparison with other studies.
- ⇒ The limitations of this study were that data were derived from cross-sectional studies and that the relationship was not necessarily identified as causal.
- ⇒ Use of self-reported data might result in recall bias.
- ⇒ A half of initial cohort has been excluded in this study, which will increase the variance of the OR estimates.

investigators have used the Framingham and SCORE risk estimation systems to assess a patient's risk for CVD.<sup>3 4</sup> These risk scores are primarily derived from the development and establishment of effective primary and secondary prevention interventions for high-risk populations. However, individuals with significantly elevated levels of risk factors are relatively uncommon in the population. Most CVD and stroke events occur in individuals with average or only slightly unfavourable levels of risk factors. Therefore, the concept of cardiovascular health (CVH) was introduced to reduce the risk of cardiovascular mortality in 2010.<sup>5</sup> CVH includes seven metrics, including body mass index (BMI), cigarette smoking, physical activity, dietary intake, total cholesterol level, blood pressure, and fasting glucose level.<sup>5</sup> The beneficial effects of ideal CVH metrics are widely supported by mounts of scientific research.<sup>6</sup> However, a recent study showed that the prevalence of ideal CVH status is low on some metrics, such as dietary pattern.<sup>7</sup> Moreover, a study involving the offspring of Framingham participants showed that the decreasing presence of ideal CVH metrics over the past 20 years has resulted in increasing risks of

subclinical diseases, CVDs and death.<sup>8</sup> Therefore, there is a long way to go regarding the ‘Strategic Impact Goals for 2030 and Beyond’ issued by the American Heart Association (AHA).

Previous studies have suggested that an ideal CVH is negatively associated with age-related diseases.<sup>9</sup> Sarcopenia, marked by the age-related loss of muscle mass, strength and function, has become a severe medical problem in the current ageing society. A meta-analysis indicated that patients with sarcopenia have decreased function, and higher rates of falls and hospitalisation.<sup>10</sup> Sarcopenia shares many common pathogenic mechanisms with CVDs, such as hormonal changes, inflammation and oxidative stress.<sup>11</sup> Studies have confirmed that sarcopenia is significantly associated with increased cardiovascular events or mortality,<sup>12</sup> and patients with CVDs are also more likely to develop sarcopenia than age-matched controls.<sup>13</sup>

Although several studies have explored the relationship between cardiovascular risk factors and sarcopenia,<sup>14</sup> it remains unclear whether ideal CVH metrics are beneficial in sarcopenic populations.

This study aimed to determine the relationship between CVH and sarcopenia by using the 2011–2018 National Health and Nutrition Examination Survey (NHANES) data to contribute to the accomplishment of the AHA 2030 goals.

## METHODS

### Patient and public involvement

NHANES is a nationally representative health survey designed and administered by the National Center for Health Statistics (NCHS) at the Centers for Disease Control and Prevention and was approved by the NCHS Research Ethics Review Board (protocols Numbers: NHANES Protocol #2011-17 and NHANES Protocol #2018-01). The NHANES was designed to represent the civilian non-institutionalised US population using a complex multi-stage probability sampling methodology. We conducted a retrospective analysis of a cohort of US population of the NHANES from 2011 to 2018. Written informed consent was acquired from each NHANES participant. The NHANES includes extensive demographic data, physical examinations, laboratory tests, health-related questionnaires and lists of prescription medications, which were measured at the start of the study. Further details on the data collection procedure and analytical guidelines are publicly available on the NHANES website.<sup>15</sup> As shown in online supplemental figure S1, this study included participants with reliable first 24 hours dietary recall and  $\geq 20$  years of age during NHANES 2011–2018 ( $n=21\,128$ ). Of these participants, 11 802 were excluded based on the following: (1) no reliable dual-energy X-ray absorptiometry (DXA) and BMI data; and (2) insufficient data to calculate the CVH scores. Thus, 9326 participants were enrolled in the present study.

### DXA, appendicular skeletal muscle mass, and the definition of sarcopenia

DXA whole-body scans were performed on participants 8–59 years of age using Hologic Discovery model A densitometers (Hologic, Bedford, Massachusetts, USA). DXA exclusion criteria included pregnancy, weight  $>300$  pounds (136 kg, because of the weight limit of the scanner), height  $>6'5"$  (DXA table limitations), history of radiographic contrast material (barium) used in the past 7 days, or nuclear medicine studies in the past 3 days. Hologic software (V.8.26: a3\*) was used to administer all scans.

Appendicular skeletal muscle mass was measured using DXA. The sarcopenia index was calculated as follows: sarcopenia index = total appendicular skeletal muscle mass (in kg)/BMI ( $\text{kg}/\text{m}^2$ ).

Sarcopenia was defined as the lowest for sex-specific sarcopenia index cut-off values (0.789 for men and 0.512 for women), based on the National Institutes of Health (FNIH).

### CVH metrics

CVH metrics include four health behaviours (cigarette smoking, physical activity, healthy dietary scores, and BMI) and three health factors (total cholesterol level, blood pressure, and fasting plasma glucose level).<sup>5</sup> The definitions of ideal, intermediate, and poor CVH metrics for adults are presented in table 1. We used the Healthy Eating Index 2010 (HEI-2010) scores as a proxy of healthy dietary scores, which were calculated using first-day 24-hour dietary recall. HEI-2010 scores were based on a 12-component index, with total scores ranging from 0 to 100, and a higher score indicating a healthier diet: total fruit; whole fruit; total vegetables; grains and beans; whole grains; dairy; total protein foods; seafood and plant protein; fatty acids; refined grains; sodium; and empty calories. Participants with an HEI-2010 score  $<50$  were assigned to poor health, those with a score of 51–80 to intermediate health, and those with a score  $>81$  to ideal health.

Although the AHA relies on fasting glucose to determine hyperglycaemic, we use HbA1c (HbA1c) concentrations for two reasons. First, recent recommendations from the American Diabetes Association allow the use of HbA1c to diagnose diabetes. Second, a significant percentage of NHANES participants who took the test did not fast. Therefore, we used HbA1c values  $<5.7\%$ ,  $5.7\%–6.4\%$ , and  $>6.5\%$  as a proxy for fasting plasma glucose levels  $<100$  mg/dL, 100 to  $<126$  mg/dL, and  $>126$  mg/dL. Participants who reported having diabetes or being treated with insulin or an oral medication to lower blood glucose and had an HbA1c concentration between  $5.7\%$  and  $6.4\%$  were categorised as intermediate health. Similarly, participants who reported taking cholesterol lowering or anti-hypertensive medications and were treated to goal were categorised as ‘intermediate’, whereas participants with these conditions who were untreated or who were not treated to goal were categorised as ‘poor’ for that health

**Table 1** Distribution of ideal, intermediate and poor CVH for each metric for adults free of CVD, NHANES 2011–2018<sup>5</sup>

AHA definitions of CVH for each metric		Total sample (n=9326)
Smoking status, n (%)		
Ideal	Never or quit>12 mo	7003 (75.2)
Intermediate	Former≤12 mo	216 (2.9)
Poor	Yes	2107 (22.0)
Body mass index, n (%)		
Ideal	< 25 kg/m <sup>2</sup>	2890 (31.1)
Intermediate	25–29.9 kg/m <sup>2</sup>	2937 (32.6)
Poor	≥ 30 kg/m <sup>2</sup>	3499 (36.3)
Physical activity, n (%)		
Ideal	≥150 min/week moderate intensity or ≥75 min/week vigorous intensity or ≥150 min/week moderate+vigorous	3660 (41.9)
Intermediate	1–149 min/week moderate intensity or 1–74 min/week vigorous intensity or 1–149 min/week moderate+vigorous	625 (7.6)
Poor	None	5041 (50.5)
Healthy diet score *, n (%)		
Ideal	4–5 components	201 (2.2)
Intermediate	2–3 components	4046 (44.3)
Poor	0–1 components	5079 (53.5)
Total cholesterol, n (%)		
Ideal	< 200 mg/dL	5213 (54.1)
Intermediate	200–239 mg/dL or treated to goal	2548 (28.7)
Poor	≥ 240 mg/dL	1565 (17.2)
Blood pressure, n (%)		
Ideal	<120/<80 mm Hg	4474 (49.1)
Intermediate	SBP 120–139 or DBP 80–89 mm Hg or treated to goal	2933 (31.9)
Poor	SBP≥140 or DBP≥90 mm Hg	1919 (20.0)
Glycated haemoglobin A1c, n (%)		
Ideal	< 5.7%	6509 (76.0)
Intermediate	5.7%–6.4% or treated to goal	1945 (16.8)
Poor	> 6.4%	875 (7.2)

\*AHA's healthy diet score includes five components: fruits and vegetables, whole grain, fish, sodium, and sugar-sweetened beverage, and a very small proportion (<0.5%) of US adults meet the ideal healthy diet. HEI-2010 is a continuous score consisting of 12 components representing major food groups including fruit and vegetables, whole grains, proteins, dairy, oils, sodium and empty calories. HEI-2010 score ranges from 0 to 100 with a higher score indicates more healthy diet. HEI-2010 has been validated to represent the diet quality in population. We used HEI-2010 as a proxy for AHA's healthy diet score with ideal diet: HEI-2010≥81; intermediate diet: 51–80; and poor diet: ≤ 50. CVH, cardiovascular health; CVD, cardiovascular disease; NHANES, National Health and Nutrition Examination Survey; AHA, The American Heart Association; DBP, diastolic blood pressure; SBP, systolic blood pressure.

factor. Use of antihypertensive, cholesterol-lowering, and glucose-lowering medications were self-reported. Total cholesterol and plasma glucose levels were measured with enzymatic methods (<https://www.cdc.gov/nchs/nhanes/index.htm>). BMI was calculated as the weight in kilograms divided by the height in metres squared. The mean blood pressure was estimated from up to three readings obtained under standard conditions during a single physical examination.

For each metric, participants received 0, 1, or 2 points, representing poor, intermediate or ideal categories, respectively. Participants with overall scores of 0–7, 8–11, or 12–14 points were categorised as having poor, intermediate, or ideal CVH, respectively. Owing to the relatively low number of people with an ideal CVH score in this sample, the intermediate and ideal CVH categories were combined.

### Statistical analysis

We used the NHANES recommended weights to account for planned oversampling of specific groups. The continuous variables were expressed as the mean±SE, and the categorical variables were presented as counts (percentages). Baseline characteristics between the two CVH groups were compared using a t-test for continuous variables and a  $\chi^2$  test for categorical variables.

Multiple logistic regression was used to examine the independent influence of CVH on sarcopenia comparing poor CVH versus intermediate or ideal CVH after adjustments for potential confounders, such as age, sex, and race/ethnicity, educational level, alcohol, congestive heart failure, coronary heart disease, angina and cancer. The OR and 95% CI were computed. We explored the relationship between CVH and sarcopenia in different subgroups (age, sex, race/

ethnicity and education level). We also separately estimated the association between individual components of the CVH metrics and sarcopenia. When assessing the role of individual components, the age, sex and race/ethnicity, educational level, alcohol, congestive heart failure, coronary heart disease, angina and cancer were adjusted. Furthermore, we used multiple logistic regression analysis to assess the effect of a different number of ideal cardiovascular health metrics (ICVHMs) on the incidence of sarcopenia. A two-sided  $p$  value  $<0.05$  indicated significance for all analyses. All data analyses were performed using SAS Release V.9.4 (SAS Institute) and Survey package in R software (V.4.0.4; R Foundation for Statistical Computing, Vienna, Austria).

## RESULTS

### Baseline characteristics

This study shown that only 2.2% of the participants met the ideal diet criteria. The frequency in the present sample of participants meeting the ideal level for the remainder of CVH metrics were cigarette smoking (weighted, 75.2%), HbA1c (weighted, 76.0%), total cholesterol level (weighted, 54.1%), blood pressure (weighted, 49.1%),

physical activity (weighted, 41.9%), and BMI (weighted, 31.1%) (table 1).

This cohort study involved 9326 adults  $>20$  years of age, comprising 4733 females (weighted, 50.0%) and 4593 males (weighted, 50.0%), with a weighted mean (SE) age of  $39.3 \pm 0.3$  years. 3323 (weighted, 60.3%) were of non-Hispanic white ancestry, 967 (weighted, 7.3%) of Hispanic ancestry, and 1955 (weighted, 11.3%) of non-Hispanic Black ancestry. The study population characteristics are listed in table 2 by CVH metrics. The number of intermediate or ideal and CVH participants was 5654 and 3672, with mean CVH metrics of  $9.70 \pm 0.03$  and  $5.66 \pm 0.04$ , respectively. The differences of CVH metrics were significant for age, race/ethnicity, and education ( $p < 0.001$ ). The frequency in the present sample of sarcopenia in participants with poor CVH metrics was 12.3%, nearly threefold as participants with intermediate or ideal CVH metrics (4.8%). Moreover, we analysed the characteristics of this study population by sarcopenic status. Sarcopenia was identified in 45.9% of 403 females based on the sarcopenia criteria and the non-Hispanic white ancestry more like to develop sarcopenia (47.5%) compared with other races/ethnicities. Furthermore, the patient with sarcopenia had poor education level, BMI risk, healthy diet

**Table 2** Baseline characteristics of the study population

Characteristics	Total (n=9326)	Intermediate or ideal CVH (n=5654)	Poor CVH (n=3672)	P value
Age, mean (SE), years	39.3 (0.3)	36.4 (0.3)	44.3 (0.3)	$< 0.001$
Female, n (%)	4733 (50.0)	2933 (50.9)	1800 (48.5)	0.078
Race/ethnicity, n (%)				
Mexican American	1406 (11.1)	846 (11.4)	560 (10.5)	$< 0.001$
Other Hispanic	967 (7.3)	598 (7.5)	369 (7.0)	
Non-Hispanic White	3323 (60.3)	2067 (61.3)	1256 (58.5)	
Non-Hispanic Black	1955 (11.3)	1001 (9.5)	954 (14.7)	
Other	1675 (9.9)	1142 (10.3)	533 (9.3)	
Heavy use of alcohol, n (%) *				
$< 12$	6636 (97.8)	4111 (98.5)	2525 (96.6)	0.173
$\geq 12$	156 (2.2)	75 (1.5)	81 (3.4)	
Education levels, n (%)				
$< 12$	3675 (34.6)	1989 (31.0)	1686 (40.9)	$< 0.001$
12	3092 (34.0)	1863 (33.6)	1299 (34.6)	
$> 12$	2557 (31.4)	1800 (35.4)	757 (24.5)	
Scores of CVH metrics, mean (SE)	8.24 (0.04)	9.70 (0.03)	5.66 (0.04)	$< 0.001$
No	8519 (92.5)	5326 (95.2)	3193 (87.7)	
Congestive heart failure	97 (1.0)	23 (0.2)	74 (1.8)	$< 0.001$
Coronary heart disease	92 (1.0)	17 (0.2)	75 (2.2)	$< 0.001$
Angina	94 (1.0)	30 (0.4)	64 (2.2)	$< 0.001$
Cancer	349 (5.0)	158 (3.9)	191 (6.8)	$< 0.001$
Sarcopenia, n (%)				
Yes	807 (7.5)	328 (4.8)	479 (12.3)	$< 0.001$
No	8519 (92.5)	5326 (95.2)	3193 (87.7)	

\*Data missing  $>5\%$   
CVH, cardiovascular health;



**Table 3** The association between CVH metrics and sarcopenia by selected subgroups

Variable	No. (%)	Intermediate or Ideal CVH OR (95% CI) *	P value	Intermediate or Ideal CVH OR (95% CI) †	P value	P for interaction
Continuous						
CVH (per 1 score)	807/9326	0.77 (0.74 to 0.79)	<0.001	0.75 (0.71 to 0.79)	<0.001	–
Categories *						
Poor CVH	479/3672	1 (Ref)	–	1 (Ref)	–	–
Intermediate or Ideal CVH	328/5654	0.36 (0.29 to 0.44)	<0.001	0.36 (0.26 to 0.50)	<0.001	–
Subgroup						
Age						
<45	211/4200	0.41 (0.31 to 0.54)	<0.001	0.38 (0.27 to 0.52)	<0.001	0.189
45–59	117/1454	0.37 (0.27 to 0.52)	<0.001	0.32 (0.19 to 0.53)	<0.001	
Sex						
Male	157/2721	0.33 (0.24 to 0.45)	<0.001	0.36 (0.24 to 0.53)	<0.001	0.827
Female	171/2933	0.40 (0.30 to 0.54)	<0.001	0.35 (0.21 to 0.58)	<0.001	
Race						
Mexican American	127/864	0.41 (0.28 to 0.60)	<0.001	0.43 (0.28 to 0.67)	<0.001	0.704
Other Hispanic	44/598	0.32 (0.18 to 0.58)	<0.001	0.37 (0.20 to 0.70)	0.003	
Non-Hispanic White	80/2067	0.30 (0.22 to 0.40)	0.019	0.31 (0.17 to 0.56)	<0.001	
Non-Hispanic Black	8/1001	0.15 (0.07 to 0.32)	<0.001	0.21 (0.08 to 0.50)	<0.001	
Other	69/1142	0.40 (0.23 to 0.69)	0.001	0.45 (0.22 to 0.95)	0.036	
Education levels						
<12	179/1989	0.43 (0.33 to 0.57)	<0.001	0.49 (0.33 to 0.74)	<0.001	0.014
12	87/1863	0.39 (0.25 to 0.61)	<0.001	0.32 (0.19 to 0.54)	<0.001	
>12	62/1800	0.30 (0.18 to 0.50)	<0.001	0.20 (0.10 to 0.40)	<0.001	

Poor CVH: CVH metrics scores 0–7; intermediate or ideal CVH: CVH metrics scores 8–14.

\*Unadjusted model.

†Analyses were adjusted for age, sex, race/ethnicity, education level, alcohol use, congestive heart failure, coronary heart disease, angina and cancer. CVH, cardiovascular health.

score risk, blood pressure risk, HbA1c risk and overall CVH metrics. And more detailed analyses are presented in online supplemental table S1.

### Association between CVH metrics and sarcopenia

The intermediate or ideal CVH was associated with an odds reduction of sarcopenia than poor CVH (OR 0.36, 0.29 to 0.44,  $p<0.001$ ; table 3). After adjusting for age, sex, race/ethnicity, education level, alcohol use, congestive heart failure, coronary heart disease, angina and cancer, intermediate or ideal CVH was associated with an odds reduction of sarcopenia than poor CVH (adjusted OR (aOR): 0.36, 95% CI 0.26 to 0.50,  $p<0.001$ ). In the fully adjusted model, the odds of sarcopenia were significantly lower for each incremental increase of 1 in CVH metrics (aOR: 0.75, 95% CI 0.71 to 0.79,  $p<0.001$ ). Further stratified and interaction analyses were performed for age, sex, race/ethnicity, and education level. And the association between intermediate or ideal CVH and sarcopenia was significant in different subgroups. Notably, the age group also showed stronger association in the subgroup aged<45 years (aOR: 0.38, 95% CI 0.27 to 0.52,  $p<0.001$ ). Further, among subgroups of non-Hispanic Black, the odds of sarcopenia decreased by 79% in participants with intermediate or ideal CVH than in participants with poor CVH (aOR: 0.21, 95% CI 0.08 to 0.50,  $p<0.001$ ; table 3).


For all of subgroups, there was no significant interaction (all  $p$  values for interaction>0.05), except of education levels ( $p$  for interaction=0.014).

### Association between number of ICVHMs and sarcopenia

Twenty-one percent of participants with sarcopenia had only 1 ICVHM and 5% had 5 ideal ICVHMs. In participants without sarcopenia, up to 70% had>3 ICVHMs (online supplemental figure S2). Logistic regression of the ICVHM number and the odds of sarcopenia revealed that the higher the number of ICVHMs, the lower the odds of sarcopenia. When participants had three ideal CVH metrics, the odds of sarcopenia decreased by 50% compared with participants with non-ideal CVH metrics (aOR: 0.50, 95% CI 0.32 to 0.78). If the number of ICVHMs was>5, the odds of sarcopenia decreased by up to 84% (aOR: 0.16, 95% CI 0.08 to 0.30; figure 1).

### Association between different individual CVH components and sarcopenia

In the subgroup analysis of the seven individual CVH components, participants defined as intermediate or poor CVH had a higher odds of sarcopenia odds than those with ideal CVH in all CVH metric subgroups except for the subgroup with cigarette smoking status, total cholesterol and physical activity. Especially in the BMI



**Figure 1** Association between number of ICVHMs and sarcopenia. ICVHMs, Ideal cardiovascular health metrics. Model: adjusted by age, sex, and race/ethnicity, educational level, alcohol use, congestive heart failure, coronary heart disease, angina and cancer.\* $P < 0.05$ .

and healthy diet score subgroups, the odds of sarcopenia decreased >80% (BMI: (aOR: 0.08, 95% CI 0.05 to 0.13,  $p < 0.001$ ); healthy diet score: (aOR: 0.18, 95% CI 0.06 to 0.54,  $p = 0.003$ )). A decreasing odd of sarcopenia trends were observed between increasing levels of CVH components for BMI, healthy diet score, HbA1c and blood pressure (all  $p$  for trend  $< 0.05$ ; table 4).

## DISCUSSION

This study used nationwide, population-based, cross-sectional data to demonstrate a significant association between CVH and sarcopenia and showed a significantly 64% decreased adjusted risk of sarcopenia in subjects with better CVH metrics. For each unit increase in the metrics of CVH, the risk of CVDs decreased by 25%. Furthermore, higher intermediate or ideal CVH metrics were associated with a lower prevalence of sarcopenia.

Our study yielded several interesting findings. First, the CVH metrics were not only associated with CVDs, but also non-CVDs, including sarcopenia. This result agreed with Han *et al*,<sup>16</sup> who also reported that sarcopenia was independently associated with cardiovascular risk factors, including diabetes and hypertension. And these risk factors were shown to be associated with the prevalence of sarcopenia defined by the recommended algorithm of the Asian Working Group in the Chinese elderly.<sup>14</sup> However, these results may only be applicable in patient with high-risk cardiovascular risk factors. In order to explore the association between sarcopenia and

the common individual with average or only slightly unfavourable levels of risk factors, we chose CVH and elaborated on the detail and found that higher intermediate or ideal CVH metrics were associated with a lower prevalence of sarcopenia, as defined by the recommended algorithm of the FNIH in American adults. This finding suggests that the level of CVH influences the incidence of sarcopenia and emphasises the greater importance of CVH for healthcare and medical conditions. A previous study showed that the presence of more desirable CVH indicators was associated with a significant reduction in CVD morbidity and mortality.<sup>17</sup> Our study broadens the application value of the CVH metrics; specifically, the higher the number of intermediate or ideal CVH metrics, the lower the incidence of sarcopenia. It showed that only a small percentage of American adults met the ideal criteria for six or seven ideal health metrics. This result is disappointing, but perhaps not surprising. Furthermore, this result challenges clinical and public health professionals to keep steering the health metrics in the desired direction. In the meantime, additional research is warranted in the future to explore CVH and non-cardiovascular fields to increase public awareness of CVH and promote achievement of AHA 2030 goals.

Second, we further observed the effects of CVH metrics on sarcopenia in different subgroups. We have reported that CVH influences the incidence of sarcopenia not only in the elderly population,<sup>14</sup> but in the younger population. In addition, we demonstrated similar results in the sex and ethnicity subgroups. Surprisingly, it appeared that the ideal CVH metrics affect different levels of participant with different levels of education. Recent study shown that low education compared with high education was associated with lower odds of having ideal CVH.<sup>18</sup> However, it appears that participants with higher levels of education are able to benefit more from the ideal CVH. At the same time, participants with low education levels also reduced the prevalence of sarcopenia by nearly 50% from the ideal CVH. Therefore, we not only need to focus on the ideal level of CVH for participants with low education levels, but also need to further increase the attainment rate of ideal CVH for participation with high education levels to achieve further benefits.

Third, we attempted to determine the effect of each indicator in CVH alone on sarcopenia in this study. Our study showed that reduced HbA1c levels were associated with a decreased risk of sarcopenia. This was consistent with the results of previous studies.<sup>19</sup> This finding may be attributed to the fact that higher blood glucose levels accelerate the loss of muscle mass and strength.<sup>20</sup> In addition, ideal blood pressure was the second significant feature associated with sarcopenia. Han *et al*<sup>14</sup> also found that hypertension is an independent risk factor for sarcopenia. Although the mechanism underlying sarcopenia and hypertension is currently unknown, recent studies have concluded that inflammatory factors during ageing could impair blood flow by damaging the microvascular endothelium,<sup>21</sup> which exerted a detrimental effect on

**Table 4** Adjusted ORs (95% CI) of sarcopenia by individual component of CVH metrics

Variable	OR*	95% CI	P value	P for trend
Smoking status				
Poor	1 (Ref)	1 (Ref)	NA	0.201
Intermediate	0.84	0.33 to 2.12	0.706	
Ideal	1.25	0.87 to 1.80	0.223	
Body mass index				
Poor	1 (Ref)	1 (Ref)	NA	<0.001
Intermediate	0.21	0.16 to 0.29	<0.001	
Ideal	0.08	0.05 to 0.13	<0.001	
Healthy diet score				
Poor	1 (Ref)	1 (Ref)	NA	0.005
Intermediate	0.66	0.46 to 0.93	0.019	
Ideal	0.18	0.06 to 0.54	0.003	
Total cholesterol				
Poor	1 (Ref)	1 (Ref)	NA	0.054
Intermediate	0.91	0.62 to 1.36	0.650	
Ideal	0.68	0.45 to 1.03	0.069	
Glycated haemoglobin A1c				
Poor	1 (Ref)	1 (Ref)	NA	<0.001
Intermediate	0.37	0.21 to 0.63	0.001	
Ideal	0.28	0.14 to 0.36	< 0.001	
Physical activity				
Poor	1 (Ref)	1 (Ref)	NA	0.401
Intermediate	0.93	0.56 to 1.54	0.774	
Ideal	0.89	0.67 to 1.18	0.402	
Blood pressure				
Poor	1 (Ref)	1 (Ref)	NA	<0.001
Intermediate	0.62	0.43 to 0.89	0.010	
Ideal	0.37	0.26 to 0.52	<0.001	

\*Analyses were adjusted for age, sex, race/ethnicity, education level, alcohol use, congestive heart failure, coronary heart disease, angina and cancer. CVH, cardiovascular health; NA, Not Applicable.

the body of the elderly. Additional studies are needed to elucidate the causal relationship between hypertension and sarcopenia. Healthy eating is significantly associated with sarcopenia. The Papaioannou study<sup>22</sup> highlighted the beneficial link between healthy eating and sarcopenia risk. There are several possible mechanisms to explain the beneficial effects of a healthy diet on skeletal muscle. First, a healthy diet rich in fruits and vegetables prevents metabolic acidosis and reduces protein hydrolysis and amino acid catabolism, thus reducing the risk of sarcopenia.<sup>23</sup> In addition, unfavourable dietary patterns, including foods rich in saturated fats, may be detrimental to the maintenance of muscle health,<sup>24</sup> while a fiber-rich diet reduces the risk of sarcopenia.<sup>25</sup> Some studies, however, suggest that a lower BMI indicates the presence of sarcopenia and malnutrition and is associated with higher mortality in the older population.<sup>26</sup> Conversely, obese patients may have a survival benefit.<sup>27</sup> However, our study still found that being overweight or obese can significantly increase the risk of sarcopenia. The poor prediction of physical activity in the present study was unexpected, in contrast to previous studies<sup>28</sup> that suggested only ideal physical activity does appear to be associated with the onset of sarcopenia. This finding might be due to the population in our study cohort included only young and middle-aged adults. Physical activity may be crucial for the occurrence of sarcopenia in the elderly population.

Our study has several limitations. First and foremost, cigarette smoking, physical activity, diseases, and diet were self-reported, and subjected to misclassification and recall bias, which can lead to an overestimated or underestimated association between CVH and sarcopenia. Second, as noted above, for practical reasons, we were not fully compliant with all of the AHA 2020 health indicators. Moreover, our study was cross sectional, so the association between CVH and sarcopenia cannot be interpreted as a direct cause-and-effect relationship. Finally, a half of initial cohort has been excluded in this study, which will increase the variance of the OR estimates. However, our results are still relatively reliable after weighting, since the main missing data are due to missing sampling.

## CONCLUSION

In conclusion, our findings suggested a relationship between CVH indicators and the prevalence of sarcopenia among US adults. Our analysis confirms that CVH extends beyond protection against CVD. More research is needed to clarify the association between CVH and other non-CVDs. The results of our study can help facilitate the 2030 goal of achieving CVH for all because the AHA 2030 goal may be supported by efforts to reduce the prevalence of sarcopenia.

## Author affiliations

<sup>1</sup>Department of Cardiology, Longyan First Affiliated Hospital of Fujian Medical University, Longyan, China

<sup>2</sup>Department of Cardiac Surgery, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

<sup>3</sup>Department of Cardiology, Beijing Friendship Hospital, Capital Medical University, Beijing, China

<sup>4</sup>Department of Industrial Engineering, Tsinghua University, Beijing, China

**Acknowledgements** The authors thank all the participants and staff of the NHANES for their valuable contributions.

**Contributors** WC: was responsible for designing the study, interpretation of the data, drafting the manuscript, revising the manuscript, and the approval of the final version. SS: participated in the design of analyses, data analysis, revising the manuscript, and approval of the final version. YJ: drafting the manuscript, revising the manuscript, and the approval of the final version. YL: interpretation of the data and the approval of the final version. KC: participated in formulating the research question, design of analyses, revising the manuscript, and the approval of the final version. RH and KH: participated in formulating the research question, design of analyses, data analysis, interpretation of the data, and the approval of the final version. All authors: read and approved the final version of the manuscript and are responsible for all aspects of the manuscript.

**Funding** This study was supported by grants from the Summit Talent Plan, Beijing Hospital Management Center (plan no: DFL20190101) (Beijing, China), and the Natural Science Foundation of Fujian Provincial Science and Technology Department (2018J01405).

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by NHANES was approved by the NCHS Research Ethics Review Board (protocols Numbers: NHANES Protocol #2011-17 and NHANES Protocol #2018-01). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. The data link is <https://www.cdc.gov/nchs/nhanes/index.htm>

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Weihua Chen <http://orcid.org/0000-0002-7670-4165>

Kun Huang <http://orcid.org/0000-0002-7418-483X>

## REFERENCES

- 1 Mehta NK, Abrams LR, Myrskylä M. Us life expectancy stalls due to cardiovascular disease, not drug deaths. *Proc Natl Acad Sci U S A* 2020;117:6998–7000.
- 2 Mozaffarian D, Benjamin EJ, et al, Writing Group Members. Heart disease and stroke Statistics-2016 update: a report from the American heart association. *Circulation* 2016;133:e38–60.
- 3 D'Agostino RB, Grundy S, Sullivan LM, et al. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. *JAMA* 2001;286:180–7.
- 4 Conroy RM, Pyörälä K, Fitzgerald AP, et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the score project. *Eur Heart J* 2003;24:987–1003.
- 5 Lloyd-Jones DM, Hong Y, Labarthe D, et al. Defining and setting national goals for cardiovascular health promotion and disease



- reduction: the American heart association's strategic impact goal through 2020 and beyond. *Circulation* 2010;121:586–613.
- 6 Ommerborn MJ, Blackshear CT, Hickson DA, *et al.* Ideal cardiovascular health and incident cardiovascular events: the Jackson heart study. *Am J Prev Med* 2016;51:502–6.
- 7 Peng Y, Cao S, Yao Z, *et al.* Prevalence of the cardiovascular health status in adults: a systematic review and meta-analysis. *Nutrition, Metabolism and Cardiovascular Diseases* 2018;28:1197–207.
- 8 Enserro DM, Vasan RS, Xanthakis V. Twenty-Year trends in the American heart association cardiovascular health score and impact on subclinical and clinical cardiovascular disease: the Framingham offspring study. *J Am Heart Assoc* 2018;7. doi:10.1161/JAHA.118.008741. [Epub ahead of print: 17 05 2018].
- 9 Samieri C, Perier M-C, Gaye B, *et al.* Association of cardiovascular health level in older age with cognitive decline and incident dementia. *JAMA* 2018;320:657–64.
- 10 Beaudart C, Zaaria M, Pasleau F, *et al.* Health outcomes of sarcopenia: a systematic review and meta-analysis. *PLoS One* 2017;12:e0169548.
- 11 Curcio F, Testa G, Liguori I, *et al.* Sarcopenia and heart failure. *Nutrients* 2020;12:211.
- 12 Han P, Chen X, Yu X, *et al.* The predictive value of sarcopenia and its individual criteria for cardiovascular and all-cause mortality in Suburb-Dwelling older Chinese. *J Nutr Health Aging* 2020;24:765–71.
- 13 Fülster S, Tacke M, Sandek A, *et al.* Muscle wasting in patients with chronic heart failure: results from the studies investigating co-morbidities aggravating heart failure (SICA-HF). *Eur Heart J* 2013;34:512–9.
- 14 Han P, Yu H, Ma Y, *et al.* The increased risk of sarcopenia in patients with cardiovascular risk factors in Suburb-Dwelling older Chinese using the AWGS definition. *Sci Rep* 2017;7:9592.
- 15 Centers for disease control and prevention, National center for health statistics. About the National health and nutrition examination survey. Available: [https://www.cdc.gov/nchs/nhanes/about\\_nhanes.htm](https://www.cdc.gov/nchs/nhanes/about_nhanes.htm)
- 16 Han E, Lee Y-H, Kim YD, *et al.* Nonalcoholic fatty liver disease and sarcopenia are independently associated with cardiovascular risk. *Am J Gastroenterol* 2020;115:584–95.
- 17 Ford ES, Greenlund KJ, Hong Y. Ideal cardiovascular health and mortality from all causes and diseases of the circulatory system among adults in the United States. *Circulation* 2012;125:987–95.
- 18 Alam MT, Echeverria SE, DuPont-Reyes MJ, *et al.* Educational Attainment and Prevalence of Cardiovascular Health (Life's Simple 7) in Asian Americans. *Int J Environ Res Public Health* 2021;18:1480.
- 19 Han P, Kang L, Guo Q, *et al.* Prevalence and factors associated with sarcopenia in Suburb-dwelling older Chinese using the Asian Working group for sarcopenia definition. *J Gerontol A Biol Sci Med Sci* 2016;71:529–35.
- 20 Morley JE, Malmstrom TK, Rodriguez-Mañas L, *et al.* Frailty, sarcopenia and diabetes. *J Am Med Dir Assoc* 2014;15:853–9.
- 21 Han K, Park Y-M, Kwon H-S, *et al.* Sarcopenia as a determinant of blood pressure in older Koreans: findings from the Korea National health and nutrition examination surveys (KNHANES) 2008–2010. *PLoS One* 2014;9:e86902.
- 22 Papaioannou K-G, Nilsson A, Nilsson LM, *et al.* Healthy eating is associated with sarcopenia risk in physically active older adults. *Nutrients* 2021;13:2813.
- 23 Kim J, Lee Y, Kye S, *et al.* Association of vegetables and fruits consumption with sarcopenia in older adults: the fourth Korea National health and nutrition examination survey. *Age Ageing* 2015;44:96–102.
- 24 Montiel-Rojas D, Santoro A, Nilsson A, *et al.* Beneficial role of replacing dietary saturated fatty acids with polyunsaturated fatty acids in the prevention of sarcopenia: findings from the NU-AGE cohort. *Nutrients* 2020;12:3079.
- 25 Montiel-Rojas D, Nilsson A, Santoro A, *et al.* Dietary fibre may mitigate sarcopenia risk: findings from the NU-AGE cohort of older European adults. *Nutrients* 2020;12:1075.
- 26 Di Angelantonio E, Bhupathiraju S, *et al.* Global BMI Mortality Collaboration. Body-Mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet* 2016;388:776–86.
- 27 Amundson DE, Djurkovic S, Matwiyoff GN. The obesity paradox. *Crit Care Clin* 2010;26:583–96.
- 28 Bosaeus I, Rothenberg E. Nutrition and physical activity for the prevention and treatment of age-related sarcopenia. *Proc Nutr Soc* 2016;75:174–80.




Figure S1. Flowchart.




Figure S2. Proportion of ICSVHMs in sarcopenia and non-sarcopenia.

**Table S1 Baseline characteristics of the study population by sarcopenic.**

<b>Characteristics</b>	<b>Non-sarcopenic (n=8519)</b>	<b>Sarcopenic (n=807)</b>	<b>p-value</b>
<b>Age, mean (SE), years</b>	39.0 (0.3)	42.9 (0.6)	0.002
<b>Female, n (%)</b>	4330 (50.3)	403 (45.9)	0.060
<b>Race/ethnicity, n (%)</b>			
Mexican American	1124 (9.9)	282 (26.1)	< 0.001
Other Hispanic	846 (6.9)	121 (12.0)	
Non-Hispanic White	3103 (61.3)	220 (47.4)	
Non-Hispanic Black	1905 (11.9)	50 (3.4)	
Other	1541 (9.9)	134 (11.1)	
<b>Heavy use of alcohol, n (%)</b>			
<12	6174 (97.9)	426 (95.8)	0.026
≥12	135 (2.1)	21 (4.2)	
<b>Education level, n (%)</b>			
Less Than High School	3214 (33.2)	461 (52.0)	< 0.001
High School Diploma	2878 (34.1)	214 (32.2)	
More Than High School	2425 (32.7)	132 (15.8)	
<b>Smoking risk, n (%)</b>			
Ideal	1969 (22.2)	138 (18.8)	0.085
Intermediate	202 (2.9)	14 (1.8)	
Poor	6348 (74.8)	655 (79.4)	
<b>Body mass index risk, n (%)</b>			
Ideal	2826 (33.1)	64 (6.5)	< 0.001
Intermediate	2746 (33.5)	191 (21.1)	
Poor	2947 (33.4)	525 (72.4)	



Physical activity risk, n (%)				
Ideal	3371 (42.1)	289 (39.1)	0.274	
Intermediate	579 (7.7)	46 (6.5)		
Poor	4569 (50.2)	472 (54.4)		
Healthy diet score risk, n (%)				
Ideal	188 (2.3)	13 (1.0)	0.001	
Intermediate	3716 (44.7)	330 (38.8)		
Poor	4615 (52.9)	464 (60.0)		
Total cholesterol risk, n (%)				
Ideal	4853 (54.8)	360 (45.7)	0.002	
Intermediate	2292 (28.4)	256 (32.1)		
Poor	1374 (16.8)	191 (22.2)		
Blood pressure risk, n (%)				
Ideal	4203 (46.6)	271 (25.7)	< 0.001	
Intermediate	2650 (33.3)	283 (37.6)		
Poor	1666 (20.1)	253 (36.8)		
Fasting plasma glucose risk, n (%)				
Ideal	6103 (50.4)	406 (32.3)	< 0.001	
Intermediate	1711 (31.8)	234 (33.9)		
Poor	705 (17.8)	167 (33.8)		
Scores of seven healthy metrics, mean (SE)		8.4 (0.0)	6.8 (0.1)	< 0.001
Overall CVH metrics, n (%)				
Poor	3193 (34.3)	479 (59.3)	< 0.001	
Intermediate or Ideal	5326 (65.7)	328 (40.7)		

**Abbreviations:** CVH, cardiovascular health.

Poor CVH: CVH metrics scores 0-7; Intermediate or Ideal CVH: CVH metrics scores 8-14.