To cite: Spinnewijn L,

Aarts JWM, Braat D, et al.

decision-making adoption:

a qualitative exploration

bmjopen-2023-080765

Prepublication history

and additional supplemental

available online. To view these

online (https://doi.org/10.1136/

files, please visit the journal

bmjopen-2023-080765).

Received 10 October 2023

Accepted 03 May 2024

material for this paper are

Unravelling clinicians' shared

through the lens of diffusion of

innovations theory. BMJ Open

2024;14:e080765. doi:10.1136/

BMJ Open Unravelling clinicians' shared decisionmaking adoption: a qualitative exploration through the lens of diffusion of innovations theory

Laura Spinnewijn ^(D), ^{1,2} Johanna WM Aarts, ³ Didi Braat, ¹ Fedde Scheele^{2,4}

ABSTRACT

Objectives This study uses the diffusion of innovations (DOI) theory to comprehensively understand the adoption of shared decision-making (SDM) in clinical practice, specifically focusing on the 'knowledge' and 'persuasion' stages within DOI. We aim to understand the challenges and dynamics associated with SDM adoption, offering insights for more patient-centred decision-making in healthcare.

Design This qualitative study employs a modified framework analysis approach, integrating ethnographic and interview data from prior research, along with additional interviews. The framework used is based on the DOI theory.

Study setting and participants This study was conducted in the obstetrics and gynaecology department of a tertiary teaching hospital in the Eastern region of the Netherlands. It included interviews with 20 participants, including gynaecologists, obstetrics registrars and junior doctors currently practising in the department. Additionally, data from prior research conducted within the same department were incorporated, ensuring the maintenance of contextual consistency.

Results Findings reveal a complex interplay between SDM's benefits and challenges. Clinicians value SDM for upholding patient autonomy and enhancing medical practice, viewing it as valuable for medical decisionmaking. Decision aids are seen as advantageous in supporting treatment decisions. Challenges include compatibility issues between patient and clinician preferences, perceptions of SDM as time-consuming and difficult and limitations imposed by the rapid pace of healthcare and its swift decisions. Additionally, perceived complexity varies by situation, influenced by colleagues' attitudes, with limited trialability and sparsely observed instances of SDM.

Conclusions Clinicians' decision to adopt or reject SDM is multifaceted, shaped by beliefs, cognitive processes and contextual challenges. Cognitive dissonance is critical as clinicians reconcile their existing practices with the adoption of SDM. Practical strategies such as practice assessments, open discussions about SDM's utility and reflective practice through professional development initiatives empower clinicians to make the best informed decision to adopt or reject SDM.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The study builds on a substantial body of prior research and well-established psychological theories.
- ⇒ A thorough and structured approach was used, incorporating diverse data sources and employing framework analysis for data analysis.
- ⇒ While the risk for construction and interpretation related to the framework analysis approach was mitigated by reflecting on researchers' roles and inherent biases, it remains a potential concern.
- ⇒ The study's reliance on data from a single obstetrics and gynaecology department may limit its ability to fully capture the diversity of challenges and facilitators present in implementing shared decisionmaking within different healthcare contexts.

INTRODUCTION

Shared decision-making (SDM) emerged as a transformative approach in medicine in the past decades, revolutionising the decision-making process by placing patients at the centre.¹ It lets patients and healthcare providers make healthcare decisions together, incorporating evidence alongside patients' preferences and values.² Unlike evidence-based medicine, which quickly gained traction and transformed medical practice,^{3 4} SDM has faced and still faces challenges in achieving widespread adoption.^{5 6}

Many studies have reported barriers to SDM technologies implementation, including a non-supportive healthcare system, time constraints, lack of support and the complexity inherent in both healthcare professionals' and patients' abilities to engage effectively in SDM.^{7–11} To overcome some of these barriers, studies have highlighted the necessity of changes in organisational culture and healthcare policies to foster an environment conducive to SDM implementation, including an emphasis on organisational leadership activities aimed at promoting SDM implementation.^{8–11} Additionally, various facilitators

Protected by copyright, including for uses related to text and

employer(s)) 2024. Re-use permitted under CC BY-NC. No

C Author(s) (or their

commercial re-use. See rights and permissions. Published by BMJ. ¹Obstetrics and Gynaecology,

Check for updates

Radboudumc, Nijmegen, The Netherlands

²VU Amsterdam Athena Institute, Amsterdam, North Holland, The Netherlands

³Gynaecological Oncology, Amsterdam UMC Location AMC, Amsterdam, The Netherlands ⁴Obstetrics and Gynaecology, OLVG, Amsterdam, Noord-Holland, The Netherlands

Correspondence to

Laura Spinnewijn; lauraspinnewijn@gmail.com have been identified to promote SDM implementation. These include increasing awareness and knowledge of SDM, providing training and educational programmes on SDM, incorporating SDM into clinical guidelines and quality standards and using decision support tools and aids to facilitate the SDM process.⁷⁻¹¹

Despite concerted efforts, the implementation of SDM remains challenging, laborious and inconsistent.⁶⁻⁸ SDM's complexity necessitates a comprehensive approach to understanding the intricate dynamics involved.⁷⁹ To investigate SDM's enigma, this study employs the diffusion of innovations (DOI) theory, developed by Rogers over 50 years ago, to provide a comprehensive framework for a deeper insight into the SDM adoption process in clinical practice.¹

DOI theory introduces the concept of innovation, defined as an idea or technology perceived as new by potential adopters, and highlights diffusion as the process through which innovations spread.¹² The diffusion process encompasses both planned and spontaneous dissemination of ideas, ultimately leading to social change within a given system.¹² One illustrative example provided by Rogers himself is the dissemination of family planning methods and birth control techniques, which revolutionised reproductive health practices and contributed to declines in birth rates, and improvements in maternal and child health worldwide.¹² In addition to tangible innovations such as birth control, Rogers also discusses more abstract concepts, such as ideas or practices. For instance, the diffusion of the idea of sustainable agriculture within the agricultural community represents a shift towards environmentally friendly, economically viable and socially responsible farming practices, away from conventional, intensive farming techniques.¹²

Despite its relative unfamiliarity among medical professionals, DOI theory has gained substantial recognition and credibility. Both within and outside the medical and health research domains, numerous studies have successfully employed DOI theory-based models to investigate the introduction of new technologies and ideas; however, the majority of these studies have focused on categorising individuals based on their propensity to adopt a new innovation.^{13–19} The advantage of DOI theory in disseminating abstract concepts such as SDM lies in another aspect of the theory, known as the 'diffusion of innovation model', or the innovation-decision process.¹² This model identifies five essential stages in the diffusion process: 'knowledge', 'persuasion', 'decision', 'implementation' and 'confirmation'. In the first stage, 'knowledge', individuals become aware of the innovation's existence and gain an understanding of how it functions. In the 'persuasion' stage, individuals form attitudes towards the innovation, weighing its advantages and disadvantages to determine whether to adopt it. The 'decision' is then made to either adopt or reject the innovation based on the assessment made during the persuasion stage. 'Implementation' involves individuals using the innovation in practice, integrating it into their routines and workflows, while

BMJ Open: first published as 10.1136/bmjopen-2023-080765 on 21 June 2024. Downloaded from http://bmjopen.bmj.com/ on June 11, 2025 at Department GEZ-LTA Erasmushogeschool

'confirmation' entails seeking reinforcement for their decision to adopt the innovation and adjusting their usage accordingly.

While all five stages of the diffusion process are significant, our study places particular emphasis on the first two stages, 'knowledge' and 'persuasion', as they precede the critical decision by any potential adopter to either adopt or reject the innovation. Therefore, we used these stages in our attempt to comprehensively assess and understand SDM adoption or rejection among healthcare professionals. Our goal was to uncover the underlying mechanisms influencing clinicians' decisions to adopt or reject. Drawing on one obstetrics and gynaecology department in the Netherlands as an illustrative case, which has **2** been previously examined in SDM studies resulting in a 8 wealth of qualitative data,^{20 21} our study aims to deepen the understanding of SDM's implementation challenges. Subsequently, when considered alongside relevant other Subsequently, when considered alongside relevant other theoretical frameworks, the findings of this study could g efforts decision-decision-analysis potentially inform policy and practice, guiding efforts towards promoting enhanced, patient-centred decisionmaking approaches.

METHODOLOGY Study design

This study uses a modified framework approach,^{22–24} drawing on two sources of data: ethno-graphic and interview data from two previous studies^{20 21} and separate data obtained through additional interviews. The framework approach provides a systematic and transparent methodology for analysing qualitative data.²² It comprises five distinct research phases: 'familiarisation', 'identifying a thematic framework', 'indexing', 'charting and mapping' and 'interpretation'.²³ The modification consisted of conducting supplementary interviews and using those as an important additional data source after the process of data familiarisation with the previously existing data. This alteration was necessary because it became evident that the existing ethnographic and interview data did not adequately address all components outlined within the DOI theory-based framework.

Setting

, Al training, and similar tec This study took place in the obstetrics and gynaecology department of a tertiary teaching hospital in the Eastern region of the Netherlands. The participants comprised both experienced gynaecologists and less experienced clinicians, including registrars in obstetrics and gynaecology.

Framework analysis

Phase I: familiarisation

In this phase, a team of three researchers (LS, JWMA and FS), all providing gynaecological care and familiar with ethnographic research, reviewed data from two prior research projects on SDM in the same department.^{20 21} The first study was a hospital ethnography that aimed to study physician culture in relation to SDM practice.²⁰ The second study investigated registrars' views on SDM, as an addition to pilot testing a patient feedback tool.²¹ The data were collected in 2018 and 2019, respectively. They included written observations and field notes from witnessing consultations and multidisciplinary meetings, as well as 20 interview transcripts from interviews focusing on clinicians' and nurses' attitudes and behaviours regarding SDM. No patients were interviewed in these research projects.

Our research team discussed these data in light of the DOI theory to feed the subsequent phase in our framework analysis. Specifically, we conducted a preliminary thematic examination, aligning with the initial two stages of the innovation-decision process outlined by Rogers, namely 'knowledge' and 'persuasion'.¹² It became apparent that more than existing data were needed to fill a potential DOI theory-based framework. For example, the way clinicians interpret the attributes of SDM was not consistently apparent in the previous on-site observations, and its coverage in earlier interviews was insufficient. Therefore, additional data were gathered through focused interviews with clinicians working in the department and not interviewed before.

Modification: additional data collection

A semistructured interview guide was developed, incorporating elements from the 'knowledge' and 'persuasion' stages and exploring physician attitudes towards SDM in-depth. Additionally, to gain deeper insights into physician attitudes towards SDM, the same interviewees were asked to complete the Attitudes toward Decision aids fOr PatienTs (ADOPT) adjective checklist. The ADOPT list of adjectives is specifically designed to explore attitudes towards SDM.²⁵ Interviewees were invited to highlight those adjectives they associated with SDM and reflect on their choices. The complete interview guide can be found in the online supplemental appendix A. Two junior interviewers, under the supervision of an experienced researcher (LS), conducted the interviews between June and July 2020. All interviews were recorded and subsequently transcribed verbatim. We obtained informed consent from all participants to use interview data for research purposes. Data saturation was achieved after conducting 20 additional interviews, indicating that further interviews were unnecessary to obtain new insights. Eight interviewees (40%) were fully trained gynaecologists, with an average age of 40.2 years (ranging from 35 to 50 years). 10 (50%) were registrars, with an average age of 33.0 years (ranging from 27 to 37 years). Two (10%) were junior doctors who had completed medical school but had yet to start their specialty training, a common occurrence in the Dutch medical training system. The first junior doctor was 27 years old, and the second was 30 years old.

Phase II: identifying and establishing a thematic framework

The framework used in this study aligns with the initial two stages of the innovation-decision process, namely 'knowledge' and 'persuasion', which were also employed during phase I, the familiarisation phase, of our current study.¹² Consequently, the framework in our study was named the innovation-decision framework. During the familiarisation phase and drawing on Rogers' manuscript, significant subthemes were identified within the broader themes 'knowledge' and 'persuasion'. For further elaboration on these themes and subthemes, refer to Table 1. Additionally, a third theme, 'prior conditions', was incorporated into the framework. This theme pertains to the circumstances and factors preceding the introduction of *E* an innovation. According to Rogers, prior conditions are 8 crucial in adoption and diffusion dynamics within a social system, as they establish a conducive or hindering environment for the adoption process to unfold more smoothly and effectively.¹² These prior conditions involve socioeconomic factors, such as education level or income, cultural norms, the presence or absence of leaders advocating Bul the innovation and the alignment of current policies and regulations with the diffusion of an innovation.¹² Once uses related again, table 1 provides detailed insights into all themes and subthemes within our framework. For a comprehensive rationale behind the inclusion of each theme in our DOI-based framework, please refer to online supplemental appendix B. to text

Phase III: indexing

anc The additional focused interviews were initially open coded by two interviewers (KR and IJ) using the Atlas.ti software programme.²⁶ Collaborative discussions with LS led to the development of a comprehensive codebook, ning, Al training, ensuring consensus on specific coding decisions. This iterative process involved alternating between interviews, transcription and coding.

Phase IV: charting and mapping

Following the coding phase, the codes were organised into subthemes aligned with the innovation-decision framework. The organisation of codes into subthemes was a collaborative effort between KR and LS for subthemes relating to 'prior conditions' and 'knowledge,' and between IJ and LS for subthemes relating to 'persuasion'. Throughout this process, agreements were reached on the mapping of data to ensure consistency and accuracy in the analysis. Next, data from the previous research $\ensuremath{\underline{\mathsf{G}}}$ projects were integrated into these subthemes where **3** applicable, guided by the outcomes of the preliminary thematic examination conducted during phase 1 of our study. The results section provides a narrative description of the findings.

Phase V: interpretation

Data interpretation occurred throughout all phases of the research process, as is common in qualitative research. This approach ensured a comprehensive depth Table 1

THEME	'Prior conditions' The contextual factors within a social system that shape the status quo when introducing an innovation.	Phase 1: Knowledge; Influenced by 'Characteristics of the decision- making unit' The attributes, traits, and dynamics of the potential adopters	Phase 2: Persuasion; Determined by "Perceived characteristics of the innovation The subjective perceptions and evaluations of an innovation formed by potential adopters
SUB-THEME	'Previous practice' The established behaviours, routines, and methods individuals or social systems use before introducing an innovation.	'Socioeconomic characteristics' The demographic and economic factors that shape individuals' societal positions and roles.	'Relative advantage' The perceived benefits and advantages of adopting an innovation compared with the existing alternatives or practices.
	'Felt needs/problems' The perceived gaps or challenges that individuals or social systems recognise and desire to address.	'Personality variables' Individual traits, attitudes, and psychological characteristics that influence the adoption and diffusion of innovations.	'Compatibility' The degree to which an innovation is perceived as consistent, harmonious, and compatible with potential adopters' existing values, beliefs, needs, and experiences.
	'Innovativeness' The degree to which an individual or social system is open to adopting new ideas or innovations.	'Communication behaviour' How information and messages about an innovation are disseminated and exchanged among individuals or within a social system.	'Complexity' The perceived difficulty, intricacy, or complexity associated with adopting and using an innovation.
	'Norms of the social system' The established standards, values, and expectations that guide behaviour within a particular group or community.		'Trialability' The degree to which an innovation can be experimented with or tested on a limited basis before fully committing to its adoption.
			'Observability' The extent to which the results or benefits of adopting an innovation are visible and easily observable by others within a social system.

Innovation-decision framework, based on work from Rogers¹²

of understanding. After all data were collected, the final interpretation was conducted by the broader research team (LS, JWMA, DB and FS), contributing diverse perspectives and enriching the overall insights drawn from the study.

Research team and reflexivity

All authors bring experience from their work within a gynaecology department, potentially introducing bias into their work. This bias was mitigated in one of the previous datasets, as one study was conducted in partnership with independent, external medical anthropologists.²⁰ Additionally, the interviewers and coders in this study had yet to acquire clinical experience. LS is a trained medical educationalist who was an obstetrics and gynaecology registrar during data collection. JWMA is a gynaecological oncologist with expertise in qualitative research and extensive experience conducting research focused on person-centred care. She has worked both domestically and internationally, thereby cultivating a deep understanding of various healthcare systems and the complexities associated with their implementation. DB is a retired gynaecologist and a Professor in Reproductive Medicine. FS is a gynaecologist and a Professor in Health Systems Innovation and Education.

RESULTS

Results are presented in alignment with the (sub)themes from our DOI-based framework.

Prior conditions: previous practice

Protected by copyright, including for uses related to text Drawing primarily from data from the two previous and studies, it was observed that medical evidence from clinical trials played and still plays a significant role in practice guidelines and clinicians' daily practice. Patient perspectives were not consistently prioritised in meetings or initial treatment decisions. The training of medical professionals emphasised medical knowledge and guide-. ح lines rather than coaching on handling patient preferences. In situations with limited medical evidence, clinicians strived to reach a consensus within the team before discussing treatment options with patients. While nd clinicians often assumed they knew what was best for their patients, they also demonstrated dedication to patient well-being by actively seeking to understand their needs and providing comprehensive support.

Prior conditions: felt needs

technolog Accurately identifying genuine felt needs for using SDM posed a challenge, as felt needs are less apparent in interviews and observational data. However, indications of an intrinsic desire to involve patients in their care pathways were observed. An example is the patients' presence and active participation during morbidity and mortality meetings at the department,²⁷ which is not common in the Dutch healthcare system.

In interviews, physicians demonstrated a strong awareness of patients' wishes and preferences, highlighting the necessity for tailored decision-making. Explicitly

related

and

mentioned reasons supporting the need for SDM included patient autonomy, adherence to good medical practice, the perception of SDM as the gold standard and the belief that SDM is the optimal approach in medical decision-making. The explicit incorporation of SDM into patient consultations, such as through decision aids and other SDM-supporting materials, through initiation by patients, or promoted by clinicians or their supervisors, increased the perceived need for SDM. One registrar made a concrete suggestion on how to foster SDM integration into group meetings dedicated to discussing patient cases:

I think one sentence should be used: 'The patient prefers:...' That says enough, and one can carry that with oneself. Source: additional interviews, registrar #2

However, there were also instances when a limited need for SDM was expressed. Most clinicians believed that SDM should primarily be employed when multiple comparable treatment options are available or when choices are unclear while still ensuring safety boundaries:

I think it is important that you stay within medically safe options. However, if there are comparable options: yes. And the patient has a choice: Yes, it is necessary. They have to deal with the result, not us. Source: additional interviews, gynaecologist #7

Prior conditions: innovativeness

The level of innovativeness was hard to establish from the data as well. However, based on a few SDM-related initiatives initiated and visible within the department during data collection, an innovative nature was suggested. For example, the gynaecological oncology department was in the middle of a research project designing cancer-specific decision aids,²⁸ and initiated the previously mentioned patient participation at morbidity and mortality meetings.²⁷ In general, we observed many interactions concerning the discussion of new (scientific) insights and the application of up-to-date scientific knowledge.

Prior conditions: norms of the social system

As in any social system, cultural and language differences between clinicians and patients influence decisionmaking. Direct communication was common in the department under study, and gynaecologists were typically approachable, involved and open to new insights. Empathy and involvement were prominent traits among gynaecologists towards both patients and colleagues. However, a hierarchical structure was still observed within team meetings, with gynaecologists generally holding decision-making power over others. Nurses were present in meetings but usually did not actively participate in discussions or treatment decision-making.

Clinicians generally valued the well-being and opinions of patients. However, some prioritised their own opinions over patients' wishes, for example, prioritising survival chances over the quality of life during decisionmaking. We also observed that dealing with uncertainty was a concern for clinicians, and uncertainties were more frequently discussed among physicians than with patients.

Knowledge: socioeconomic characteristics

The group of clinicians, both gynaecologists and registrars, were very homogeneous. Specifically, they exhibited similarities regarding their higher socioeconomic status and cultural backgrounds.

Knowledge: personality variables

Protected Certain clinicians demonstrated higher interest and by copyright, including engagement in SDM than others. Through our observations, we noticed distinct personality differences among physicians, with some individuals displaying more assertiveness and expressing their opinions more firmly than others. The doctors themselves also acknowledged these differences.q

Knowledge: communication behaviour

₫ We identified a notable absence of emphasis on SDM in local protocols and national guidelines. Moreover, **G** during meetings and handovers, patients' preferences were lacking in discussion, meaning SDM was hardly integrated. Additionally, we observed variations in engagement and assertiveness among participants during group discussions. It became evident that more proactive and assertive individuals often assumed leadership roles when shaping policies and making decisions.

Persuasion: relative advantage

data m The use of decision aids offered specific advantages, benefiting both clinicians and patients in the process of making treatment decisions. They provided valuable a structure, assisting in clarifying the patient's preferences training, and desires. They were crucial in presenting a comprehensive and realistic understanding of treatment options and their potential outcomes to the patient. As one partic-, and similar technol ipant highlighted:

Well, for example, I use the decision aid for heavy menstrual blood loss (...). That also gives me structure. One uses it as a handle to have that conversation. Source: previous interview data,²¹ registrar #7

Participants considered it advantageous that SDM allowed patient interests to be visible, leading to better understanding and insight into the patient's thoughts **\$** and feelings. It led to a stronger doctor-patient relationship, with patients appreciating the shared responsibility between clinicians and patients in the decision-making process, as well as improved health outcomes, better quality of care and providing patients with better explanations and informed decision-making. SDM involved patients more consciously in the decision-making process, making them aware of disadvantages and reducing regret or blame in case of adverse outcomes:

You also create clear expectations with the patient by explaining which treatment option has risks and what success rates comprise. So, I also think that, ultimately, if a treatment does not work, patients will at least appreciate being included in the process. Source: additional interviews, gynaecologist #3

Multiple interviewees suggested that SDM improved compliance and patient motivation, leading to potentially more effective treatment. According to them, SDM was associated with increased patient satisfaction, a better feeling about the treatment, and reduced regret. According to the clinicians, patients felt seen, heard and taken seriously during SDM.

The exercise with the adjective list, where interviewees had to indicate which adjectives described their attitude towards SDM best, underlined this positive stance towards SDM. There, the adjective 'rewarding' was chosen by eleven out of 20 participants, against none of them choosing the adjective 'unrewarding'.

All these relative advantages, however, were downsized by felt disadvantages. For example, the often-mentioned belief that SDM is 'time-consuming'; eight participants chose this adjective for SDM. Others, however, described that although it may initially take time, SDM ultimately saves time by fostering patient awareness and understanding, preventing future difficulties or regrets:

At first, I thought: 'Geez, that takes much time.' However, one has a satisfied patient who does not come up with many questions afterwards, saving time. So that is also an advantage. Source: additional interviews, registrar #4

Other often chosen adjectives from the ADOPT adjectives list describing SDM as potentially disadvantageous were 'difficult' or 'laborious', each chosen by three out of 20 participants. Furthermore, although clinicians often felt more appreciation from patients when applying SDM principles, varying levels of interest from colleagues for SDM were described. In intercollegial interactions, negative aspects associated with SDM often received more attention than success stories.

Persuasion: compatibility

During our practice observations, we noticed that physicians dedicated considerable attention to providing proper patient information during consultations. Some clinicians extended consultation times to ensure adequate patient education. However, SDM was predominantly observed in follow-up consultations when care was less guideline-driven or clear-cut. Examples highlighted the challenges of balancing SDM and the performance of simple medical actions, particularly in diagnostic procedures. For instance, a biopsy was taken during a routine patient consultation without extensive patient involvement in decision-making.

Another compatibility issue arose when a patient's wishes diverged from the doctor's belief regarding this and

data

Ξ

<u>0</u>

logies

patient's optimal course of action. Then, varying levels of support were observed among clinicians, and instances indicated that the patients' opinions were not always prioritised. Overall, there were considerable variations between individual physicians regarding how extensively patients' needs and wishes were addressed.

In our additional interviews, the adjective 'realistic' was chosen seven times, and most participants believed that SDM aligned with their current practice. However, according to the interviewees, not all patients desire SDM, and team policies or strong advice can override it. A few clinicians expressed that SDM was unsuitable for certain patients who find it burdensome or are unable to make Š choices, suggesting that a paternalistic approach might be more suitable in such cases. Some admitted to being more 8 controlling in specific situations, such as when there is a superior medical option, when patients explicitly request the doctor to decide or when patients face difficulties in including for uses related to text decision-making. Emergencies were also perceived as less suitable for SDM, and time constraints were mentioned as a reason to be more controlling:

Well, in my experience, if you have a hectic consultation schedule and end up running forty-five minutes behind at the end of the day, that [SDM] becomes constrained by time pressures. You think: Well, [a particular treatment option] would suit this woman. Consequently, you start steering towards a specific course of action more rapidly, rather than calmly engaging with the decision aid, discussing the options, and allowing the patient to leave and revisit the matter in the following week. Source: additional interviews, gynaecologist #3

We observed several limitations concerning time constraints: short consultation times, swift treatment protocols limiting the available time for clinicians and their patients to engage in a collaborate decision-making ≥ training, and process and a fast-paced discussion of individual patient cases during meetings. As a result, there was limited space for SDM.

Persuasion: complexity

Our interviews showed that levels of experience in gynaecology care provision impacted the perceived difficulty of SDM, as registrars more frequently mentioned finding SDM challenging compared with senior clinicians. The perceived complexity of executing SDM was situationally determined and aligned with the compatibility issues mentioned earlier. Factors such as patient intelligence and their ability to understand the SDM process and content, influenced by language barriers and cultural differences as well, played a significant role in determining the difficulty for healthcare professionals. Difficulties arose when options were unequal, patient preferences were unclear, unrealistic patient wishes existed or clinicians were biased due to their opinions.

If you feel that a patient makes a choice that is very remote from you, which you do not support yourself,

Protected

6

then it gets difficult. Because then, as a doctor, I might not want to participate in this treatment because that is not good for this patient. Is that shared decisionmaking? To me, it is not shared decision-making because we did not do it together. Source: additional interviews, junior doctor #2

Sometimes, colleagues complicate the SDM process as well. Some participants expressed a lack of appreciation and felt the need to justify their decisions to colleagues when their decision-making deviated from the usual or medically preferred practice. As one registrar puts it:

When I did that [SDM] with that patient, I felt very good about it. I thought, 'Wow, I had a good conversation with that lady. (...) We weighed everything from both sides. [We made] a good decision'. However, I had to discuss it with the obstetrician for another half hour, who ultimately could not proceed. (...) That is almost bizarre that you get so much resistance. Source: additional interviews, registrar #10

Persuasion: trialability

In the focused interviews, we found that while some participants (nine out of 20) did not actively or consciously experiment with SDM or faced challenges, many felt they had the freedom to explore different approaches and techniques. However, the opportunity to experiment was influenced by factors such as their level of training or experience, which supervisor was involved, or the specific cases they encountered:

In more complex cases, the supervisor usually leads discussions. So, I have not had much experience with those academic care problems yet. I think I have not had the opportunity to start experimenting [with that] on my own. However, depending on the supervisor, I might get the chance if I ask for it. Source: additional interviews, registrar #3

A few participants mentioned a lack of knowledge and guidance on effective experimentation as barriers to exploring SDM. Time pressure was also identified as a significant limitation to SDM experimentation.

Persuasion: observability

Our observations showed a general lack of discussion about SDM in teaching moments, handover discussions and tumour board meetings. Furthermore, it was only sporadically observed in other meetings.

Most participants mentioned discussing SDM with their colleagues sometimes. These discussions tended to focus more on the medical aspect and patient perspective rather than specific conversation techniques related to SDM:

For example, if you have a difficult case or difficulty deciding with a patient, I sometimes talk about that with colleagues. We then discuss what I found particularly difficult in reaching a decision (...). However,

we also discuss medical content, like 'What would you do in this situation? Would you give these drugs or something else?' Source: additional interviews, junior doctor #1

10 out of 20 interviewed physicians did not observe the use of SDM by their colleagues, mainly because it occurred during individual patient consultations without colleagues being present. Some participants had suspicions about its application or the lack thereof.

DISCUSSION

The adoption or rejection of SDM is a complex process by that requires careful consideration of its benefits alongside acknowledgement of existing limitations and challenges. Using Rogers' framework has been instrumental in examining this intricate process, providing a comprehensive approach to identifying the key attributes influencing SDM adoption within our study's context. Through evaluating the characteristics, traits and dynamics of potential adopters, as well as assessing subjective perceptions and evaluations of the innovation, we gained deeper insights into clinicians' decisions to embrace or decline SDM.

In our study setting, clinicians acknowledged the need to use SDM, to uphold patient autonomy and adhere to good medical practice. They perceived SDM as a potentially valuable approach to medical decision-making, with ð decision aids offering specific advantages for both clinie cians and patients in treatment decision-making and SDM adoption. However, clinicians also recognised the limitations and challenges associated with SDM. Disadvantages found are perceptions that it is time-consuming, difficult or laborious. Compatibility issues arose when patients' wishes differed from clinicians' views or when equivalent treatment options were unavailable, and the pace ≥ of healthcare practice sometimes hindered SDM executrain tion. The perceived complexity of SDM varied depending on the situation and often aligned with compatibility ğ issues. Colleagues also contributed to SDM's perceived complexity, as clinicians felt pressure to justify their decisions towards them, and there was a lack of appreciation for good SDM practice. Additionally, clinicians did not recognise the trialability of SDM, and SDM was sparsely observed, both in meetings and doctor-patient encounters.

The use of theory to explain implementation failure **D** is not new in itself. Multiple theories have been applied to capture factors influencing SDM implementation, including normalisation process theory (NPT).²⁹⁻³¹ NPT examines the mechanisms by which new and complex practices, technologies or interventions become routine and normalised within social contexts, focusing on the processes of implementation, embedding and integration.^{29 30} It has often been used for evaluations of innovation implementation.³¹ Examples from within SDM research include the evaluation of routine embedment of decision aids within healthcare settings,²⁹ and the exploration of healthcare providers' perspectives on implementing SDM in routine practice.³⁰ However, NPT highlights innovation benefits rather than delving into reasons for rejection.³¹ In our perspective, recognising and understanding the adverse influences impacting SDM adoption is pivotal for establishing a foundation to drive innovation. Therefore, NPT was deemed less beneficial for the development of our framework and DOI theory was chosen to feed the framework's content instead.

Moreover, current change practices and evaluation theories such as NPT tend to overlook complexities associated with behavioural change, often adopting a reductionist approach when addressing barriers to SDM. The reductionist approach simplifies the determinants of change by treating them as measurable independent variables, assuming that addressing these variables successfully will automatically result in a shift in practice.³² This linear and causal perspective fails to encompass the complete intricacy of SDM adoption. Our study introduces a novel framework for evaluation based on the DOI theory, aiming to comprehend better why clinicians fail to adopt SDM. The next logical step would involve better facilitation of behavioural change, yet without assuming the causal relationships mentioned above.

Efforts to enhance the implementation of SDM have been ongoing, as overviewed in a series of articles following the International SDM Conference in 2022.³³ Furthermore, a 2021 review thoroughly examined strategies for integrating patient decision aids into routine healthcare settings, drawing insights from 23 implementation studies.³⁴ Key success factors identified included collaborative development with patient groups, adaptation of aids to local contexts, securing senior-level endorsement and demonstrating SDM benefits through outcome measures. However, despite their significance, these findings may oversimplify the complexities of real-world scenarios. As the same researchers acknowledged, results from controlled studies may not accurately mirror practical experiences, potentially leading to underperformance and slow adoption of decision aids.³⁴ Thus, understanding the root causes of implementation delays is crucial, extending beyond the factors highlighted for successful implementation. Moreover, it is essential to recognise that implementing decision aids does not inherently ensure successful SDM; rather, decision aids serve as supportive tools for SDM but do not substitute for the practice itself. Consequently, despite our framework analysis, it is imperative to delve into the reasons behind unsuccessful SDM implementation. This involves considering insights from psychology and behavioural change.

Cognitive dissonance theory, initially proposed by Festinger in 1957, offers valuable insights for understanding behavioural change.³⁵ This theory explains the

psychological discomfort individuals experience when they hold conflicting beliefs or attitudes.^{35,36} In the context of SDM, clinicians who recognise the importance of patient involvement may experience cognitive dissonance when confronted with their current practices, which lack SDM. This discomfort can compel them to seek justifications for their existing approaches. For example, they may cite time constraints or difficulties in implementing SDM for patients with low health literacy as good reasons for not executing SDM, as clinicians participating in our study mentioned as well. Additionally, clinicians may find reassurance in patient satisfaction with their current practices, which further discourages using SDM. Moreover, the preference for the status quo, commonly known as the status quo bias, reinforces these arguments against SDM adoption in certain situations.³⁷

SDM adoption in certain situations.³⁷ Despite the potential adverse effects, cognitive dissonance can also serve as a powerful motivator for behavioural change, particularly when individuals hold conflicting solid beliefs.^{38 39} In our opinion, it is crucial to carefully address these dissonances in order to achieve practice change successfully. The following steps can aid in this process.

First, it remains essential to identify professionals' current cognitions and beliefs. A thorough understanding of the consonant and dissonant beliefs influencing professionals' decisions to adopt or reject SDM is essential to address them effectively. Conducting practice assessments, similar to the approach used in our study, can be instrumental in uncovering these often covert beliefs.

Second, there needs to be more open discussions as surrounding these beliefs, particularly regarding the usefulness of SDM. While open communication within healthcare settings has been examined, mainly about professional or patient safety concerns and the importance of speaking up during incidents,^{40 41} research on tively limited.⁴² However, open communication is closely intertwined with significant, well-described psychological **g** constructs, such as those outlined in social constructs, such as those outlined in social interdependence theory.^{43–45} This theory explores how individuals' actions are influenced by their perceived interdependence, whether positive through cooperation towards shared goals or negative through competition and rivalry.43 Contextual factors play a mediating role in determining whether interdependence is constructive. Mediating factors include valuing diverse perspectives, a addressing emotions alongside factual information and $\overline{\mathbf{g}}$ maintaining open-mindedness (p109–110).⁴³ These findings align with a Dutch study highlighting the significance of an 'open culture' encompassing various aspects such as thoughts, emotions, attitudes, safety and wellbeing.⁴² These insights underscore the importance of openly discussing SDM beliefs and experiences. They extend beyond what is typically stated in the literature as an existing 'culture supporting SDM'.^{8 10} Through discussions with colleagues, making an informed argument to

either adopt or reject SDM becomes possible. It is only through this process that a culture truly supportive of SDM can emerge.

Third, and in connection with the previous argumentation, fostering reflection is paramount. Reflective practice has long been recognised as an effective approach in continuous professional education, surpassing the limitations of simply prescribing or teaching new methods.⁴⁶ Through self-reflection, professionals critically evaluate their behaviours and decide whether to change or maintain them. Hence, openly addressing the cognitions and beliefs that hinder or promote the decision to adopt SDM is crucial. To foster behavioural change, training initiatives should not solely focus on straightforward situations. They should also address challenging scenarios where practitioners commonly decide against SDM due to how they resolve their dissonant beliefs. Understanding the arguments used and facilitating open-minded discussions weighing these arguments is once again crucial in the decision to either adopt or reject the use of SDM.

This need for facilitated reflections becomes even more apparent when considering more experienced professionals. Reflective practice and subsequent practice change tend to decline with increased experience, necessitating additional efforts to facilitate reflections among more experienced practitioners.⁴⁷ Particularly in these cases, advocating solely for SDM as the best approach is unlikely to yield positive results. Instead, concerted efforts should be made to challenge prevailing beliefs and practice routines, maintaining an open attitude towards potential flaws in the SDM approach in certain situations. Ultimately, regardless of external or cultural influences, the individual professional possesses the agency to either adopt or reject SDM.^{48 49} Therefore, addressing these individuals most effectively is essential.

Our study acknowledges several limitations. One limitation concerns the framework analysis approach, which inherently risks being influenced by the researchers' preconceptions and assumptions, despite our proactive measures to mitigate bias through critical reflection on our backgrounds. We further addressed this limitation by drawing on well-established psychological theories to formulate broadly relevant recommendations. Additionally, we employed DOI theory as an existing framework to guide our data analysis, providing a structured and theoretically grounded approach. Furthermore, our study focused solely on the role of clinicians within SDM, overlooking patient perspectives within this collaborative process. This decision was driven by the recognition that clinicians often determine the use of SDM. Another limitation is our study's narrow focus on obstetrics and gynaecology within a single hospital department in the Netherlands. While this specialty is dynamic and characterised by preference-sensitive healthcare decisions, the findings may not be generalizable due to the unique nature of the field. For example, decision-making in obstetrics and gynaecology often occurs under time constraints, such as

Protected

<u></u>

ſe

at

data

mining, AI training,

and similar technologies

during critical moments of childbirth or cancer diagnosis. However, it was not our goal to identify universal and generalisable study results. Instead, our findings underscore the importance of conducting comprehensive practice assessments, similar to our approach, before formulating implementation strategies for any medical specialty or department. This ensures that strategies are tailored to the unique dynamics of each local context.

CONCLUSION

Adopting or rejecting SDM is a complex process influenced by copyright by beliefs, cognitions and contextual challenges. Our DOIbased framework analysis aids in identifying these influential factors. Cognitive dissonance plays a significant role as clinicians seek justifications for their current practices or for embracing SDM. By employing strategies such as practice assessments, fostering open discussions on the usefulness of SDM, and promoting reflective practice in, for example, continuing professional development initiatives, we can empower individual clinicians to make the best choices ō regarding the adoption of SDM. Future research should focus on understanding the cognitive and behavioural factors influencing SDM adoption and developing evidencebased strategies within a certain practice setting, to empower ēd informed decision-making in adopting or rejecting SDM practice. The framework employed in our study holds of promise for conducting localised practice assessments, enabling the identification of implicit beliefs specific to each setting. Leveraging this assessment can inform the design of effective implementation strategies with the design of the set of the s effective implementation strategies suited to local contexts.

Acknowledgements The authors want to thank all clinicians interviewed for their participation in this study. Special thanks go to Kim Rutten and Iris Jansen for their contributions in data acquisition for this study.

Contributors LS: Conceptualisation, methodology, investigation, data curation, formal analysis, writing-original draft, writing-review and editing and project administration. JWMA: Conceptualisation, formal analysis and writing-review and editing. DB: Formal analysis and writing-review and editing. FS: Conceptualisation, methodology, formal analysis, writing-review and editing and supervision.

Funding This study received no (additional) funding. Some data for this study were drawn from interviews conducted in a previous study funded by the Education and Training Region East Netherlands (OOR-ON). Grant Number: NA.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. This study was judged by the regional medical ethical assessment committee and the local Radboudumc research ethics board, and it was determined to be exempt from the need for formal ethical approval (file numbers 2017-3887; 2019-5389). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information. No data are available. Data consist of written out observations, other fieldnotes and raw interview data. Data are subject to interpretation and therefore not primarily open for publication.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been

Open access

peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/bv-nc/4.0/.

ORCID iD

Laura Spinnewijn http://orcid.org/0000-0002-8708-2744

REFERENCES

- 1 Coulter A. Partnerships with patients: the pros and cons of shared clinical decision-making. J Health Serv Res Policy 1997;2:112-21.
- Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean? (or it takes at least two to tango). Soc Sci Med 1997:44:681-92.
- 3 Barratt A. Evidence based medicine and shared decision making: the challenge of getting both evidence and preferences into health care. Patient Educ Couns 2008;73:407-12.
- Djulbegovic B, Guyatt GH. Progress in evidence-based medicine: a quarter century on. Lancet 2017;390:415-23.
- 5 Stiggelbout AM, Weijden T d., Wit M, et al. Shared decision making: really putting patients at the centre of Healthcare. BMJ 2012;344:e256.
- Couët N, Desroches S, Robitaille H, et al. Assessments of the extent to which health-care providers involve patients in decision making: a systematic review of studies using the OPTION instrument. Health Expect 2015;18:542-61.
- 7 Joseph-Williams N, Elwyn G, Edwards A. Knowledge is not power for patients: a systematic review and thematic synthesis of patientreported barriers and Facilitators to shared decision making. Patient Educ Couns 2014;94:291-309.
- Waddell A, Lennox A, Spassova G, et al. Barriers and facilitators 8 to shared decision-making in hospitals from policy to practice: a systematic review. Implement Sci 2021;16:74.
- van Veenendaal H, van der Weijden T, Ubbink DT, et al. Accelerating q implementation of shared decision-making in the Netherlands: an exploratory investigation. Patient Educ Couns 2018;101:2097-104.
- Scholl I, LaRussa A, Hahlweg P, et al. Organizational- and system-10 level characteristics that influence implementation of shared decision-making and strategies to address them - a scoping review. Implement Sci 2018;13:40.
- Légaré F, Stacey D, Turcotte S, et al. Interventions for improving the 11 adoption of shared decision making by healthcare professionals. Cochrane Database Syst Rev 2014;2014:CD006732.
- Rogers EM. Diffusion of Innovations. New York: Free Press, 2003. 12
- 13 Greenhalgh T, Robert G, Macfarlane F, et al. Diffusion of innovations in service organizations: systematic review and recommendations. Milbank Q 2004;82:581-629.
- Colquhoun HL, Brehaut JC, Sales A, et al. A systematic review of the 14 use of theory in randomized controlled trials of audit and feedback. Implement Sci 2013;8:66.
- Dearing JW. Applying diffusion of innovation theory to intervention 15 development. Res Soc Work Pract 2009:19:503-18.
- Lien AS-Y, Jiang Y-D. Integration of diffusion of innovation theory into 16 diabetes care. J Diabetes Investig 2017;8:259-60.
- Rizan C, Phee J, Boardman C, et al. General surgeon's antibiotic 17 stewardship: climbing the rogers diffusion of innovation curveprospective cohort study. Int J Surg 2017;40:78-82.
- Mazotti L, Adams J, Peyser B, et al. Diffusion of innovation and 18 longitudinal integrated clerkships: results of the clerkship directors in internal medicine annual survey. Medical Teacher 2019;41:347-53.
- Benson T. Digital innovation evaluation: user perceptions of 19 innovation readiness, Digital confidence, innovation adoption, user experience and behaviour change. BMJ Health Care Inform 2019;26:e000018:0:.
- Spinnewijn L, Aarts J, Verschuur S, et al. Knowing what the patient 20 wants: a hospital ethnography studying physician culture in shared decision making in the Netherlands. BMJ Open 2020;10:e032921.

- Spinnewijn L. Bolte AC. Braat DDM. et al. Structurally collecting patient feedback on trainee skills: a pilot study in obstetrics and gynaecology. Patient Educ Couns 2022;105:1276-82.
- 22 Gale NK, Heath G, Cameron E, et al. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. BMC Med Res Methodol 2013;13:117.
- Pope C, Ziebland S, Mays N. Qualitative research in health care. 23 Analysing Qualitative Data BMJ 2000;320:114-6.
- Ritchie J, Spencer I. Qualitative Data Analysis for Applied Policy 24 Research. Analyzing Qualitative Data. London: Routledge, 1994:173-94.
- Aarts JWM, Dannenberg MD, Scalia P, et al. Development of an 25 adjective-selection measure evaluating clinicians' attitudes towards using patient decision AIDS: the ADOPT measure. Patient Educ Couns 2022:105:2785-92.
- 26 Allen M, Sage P. The SAGE Encyclopedia of Communication Research Methods. 1ST Ed. 1st edn. Los Angeles, California: SAGE, 2017.
- 27 Myren BJ, Hermens R, Koksma JJ, et al. Openness to new perspectives created by patient participation at the morbidity and mortality meeting. Patient Educ Couns 2021;104:343-51.
- 28 van den Berg M, van der Meij E, Bos AME, et al. Development and testing of a tailored online fertility preservation decision aid for female cancer patients. Cancer Med 2021;10:1576-88.
- 29 Elwyn G, Légaré F, van der Weijden T, et al. Arduous implementation: does the normalisation process model explain why it's so difficult to Embed decision support Technologies for patients in routine clinical practice. Implement Sci 2008;3:57.
- 30 Lloyd A, Joseph-Williams N, Edwards A, et al. "Patchy 'coherence': using normalization process theory to evaluate a multi-faceted shared decision making implementation program (MAGIC)". Implement Sci 2013;8:102.
- 31 May C, Finch T. Implementing, embedding, and integrating practices: an outline of normalization process theory. Sociology 2009:43:535-54
- 32 Kricheldorf HR. Getting It Right in Science and Medicine: Can Science Progress through Errors? Fallacies and Facts (1 Online Resource, XII, 285 Pages 8 Illustrations, 1 Illustrations in Color). 1st edn. Cham: Springer International Publishing : Imprint: Springer, 2016.
- 33 Bravo P, Härter M, McCafferty K, eds. "Special issue "International shared decision making conference 2022"" In: International Shared Decision Making Conference; 2022
- 34 Joseph-Williams N, Abhyankar P, Boland L, et al. What works in implementing patient decision aids in routine clinical settings? A rapid realist review and update from the International patient decision aid standards collaboration. Med Decis Making 2021:41:907-37.
- 35 Festinger L. A Theory of Cognitive Dissonance. Evanston, III: Row, 1957. Aronson E. The evolution of cognitive dissonance theory: a personal 36
- appraisal. Front Soc Psychol 2007;115-35. 37 Samuelson W, Zeckhauser R. Status quo bias in decision making. J Risk Uncertainty 1988;1:7-59.
- 38 Elliot AJ, Devine PG. On the motivational nature of cognitivedissonance - dissonance as psychological discomfort. J Person Soc Psych 1994;67:382-94.
- , AI training, and 39 Cooper J. Unlearning cognitive dissonance: toward an understanding of the development of dissonance. J Exp Soc Psy 1998;34:562-75.
- Murray JS, Lee J, Larson S, et al. "Requirements for implementing a 'iust culture' within healthcare organisations: an integrative review". BMJ Open Qual 2023;12:e002237.
- Wagner C, Smits M, Sorra J, et al. Assessing patient safety culture in 41 hospitals across countries. Int J Qual Health Care 2013;25:213-21.
- 42 Malik RF, Buljac-Samardžić M, Amajjar I, et al. Open organisational culture: what does it entail? Healthcare Stakeholders reaching consensus by means of a Delphi technique. BMJ Open 2021:11:e045515.
- Johnson DW. Constructive Controversy: Theory, Research, Practice. 43 New York: Cambridge University Press, 2015.
- Deutsch M. A theory of co-operation and competition. Human Relations 1949:2:129–52.
- Johnson DW. The Social Psychology of Education. New York: Holt, 1970. 45
- 46 Brigley S, Young Y, Littlejohns P, et al. Continuing education for medical professionals: a reflective model. Postgrad Med J 1997:73:23-6.
- 47 Mamede S, Schmidt HG. Correlates of reflective practice in medicine. Adv Health Sci Educ Theory Pract 2005;10:327-37.
- 48 Bourdieu P. Outline of a Theory of Practice. Cambridge ; New York: Cambridge University Press, 1977.
- 49 Giddens A. The Constitution of Society: Outline of the Theory of Structuration. Cambridge Cambridgeshire: Polity Press, 1984.