BMJ Open Protocol of the Nurses' Mental Health Study (NMHS): a nationwide hospital multicentre prospective cohort study

Meng Ning,^{1,2} Xuting Li,¹ Zengyu Chen,³ Jiaxin Yang,^{1,4} Qiang Yu,¹ Chongmei Huang ⁽¹⁾,⁵ Yamin Chen,^{1,2} Yusheng Tian,^{1,6} Ya-Min Li ⁽¹⁾,⁷ Shuivuan Xiao^{8,9}

ABSTRACT

To cite: Ning M, Li X, Chen Z, et al. Protocol of the Nurses' Mental Health Study (NMHS): a nationwide hospital multicentre prospective cohort study. BMJ Open 2025;15:e087507. doi:10.1136/ bmjopen-2024-087507

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (https://doi.org/10.1136/ bmjopen-2024-087507).

Received 11 April 2024 Accepted 30 January 2025



C Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to Professor Ya-Min Li: aminny@csu.edu.cn

Introduction Due to high workload, shift work and shortage of human resources, nurses are vulnerable to various mental health problems which are significantly related to their physical health, patient safety and healthcare guality. The long-term monitoring of nurses' mental health and identifying its risk factors are necessary. Accounting for more than one-fifth of global nurses. Chinese nurses have a demanding healthcare workload, especially in tertiary hospitals, yet there is no national cohort concerning their mental health and associated factors. This study (Nurses' Mental Health Study, NMHS) aims to investigate the mental health of Chinese nurses in tertiary hospitals and to identify the risk and protective factors.

Methods and analysis NMHS is a multicentre prospective cohort study which plans to recruit at least 30 000 registered nurses from 62 tertiary hospitals using the cluster sampling method. Baseline data collection will be implemented through online questionnaires after obtaining informed consent from each participant. Variables including sociodemographic information, work-related factors, health status, lifestyle habits, objective experiences, family and social relations, current affairs opinions and overall evaluation will be collected. Annual follow-up will be conducted via a guestionnaire link to monitor subsequent changes in nurses' mental health and related factors.

Patient and public involvement statement No patient and public involvement in the study design.

Ethics and dissemination This study was approved by the Institute Review Board from the ethical committee of The Second Xiangya Hospital of Central South University (E20230048) and NMHS will be performed in line with Helsinki Declarations. The results of NMHS will inform nursing research and practice by providing evidence to nurse managers, healthcare organisations and the government about the target mental health outcomes for early prevention and intervention.

Trial registration number This study has been registered at the Chinese Clinical Trail Registry (ID: ChiCTR2300072142, https://www.chictr.org.cn/showproj. html?proj=192676).

INTRODUCTION

Due to high workload, shift work and shortage of human resources, nurses are vulnerable to various mental health problems, such as depression, anxiety, insomnia, burnout and

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow To the best of our knowledge, this study will be the first cohort study of Chinese nurses concerning mental health, analysing the protective factors and risk factors from multiple dimensions including environmental factors, health behaviour and physical diseases which will provide target outcomes for interventions.
- ⇒ Participants will consist of nurses from all provinces in mainland China, ensuring the representativeness of the whole sample.
- \Rightarrow Self-reported questionnaires will be used to gain most of the data which may cause recall bias.
- \Rightarrow There may be selection bias since nurses who refuse to participate in the baseline survey or lost to follow-ups may have different characteristics compared with those who volunteer and stay in the cohort.
- \Rightarrow Hospitals with beds <2000 will not be included in this study which may cause sample bias.

data mining, Al training, post-traumatic stress disorder (PTSD).¹² This situation became worse due to the COVID-19 pandemic and the associated public health pressure, like the stress and workload from frontline healthcare.³ Meanwhile, nurses are also more susceptible to offensive behaviours <u>0</u> (eg, workplace violence) than other health professionals,⁴ resulting in higher stress.⁵ Mental health problems of nurses are significantly related to their poor physical health,⁶ decreased patient safety and healthcare quality.⁷ Therefore, the long-term monitoring **&** of nurses' mental health and identifying the risk factors of mental health problems would be crucial to their health and the healthcare system. A longitudinal study would be necessary to fully understand the mental health outcomes associated with environmental exposures and work-related factors over time. More than one-fifth of nurses work in China, taking on a non-negligible healthcare role worldwide. However, only two cohort studies

and

Protected by copyright, including for uses related to text

focused on Chinese nurses and none of them were concerned with mental health.

BACKGROUND

According to WHO, mental health refers to a state of mental well-being that enables people to cope with the stress of life, realise their abilities, learn well and work well and contribute to their community.⁸ Mental health is a continuum including positive and negative states. The positive aspect involves well-being and life satisfaction, while the negative element involves psychopathology, manifesting as mental health problems.⁹ Mental health problems of nurses include mental disorders and symptoms that affect nurses which may result from biological, economic and social factors.¹⁰ Compared with other healthcare professionals, nurses are generally more likely to suffer from mental health problems due to frequent night shifts and heavy workloads, especially after the COVID-19 pandemic.¹¹ Taking care of COVID-19 patients took immense physical and psychological toll on nurses. In addition, healthcare workers were at very high risk of infection during the pandemic which brought extra stress on nurses.¹²¹³ The prevalence of overall mental health problems among nurses was 25-39% during the COVID-19 pandemic, with the prevalence of anxiety, depression, PTSD and burnout being 25.8%, 30.3%, 39.3% and 31.5%, respectively.^{11 14 15} During the postpandemic period, they still suffer from mental health problems.¹⁶ In addition, nurses are vulnerable to compassion fatigue from repeated exposure to patients' suffering and pain which causes them emotional and physical exhaustion.¹⁷ Over 90% of nurses have experienced compassion fatigue and over 70% experienced moderate-to-severe levels.¹⁸

The mental health problems of nurses negatively affect their health and are associated with poor patient safety.^{6 19} It is reported that nurses' mental health problems may increase the risk of chronic diseases (eg. coronary heart disease, hypertension), tumours (eg, lung/ colorectal cancer) and other diseases (eg, systemic lupus erythematosus, Crohn's disease).^{6 20–23} Meanwhile, nurses with mental health problems are more likely to experience adverse events than others which threatens patient safety.^{15 19} As for organisations, mental health problems among nurses are associated with increased absences (eg, sick leave and resignation),²⁴ which would exacerbate the shortage of human resources. Therefore, understanding the risk factors and protective factors of nurses' mental health problems would be urgent due to the adverse influence on nurses' health and the healthcare system.

The cohort study design is an appropriate method to clarify protective and risk factors of mental health in nurses. Several cohort studies have been conducted in developed countries, such as the Nurses' Health Study conducted in America,²⁵ the Japan Nurses' Health Study²⁶ and the Danish Nurse Cohort.²⁷ In these cohorts, their primary outcome was somatic illnesses, however,

depression, anxiety, PTSD and other mental health disorders were included as potential risk factors.⁶ Evidence has demonstrated that mental health problems (such as depression, anxiety and PTSD) are influenced by multiple factors including individual factors, behaviours, community and policy.^{6 28-30} However, previous studies of nurses' mental health mainly relied on cross-sectional evidence.^{31 32} Besides, mental health problems in nurses from different countries are diverse and findings from cohort studies in other countries are not entirely applicable to Chinese nurses. The current situation of mental health and its influencing factors on Chinese nurses remained little known.

As the world's second most populous developing country, China boasts a nurse workforce exceeding five 8 million as of 2021,³³ more than one-fifth of the global nurses.³⁴ However, the density of nurses (per 10000 pop) is 33.05 in China, far lower than in developed countries like Australia (137.99), the USA (124.68), Canada (102.70) and the UK (86.86).³⁵ Consequently, Chinese nurses grapple with a demanding healthcare workload, especially those in tertiary hospitals who provide care for patients with complex or rare diseases.³⁶ In addition, the beds and the utilisation rate of beds are higher in tertiary hospitals and over 50% of inpatients and outpatients are hospitals and over 50% of inpatients and outpatients are $\frac{1}{2}$ in these hospitals.³⁷ Thus, nurses in tertiary hospitals $\frac{1}{2}$ have heavier workloads and pressure, making them even g more vulnerable to mental health problems. Up to now, there have been only two cohorts of nurses in China, the Chinese Nurses' Health Study (The TARGET study)³⁸ and the National Nurse Health Study,³⁹ but both of them only involved nurses in one hospital or one province which data mining, Al training, and could not represent the national circumstances. Therefore, it is necessary to conduct a national multicentred large cohort study of nurses' mental health.

STUDY OBJECTIVES

This study (Nurses' Mental Health Study, NMHS) aims to investigate the mental health of Chinese nurses in tertiary hospitals and to identify the risk factors and protective factors for nurses' mental health, by establishing a largescale cohort of nurses across China. We will include nurses from 62 tertiary hospitals in 31 provinces, indicating good representativeness.

METHODS AND ANALYSIS Study design and theoretical model Study design

The NMHS is a nationwide multicentre prospective cohort study and its sample can represent nurses in tertiary hospitals in mainland China. This study will include a baseline survey and longitudinal follow-ups and has been registered on the Chinese Clinical Trial Registry (ChiCTR2300072142, https://www.chictr.org.cn/showproj.html?proj=192676). Figure 1 shows the procedure and overall flow of NMHS. The recruitment and baseline



Figure 1 The procedure and overall flow of the Nurses' Mental Health Study (NMHS).

survey were conducted between December 2023 and January 2024.

Theoretical model

The potential influencing factors of nurses' mental health will be identified according to the Ecosocial Model by Glass and McAtee⁴⁰ which directs attention to the biology–environmental interactions on health and health behaviours across the lifespan. Glass and McAtee proposed time axis and hierarchical axis to demonstrate the impact on health outcomes, from multilevel and different life stage.⁴⁰ The time axis represents one's life span or a certain period, emphasising the cumulative effects of early exposure. The hierarchical axis demonstrates a multilevel nested system

from genes to cells and organs to human behaviour and the environment.⁴⁰ The theory framework of this study is shown in figure 2.

Study population

Sample size

To detect any events with a proportion as low as 1% with $\pm 0.1\%$ precision (ie, it offered a relative precision of $\pm 10\%$ with a two-side 95% confidence level),⁴¹ this study will need to include at least 30 000 nurses.

Sampling criteria

Nurses working in tertiary hospitals in mainland China will be recruited. Inclusion criteria are as follows: (1) at



Figure 2 Theory framework for risk factors of nurses' mental health problems, adapted from Ecosocial Model.⁴⁰

least 18 years old; (2) working full-time as a registered nurse in a tertiary hospital. Exclusion criteria are (1) training nurses (eg, student nurses, nurses in preregistration or standardised training programmes); (2) nurses from other hospitals who are in short-term programmes of specialised certifications (eg, intensive care unit nursing specialists). The determinant factors for mental health are different between student nurses and full-time nurses due to their distinct roles and pressure.⁴² Nurses in short-term programmes always come from non-tertiary hospitals which are not eligible for inclusion in hospitals. For these reasons, they will be excluded from this study.

Sampling method

A cluster sampling method will be used to recruit participants. First, two tertiary hospitals will be selected from each of the 31 province-level administrative regions (hereafter, 'provinces') in mainland China, making up 62 primary sampling sites. Dividing the required sample size (30 000) equally into 62 hospitals in 31 provinces, the baseline recruitment will need to recruit at least 630 nurses in each hospital considering a potential nonresponse rate of 30%. According to the Guidelines for Medical Service Capacity of Tertiary General Hospitals (2016 Edition) proposed by the National Health and Family Planning Commission of the People's Republic of China,⁴³ the minimum nurse-bed ratio in tertiary general hospitals is 0.4, which means targeted hospitals in this study will need to have 1575 beds. Meanwhile, given that the biggest hospitals in some remote provinces (eg, Qinghai and Tibet) do not have so many beds, we conservatively decided to choose hospitals with 2000 beds in most provinces, while in others, the target hospitals will be chosen from the top five tertiary hospitals ranked by bed numbers with the same criteria. Thus, the inclusion criteria of hospitals are: (1) having at least 2000 beds

Protected by copyright, including for uses except for hospitals in remote provinces; (2) general hospitals capable of dealing with a range of disease types similar to other tertiary hospitals in the provinces. Hospirelated tals unwilling to participate or quit during follow-ups will be excluded.

With hospitals selected, all registered nurses in those **ö** tex hospitals who met the inclusion and exclusion criteria will be invited to participate the study. Since the range of beds among all tertiary hospitals in mainland China is 500~12000, the number of nurses in each primary site may vary from 200 to 4800 according to the lowest nurse-bed ratio of 0.4:1.

Procedure

Baseline recruitment

I training, Baseline recruitment and data collection will be performed through online questionnaires via the Wenjuanxing platform which has been widely used in multicentre studies.44 45 Variables including sociodemographic information, work-related factors, health status, <u>0</u> lifestyle habits, objective experiences, family and social relations, current affairs opinions, overall evaluation and identification information will be collected. Details of the variables are shown in table 1 and the questionnaire is presented in online supplemental table S2.

Directors of nursing departments in each potential primary study site will be invited first through Chinese 8 Nursing Association. Once they agree, their hospital will be officially included as the primary study site and all nurses in the hospital will be recognised as potential participants. Based on the basic hospital information provided by directors of nursing departments, a unique QR code (Quick Response Code) for each hospital, which links to informed consent and online questionnaires will be generated and distributed by two trained researchers. The QR codes will be sent by head nurses to all nursing

and

data mining, Al

		Time points				
Construction	Contents	Baseline	1y	2у	Зу	4y
Sociodemographic Information (11)	Sex, Age, Ethnicity, Height, Weight, Years of Work Experience, Professional Title, Position, Educational Level, Marital Status, Spouse's Occupation, Pregnancy Status, Childbirth Status and Intention, Religious Belief	×	×	×	×	×
Work-related information (18)	Main Work, Department, Weekly Working Hours, Shifts, Nurse-to-Patient Ratio, Paper Work, Work-Life Balance, Occupational Events	×	×	×	×	×
	Workplace Violence	x	×	×	×	×
	Work Atmosphere, Sense of Meaning in Work, Career Development Prospects, Willingness to Children being a Nurse, Intention to Leave	×	×	×	×	×
	Nursing Leadership	×	×	×	×	×
Health status (25)	Fatigue, Loneliness, Psychological Resilience and Burnout, PTSD, Compassion Fatigue	×	×	×	×	×
	Suicidal and Self-harming Behaviour	×	×	×	×	×
	Stress	×	×	×	×	×
	Depression Symptoms, Depressive Traits and Postpartum Depression	×	×	×	×	×
	Anxiety Symptoms	×	×	×	×	×
	Obsession	×	×	×	×	×
	Sleep Disorder	×	×	×	×	×
	Current/Past History and Family History of Mental Health Problems, Utilisation of Mental Health Services, Psychological Care Organisations and Participation in Mental Health Promotion Activities	×	×	×	×	×
	Past/ Present Illness History, Pain, Hospitalisation History and Constipation	×	×	×	×	×
ifestyle habits (6)	Smoking, Drinking, Milk Tea, Exercise, Pets, Handedness	×	×	×	×	×
bjective xperiences (3)	Adverse Childhood Experience, Life Events	×	×	×	×	×
amily and social elations (9)	Social Support, Work Support, Intimate Partner Violence	×	×	×	×	×
Current affairs pinions (5)	Blood Donation, Organ Donation, Nurses' Part-Time Jobs, Nurse Prescriptive Authority, Euthanasia	×				
verall evaluation 2)	Overall Evaluation and Subjective Well-being	×	×	×	×	×
tentification formation (4)	Contact Number, Last Four Digits of ID Card, Email	×	×	×	×	×
	stress disorder: v vear					

units in selected hospitals through their wechat group. Eligible nurses in these departments can participate in this study voluntarily. Once nurses log in, they will first be provided with an online informed consent claiming with the following information: (1) this is a national multicentre cohort study concerning mental health and associated factors; (2) after baseline recruitment, there will be annual follow-ups in the next 5 years or longer; (3) the participation is voluntary and they could refuse or quit at any time and could withdraw their data before data analyses; (4) data will only be used for research purposes and will be kept confidentially; (5) personal information (eg, email address, phone numbers) will be carefully stored and only used for follow-up invitation; (6) participating

to join. If the questionnaire is not completed, it is not 8 considered consent to participate and the data for the uncompleted questionnaire will not be saved. Baseline recruitment in each hospital will be finished within 2months and the whole baseline recruitment in 62 primary study sites will be finished within 3 months.

Follow-up

Participants will receive annual follow-up online questionnaires through a link. The initial duration of follow-ups

is 5 years and additional funding applications will be performed to transform this study into a continuing project. Hospitals unwilling to participate or quit during follow-ups will be excluded. However, nurses in those hospitals can still join in follow-ups with online questionnaires sent to them through a link.

Outcomes

The primary outcomes are the mental health status of nurses, including both positive aspects and mental health problems. Specifically, subjective well-being, resilience and life satisfaction will be the three main indicators of positive aspects and depression, anxiety, obsessive symptom/disorder, PTSD, compassion fatigue, self-harm and suicidal behaviours will be the main outcomes for mental health problems. The scales for different mental health variables will be passed to all the participants.

Similar to previous studies,⁶ symptom burden or incidents of mental disorders will be measured using a Boolean OR operator approach (self-reported cliniciandiagnosed disorders or severe symptoms by published cutpoints, such as Patient Health Questionnaire-9⁴⁶). All outcome indicators will be identified using sophisticated self-reported questionnaires and scales. Specifically, mental disorders will be recorded once participants reported newly diagnosed by a doctor during follow-ups. The details of measures and diagnostic criteria for primary outcomes are shown in online supplemental table S1.

Data management and statistical analysis

Data management and usage

All collected data will be uploaded and stored on an encrypted big data platform by the data manager using a dual entry system. Two independent researchers will perform data cleaning before data analyses which includes three steps. First, the duplicated questionnaires will be removed. If the birth month, phone number and the last four digits of the ID number are all the same in two questionnaires, the one submitted first will be retained. Next, the outlier of the continuous variable will be removed, including numbers less than P25-3IQR and over P75+3IQR.⁴⁷ Finally, logical conflict checks will be employed. For example, if a participant reported years of work experience more than their age, it will be considered a logical conflict. Questionnaires with over two such conflicts will be excluded. No missing data in this study because the questionnaire can only be submitted after all the items are completed.

After concealing the personal information of participants and basic data cleaning, data per hospital will be opened to themselves. Researchers from participating hospitals can also apply for access to the overall data by submitting study proposals which will be evaluated by professionals and the research team. They are also welcomed and encouraged to make suggestions for the project in light of the local reality.

<page-header><page-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header>

BMJ Open: first published as 10.1136/bmjopen-2024-087507 on 11 February 2025. Downloaded from http://bmjopen.bmj.com/ on June 12, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

project consultants and technical staff from the lead unit to ensure the quality of data collection.

Handling of adverse events and emergencies

- 1. Assessment of risk for suicide and self-harm behaviour: If nurses are at high risk for anxiety, depression, selfharm and suicide assessed by the self-reported questionnaire, the page after completing the questionnaire will show access to professional mental health services. In addition, the research team will provide them with a professional self-care manual through email and the details can be found in online supplemental table S3. They will also be recommended for further diagnosis and treatment in specialised hospitals or institutions.
- 2. Each project site should record other unintended or adverse events (eg, strong issues about this study, unexpected death of nurses by suicide or other accidents and turnover of nurses) and report to the lead unit timely. The study team will revise the operation manual based on feedback.

ETHICS AND DISSEMINATION

This study obtained ethics approval by the Institute Review Board from the ethical committee of The Second Xiangya Hospital of Central South University (Ethics approval number: E20230048). Before data collection, the purpose and content of the study will be fully explained to the participants and informed consent will be obtained from each participant. Participants in the baseline survey and follow-up studies will be offered ¥1~100 (US\$0.13701~13.701) randomly as thanks for completing questionnaires through online WeChat payment. This study will be performed in line with the protocol, Chinese law and Helsinki Declarations. Participants are free to join or refuse and they will be formally included only after giving fully informed consent. Participants will have the right to quit the study at any time, yet their data can be withdrawn only before data analyses which will not harm their health or career. The study reported in this paper will not involve the use of any animal or human tissue. All data will be kept anonymously and only used for this study. The findings of NMHS will be disseminated through peer-reviewed journals and national and international conferences.

Patient and public involvement statement

Patients or the public will not be involved in the design, or conduct, or reporting, or dissemination plans of our research.

DISCUSSION

To the best of our knowledge, this study may be the first to establish a large cohort of Chinese nurses concerning mental health. We will analyse the protective factors and risk factors including policy, health behaviour, workrelated factors and somatic disorders, and construct a risk prediction model of nurses' mental disorders. The results of our study will offer a long-term observation and an accurate understanding of the mental health trajectories of nurses over time, providing researchers and nurse managers with significant mental health outcomes which will be a scientific basis for implementing targeted interventions. Moreover, our study will provide scientific evidence for the government to formulate macro prevention and control policies for nurses' mental health problems and ultimately improve the mental health of nurses and improve nursing quality and patient safety.

Current status of this study

The baseline recruitment was conducted between December 2023 and January 2024. The follow-up proce- or dure will start one year after.

Author affiliations

¹Clinical Nursing Teaching and Research Section, The Second Xiangya Hospital of Central South University, Changsha, China

²Xiangya School of Nursing, Central South University, Changsha, China
 ³School of Nursing, University of Washington-Seattle, Seattle, Washington, USA
 ⁴Computer Science and Engineering, Central South University, Changsha, China
 ⁵School of Nursing at Ningxia Medical University, Yinchuan, China
 ⁶Department of Psychiatry, National Clinical Research Center for Mental Disorders,

the Second Xiangya Hospital of Central South University, Changsha, China ⁷Hunan Provincial People's Hospital, The first-affiliated hospital of Hunan nominal university, Changsha, China

⁸Mental Health Center, Xiangya Hospital Central South University, Changsha, China
⁹Department of Social Medicine and Health Management, Central South University, Changsha, China

Contributors All authors contributed to the study conceptualisation. MN: Writing original draft; Writing—review and editing; Visualisation. XL: Writing—review and editing; Visualisation. ZC, JY, QY, CH, YC, YT: Writing—review and editing. Y-ML: Methodology; Funding acquisition; Resources; Supervision; Validation; Project administration. SX: Methodology. Y-ML is responsible for the overall content as the guarantor.

Funding This work was supported by the STI2030-Major Projects (2021ZD0200700) and the Scientific Research Project of Health Commission of Hunan Province (W20243200). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

Author note Co-first authors: Meng Ning and Xuting Li contributed equally to this work.

ORCID iDs

Chongmei Huang http://orcid.org/0000-0001-6794-0724 Ya-Min Li http://orcid.org/0000-0002-8396-5208

REFERENCES

- 1 Zheng R, Zhou Y, Fu Y, et al. Prevalence and associated factors of depression and anxiety among nurses during the outbreak of COVID-19 in China: A cross-sectional study. Int J Nurs Stud 2021;114:S0020-7489(20)30295-9.
- 2 Rosa D, Terzoni S, Dellafiore F, et al. Systematic review of shift work and nurses' health. Occup Med (Lond) 2019;69:237–43.
- 3 Cuccia AF, Peterson C, Melnyk BM, et al. Trends in mental health indicators among nurses participating in healthy nurse, healthy nation from 2017 to 2021. Worldviews Evid Based Nurs 2022;19:352–8.
- 4 Liu J, Gan Y, Jiang H, *et al.* Prevalence of workplace violence against healthcare workers: a systematic review and meta-analysis. *Occup Environ Med* 2019;76:927–37.
- 5 Yao X, Shao J, Wang L, *et al.* Does workplace violence, empathy, and communication influence occupational stress among mental health nurses? *Int J Ment Health Nurs* 2021;30:177–88.
- 6 Trudel-Fitzgerald C, Chen Y, Singh A, *et al.* Psychiatric, Psychological, and Social Determinants of Health in the Nurses' Health Study Cohorts. *Am J Public Health* 2016;106:1644–9.
- 7 Baylina P, Barros C, Fonte C, *et al.* Healthcare Workers: Occupational Health Promotion and Patient Safety. *J Med Syst* 2018;42:159.
- 8 WHO. Mental health, 2022. Available: https://www.who.int/newsroom/fact-sheets/detail/mental-health-strengthening-our-response
- 9 Magalhães E. Dual-factor Models of Mental Health: A Systematic Review of Empirical Evidence. *Psychosoc Interv* 2024;33:89–102.
- 10 Rugulies R, Aust B, Greiner BA, et al. Work-related causes of mental health conditions and interventions for their improvement in workplaces. Lancet 2023;402:1368–81.
- 11 Pappa S, Ntella V, Giannakas T, et al. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. Brain Behav Immun 2020;88:901–7.
- 12 International Council of Nurses. ICN COVID-2019 Update: Data on infected healthcare workers and access to protective equipment vital in prevention of the virus, Available: https://www.icn.ch/news/icncovid-2019-update-data-infected-healthcare-workers-and-accessprotective-equipment-vital
- 13 Sabetian G, Moghadami M, Hashemizadeh Fard Haghighi L, *et al.* COVID-19 infection among healthcare workers: a cross-sectional study in southwest Iran. *Virol J* 2021;18:58.
- 14 Chen R, Sun C, Chen J-J, et al. A Large-Scale Survey on Trauma, Burnout, and Posttraumatic Growth among Nurses during the COVID-19 Pandemic. Int J Ment Health Nurs 2021;30:102–16.
- 15 Kakemam E, Chegini Z, Rouhi A, et al. Burnout and its relationship to self-reported quality of patient care and adverse events during COVID-19: A cross-sectional online survey among nurses. J Nurs Manag 2021;29:1974–82.
- 16 de Vroege L, van den Broek A. Post-pandemic self-reported mental health of mental healthcare professionals in the Netherlands compared to during the pandemic - an online longitudinal follow-up study. *Front Public Health* 2023;11:1221427.
- 17 Peters E. Compassion fatigue in nursing: A concept analysis. *Nurs Forum* 2018;53:466–80.
- 18 Xie W, Chen L, Feng F, et al. The prevalence of compassion satisfaction and compassion fatigue among nurses: A systematic review and meta-analysis. Int J Nurs Stud 2021;120:S0020-7489(21)00118-8.
- 19 Kakemam E, Kalhor R, Khakdel Z, et al. Occupational stress and cognitive failure of nurses and associations with self-reported adverse events: A national cross-sectional survey. J Adv Nurs 2019;75:3609–18.
- 20 Choi MY, Malspeis S, Sparks JA, *et al*. Association of Sleep Deprivation and the Risk of Developing Systemic Lupus Erythematosus Among Women. *Arthritis Care Res (Hoboken)* 2023;75:1206–12.
- 21 Ananthakrishnan AN, Khalili H, Pan A, *et al.* Association between depressive symptoms and incidence of Crohn's disease and ulcerative colitis: results from the Nurses' Health Study. *Clin Gastroenterol Hepatol* 2013;11:57–62.
- 22 Trudel-Fitzgerald C, Zevon ES, Kawachi I, et al. Depression, smoking, and lung cancer risk over 24 years among women. *Psychol Med* 2022;52:2510–9.

- 23 Trudel-Fitzgerald C, Tworoger SS, Zhang X, et al. Anxiety, Depression, and Colorectal Cancer Survival: Results from Two Prospective Cohorts. J Clin Med 2020;9:3174.
- 24 Korošec D, Vrbnjak D, Štiglic G. Prevalence of mental disorders and high rates of absenteeism from work among healthcare professionals in Slovenia: a retrospective study. *BMJ Open* 2023;13:e075718.
- 25 Bao Y, Bertoia ML, Lenart EB, et al. Origin, Methods, and Evolution of the Three Nurses' Health Studies. Am J Public Health 2016;106:1573–81.
- 26 Hayashi K, Mizunuma H, Fujita T, et al. Design of the Japan Nurses' Health Study: a prospective occupational cohort study of women's health in Japan. Ind Health 2007;45:679–86.
- 27 Hundrup YA, Simonsen MK, Jørgensen T, *et al.* Cohort profile: the Danish nurse cohort. *Int J Epidemiol* 2012;41:1241–7.
- 28 Weitzer J, Trudel-Fitzgerald C, Okereke OI, et al. Dispositional optimism and depression risk in older women in the Nurses' Health Study: a prospective cohort study. *Eur J Epidemiol* 2022;37:283–94.
- 29 Qureshi F, Stampfer M, Kubzansky LD, et al. Prospective associations between coffee consumption and psychological wellbeing. PLoS One 2022;17:e0267500.
- 30 Chen Y, Harris SK, Worthington EL Jr, et al. Religiously or Spiritually-Motivated Forgiveness and Subsequent Health and Well-Being among Young Adults: An Outcome-Wide Analysis. J Posit Psychol 2019;14:649–58.
- 31 Matricciani L, Clarke J, Wiley S, et al. Sleep of Nurses: A Comprehensive Scoping Review. J Adv Nurs 2024.
- 32 Uddin H, Hasan MK, Cuartas-Alvarez T, et al. Effects of mass casualty incidents on anxiety, depression, and post-traumatic stress disorder among doctors and nurses: a systematic review. Public Health (Fairfax) 2024;234:132–42.
- 33 National Health Commission. China health statistics yearbook (2022). Peking Unionmedical College Press, 2022.
- 34 WHO. Global strategic directions for nursing and midwifery, 2022. Available: https://iris.who.int/bitstream/handle/10665/344562/ 9789240033863-eng.pdf?sequence=1
- 35 Platform, W.N.D. National Health Workforce Accounts Data Portal, 2022. Available: https://apps.who.int/nhwaportal/Home/Index
- 36 Commission, T.N.H. National Health Commission to consolidate diagnosis and treatment systems, Available: https://english.www.gov. cn/news/202411/29/content_WS67491b13c6d0868f4e8ed837.html
- 37 China, N.H.C.o.t.P.s.R.o. China Health Statistical Yearbook 2022, Available: http://www.nhc.gov.cn/mohwsbwstjxxzx/tjtjnj/202305/6ef6 8aac6bd14c1eb9375e01a0faa1fb/files/b05b3d958fc546d98261d165 cea4adba.pdf#page=405.08
- 38 Lv X, Cao Y, Li Y, et al. The TARGET Nurses' health cohort study protocol: Towards a revolution in getting nurses' health ticked. J Adv Nurs 2022;78:1815–23.
- 39 Zhuo L, Zhang H, Geng R, et al. Protocol for the National Nurse Health Study (NNHS): a web-based ambispective cohort study. BMJ Open 2021;11:e049958.
- 40 Glass TA, McAtee MJ. Behavioral science at the crossroads in public health: Extending horizons, envisioning the future. Soc Sci Med 2006;62:1650–71.
- 41 Sawaengdee K, Tangcharoensathien V, Theerawit T, et al. Thai nurse cohort study: cohort profiles and key findings. BMC Nurs 2016;15:10.
- 42 Visier-Alfonso ME, Sarabia-Cobo C, Cobo-Cuenca AI, *et al.* Stress, mental health, and protective factors in nursing students: An observational study. *Nurse Educ Today* 2024;139:S0260-6917(24)00168-0.
- 43 National Health and Family Planning Commission of the People's Republic of China. Guidelines for Medical Service Capacity of Tertiary General Hospitals, 2016. Available: http://www.nhc.gov.cn/ ewebeditor/uploadfile/2016/10/20161020101032482.pdf
- 44 Huang C, Li L, Zhang T, *et al.* Breastfeeding education in Chinese hospitals: A cross-sectional study. *Int J Nurs Stud* 2022;133:S0020-7489(22)00139-0.
- 45 Han T, Wang D, Xie W, *et al.* Obstetricians' Attitudes Toward the Treatment of Extremely Preterm Infants in China. *JAMA Netw Open* 2022;5:e2233511.
- 46 Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. J Gen Intern Med 2001;16:606–13.
- 47 Seo S. A review and comparison of methods for detecting outliers in univariate data sets. University of Pittsburgh, 2006.
- 48 Battineni G, Sagaro GG, Chinatalapudi N, et al. Applications of Machine Learning Predictive Models in the Chronic Disease Diagnosis. J Pers Med 2020;10:21.

for uses related to text

and data mining,

Al training, and similar technologies

Protected by copyright, including