




BMJ Open Incidence and risk factors for glaucoma and its clinical, mental health and economic impact in an elderly population: a longitudinal study

Catherine Jan ^{1,2} Xin Jin ^{3,4} Mengtian Kang,⁵ Jiahao Liu,^{1,2} Wenyi Hu,^{1,2} Ruiye Chen,^{1,2} Li Li,^{1,2} Mingguang He,^{1,6} Nathan Congdon ^{7,8,9}

To cite: Jan C, Jin X, Kang M, *et al.* Incidence and risk factors for glaucoma and its clinical, mental health and economic impact in an elderly population: a longitudinal study. *BMJ Open* 2025;**15**:e096879. doi:10.1136/bmjopen-2024-096879

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-096879>).

CJ and XJ are joint first authors.

Received 20 November 2024
Accepted 01 April 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Nathan Congdon;
n.congdon@qub.ac.uk and
Dr Catherine Jan;
catherine.jan@student.unimelb.edu.au

ABSTRACT

Objectives To investigate the incidence and determinants of glaucoma in an elderly Chinese population, and clinical, mental health and economic impacts.

Design This nationally representative, longitudinal study assessed self-reported 6-year (from 2011 to 2018) incident glaucoma diagnosis by a physician and measured biological, clinical and socioeconomic participant characteristics at baseline and endline.

Setting In the first stage, 150 county-level units from across China were randomly selected with a probability-proportional-to-size sampling technique from a frame containing all county-level units nationwide. The sample was stratified by region and within region by urban district or rural county and per capita gross domestic product. The final sample of 150 counties included 30 out of 31 provinces and autonomous regions in China.

Participants Consenting, community-dwelling Chinese persons aged 50 years and older.

Primary and secondary outcome measures Incident glaucoma incidence (primary), factors associated with incident glaucoma (secondary), impact of glaucoma (secondary).

Results Among 9973 individuals, 3.4% reported a glaucoma diagnosis between 2011 and 2018; Central China had the highest incidence (3.95%) and Eastern China the lowest (2.64%) between 2011 and 2018. Those diagnosed with glaucoma during 2011 and 2018 were of older age (beta coefficient: 0.050, 95% CI: 0.001, 0.001, $p<0.001$), had higher prevalence of diabetes (beta coefficient: 0.049, 95% CI: 0.028, 0.032, $p<0.001$), hypertension (beta coefficient: 0.019, 95% CI: 0.006, 0.008, $p<0.001$), smoking (beta coefficient: 0.029, 95% CI: 0.004, 0.020, $p=0.004$), alcohol consumption (beta coefficient: 0.026, 95% CI: 0.002, 0.017, $p<0.009$) and illiteracy (beta coefficient: -0.057 , 95% CI: -0.030 , -0.015 , $p<0.001$). Logistic regression models showed significant association between incidence of the following characteristics and baseline glaucoma: poor self-reported distance vision (beta coefficient: 1.106, 95% CI: 0.701, 1.511, $p<0.001$), having hypertension (beta coefficient: 0.545, 95% CI: 0.496, 0.593, $p<0.001$), having diabetes (beta coefficient: 0.388, 95% CI: 0.326, 0.449, $p<0.001$), not having obesity (beta coefficient: -0.184 , 95% CI: -0.239 , -0.129 , $p<0.001$) and lower mean value of health utility score of residents' quality of life (beta coefficient: -0.040 , 95% CI: -0.006 , 0.776, $p<0.001$).

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The study used a large, nationally representative sample from most of China's administrative units, selected using standardised protocols.
- ⇒ Study personnel were trained uniformly to ensure consistent implementation of procedures.
- ⇒ The self-reported nature of glaucoma diagnosis may have introduced underreporting or misdiagnosis, though prior evidence suggests reasonable accuracy in self-reports.
- ⇒ The results may have been influenced by the high proportion (42.2%) of participants who did not attend follow-up examinations or provide information on their glaucoma history.

Conclusions Glaucoma incidence rate varies among geographical regions in China. Several risk factors for incident glaucoma were identified. In addition, glaucoma was found to be associated with multiple physical and psychosocial outcomes. Targeted public health strategies are needed, emphasising early detection and better vision care, to alleviate the burden of glaucoma and improve well-being.

INTRODUCTION

Glaucoma, the world's leading cause of irreversible blindness,¹ poses a significant public health challenge, especially in a rapidly ageing population such as China's. In China, the burden of glaucoma is increasingly recognised as a substantial health issue due to its impact on vision-related quality of life and the economy.^{2 3} Despite this recognition, there are no comprehensive nationwide studies exploring the incidence rates and determinants associated with incident glaucoma in mainland China.

China is a large and populous country containing many ethnicities. Understanding the epidemiology of glaucoma incidence and its associated risk factors on a national scale is crucial for developing effective preventive

strategies, early detection methods and targeted interventions. Glaucoma incidence studies from mainland China are rare.^{4–6} One study reported the 5-year cumulative incidence of primary open-angle glaucoma was 1.3% in Bai Chinese aged 55 years and above,⁴ while another study reported the 5-year cumulative incidence of primary glaucoma was 1.6% in rural northern China among people aged 30 years and older.⁵ Previous localised studies have highlighted specific risk factors contributing to glaucoma incidence in certain regions of China, such as increased age,^{4,5} elevated intraocular pressure,^{4,5} lower educational level⁴ and the presence of myopia,⁴ yet a cohesive and comprehensive nationwide analysis is essential to capture the broader landscape.

The current report aims to address this gap by providing a comprehensive assessment of the national glaucoma incidence rates in China and exploring the determinants, risk factors and impact associated with incident glaucoma across diverse geographical regions and demographic groups. This project employed a national longitudinal survey conducted in 30 out of 31 Chinese provinces and autonomous regions (except Tibet, though the study did include one Tibetan county), representative of the adult population 45 years and older between 2011 and 2018.

The aim of this research is to offer insights into the epidemiology of glaucoma in China, aiding policy-makers, healthcare providers and public health experts in formulating targeted preventive strategies, allocating resources effectively and devising tailored interventions to mitigate the burden of glaucoma and preserve ocular health nationally.

METHODS

This study uses data drawn from the same original longitudinal cohort as our previous publication on cataract surgery incidence,⁷ including the same population selection, study settings and statistical analysis framework. As such, some overlap in the Method sections is present due to the consistent methodology applied across both papers. However, the two studies address entirely distinct research questions: while the previous paper focused on the incidence and determinants of cataract surgery uptake—a healthcare service intervention—this manuscript investigates the incidence of glaucoma, a different disease entity and explores its associated clinical, psychosocial and economic consequences.

Study population

The China Health and Retirement Longitudinal Study (CHARLS) is a nationally representative longitudinal survey among Chinese persons aged 45 years and older (note for this particular paper, we used data from people aged 50 years and above because glaucoma is an age-related disease and this is the most common threshold reported by literature), that includes assessments of biological, social and economic conditions. We used data from the baseline in-person study conducted in 2011

(‘wave 1’) and an in-person follow-up conducted in 2018 (‘wave 4’). Main respondents and spouses in the baseline survey are followed throughout the life of CHARLS, or until they die.⁸ Detailed information on the methodology of CHARLS has been provided elsewhere.⁸ Among 17250 eligible participants at baseline who provided information on glaucoma status and had not been diagnosed with glaucoma in 2011, 9973 (57.8%) attended the 7-year follow-up and provided glaucoma information in 2018. The identical protocols for collection of the data used in this paper were followed during the baseline and follow-up studies.

The participant cohort was chosen through multi-stage probability sampling. In the first stage, 150 county-level units were randomly selected with a probability-proportional-to-size (PPS) sampling technique from a frame containing all county-level units in the country, with the exception of the Tibet Autonomous Region. The sample was stratified by region and within region by urban district or rural county and per capita gross domestic product (GDP). The final sample of 150 counties included 28 of China’s 31 administrative units (provinces and autonomous regions). Our sample used the lowest level of government organisation, consisting of administrative villages (cun) in rural areas and neighbourhoods (shequ or juweihui) in urban areas, as primary sampling units (PSUs). We selected three PSUs within each county-level unit, using PPS sampling (more details are reported elsewhere).⁸

Variables

Glaucoma incidence in this paper refers to the proportion of the cohort population aged 50 years and older who reported having a glaucoma diagnosis by a physician in at least one eye between 2011 and 2018, among those who had not been diagnosed in either eye at baseline. This was determined by participants’ response to the question “Have you been diagnosed with glaucoma by a physician before?”, comparing answers between the previous and present round. Age and gender were self-reported variables taken from the baseline 2011 survey.

Rural or urban place of residence was obtained from self-reported hukou. Hukou is a Chinese internal passport system, designed to regulate the movement of rural residents into urban areas; migrants with a rural hukou do not have access to local healthcare as do local urban residents. Hukou also determines the type of public health insurance a person has access to.

Three comorbidities linked with glaucoma were extracted from the CHARLS dataset. Hypertension was defined as either self-report of a physician’s diagnosis, or presence of objective biomarkers (systolic pressure ≥ 140 and/or diastolic pressure ≥ 90 mm Hg) during direct physical examination. Diabetes was also defined by self-report of prior diagnosis by a doctor or the presence of a biomarker (hemoglobin A1c, also known as glycated hemoglobin; HbA1c value $\geq 6.5\%$) on examination. Depression symptoms were measured as a continuous

variable, using the Centre for Epidemiologic Studies Depression Scale-10,⁹ with total score ranging from 0 (minimal symptoms) to 30 (most symptomatic). Other covariates included in this study are household expenditure, representative of economic status, and EuroQol five-dimension questionnaire or EQ-5D health utility scores, an indicator of quality of life, and level of attained education.

Patient and public involvement

None.

Statistical analysis

All statistical analyses were performed using Stata V.14.0 (StataCorp). The t-test was used to compare continuous variables, while Pearson χ^2 or Fisher's exact tests were used for the comparison of categorical data. To assess the accuracy of household expenditure data from the sample as an index of income in the province, a scatterplot graph was made comparing household expenditure to provincial GDP data. The 7-year incidence of newly diagnosed glaucoma stratified by age and gender was calculated. Age was defined in our analysis as the age at baseline.

Univariate and multivariate logistic regression models were used to test for associations with incident glaucoma over the 7-year follow-up period. We ran regression analyses including the whole sample of eligible participants, aged 50 years and above. Variables in univariate models with $p < 0.05$ were entered into the multivariate models. In addition, univariate logistic regression models were applied to investigate the association of baseline (2011) glaucoma with subsequent (2018) social and economic variables. We applied sample weights to the glaucoma incidence between 2011 and 2018. These weights were constructed from weights for the structure of age and gender from the baseline wave of the CHARLS study. A $p < 0.05$ was defined as statistically significant.

RESULTS

Among 9973 eligible participants aged 50 years and above (table 1), 342 (3.4%) self-reported glaucoma diagnosis by a physician, while 9631 (96.6%) did not. Those who were diagnosed with glaucoma were significantly older (62.9 ± 7.99 vs 60.8 ± 7.49 , $p < 0.001$), were more likely to

Table 1 Baseline characteristics of 9973 participants aged 50 and above who participated in both the baseline and follow-up surveys

Baseline characteristic	Glaucoma	Non-glaucoma	P value
N	342	9631	
Mean age, years (SD)	62.9 (7.99)	60.81 (7.49)	<0.001
Self-rated distance Vision, n (%)			
Poor	132 (38.6)	2125 (22.06)	<0.001
Excellent, very good, good or fair	190 (55.6)	6956 (72.23)	
Hypertension present, n (%)	157 (45.9)	3978 (41.3)	0.232
Diabetes present, n (%)	49 (14.3)	731 (7.59)	<0.001
Obesity present, n (%)	83 (24.27)	2334 (24.23)	0.512
Male gender, n (%)	134 (39.2)	4714 (49.0)	<0.001
Drinking alcohol currently and also drinking in the past year	222 (64.9)	5565 (57.8)	0.029
Currently smoking and also smoking in the past	257 (75.2)	6488 (67.4)	0.010
Literacy, n (%)			
Illiterate	155 (45.3)	2932 (30.4)	<0.001
EQ-5D score, mean (SD)	0.72 (0.23)	0.79 (0.25)	<0.001
Depression score, mean (SD)	10.5 (7.08)	8.49 (6.32)	<0.001
Rural hukou, n (%)	279 (81.6)	7728 (80.2)	0.817
Mean log household per capita expenditure in USD, mean (SD)	4.59 (0.88)	4.62 (0.88)	0.539
Reporting having health insurance, n (%)	323 (94.4)	9057 (94.0)	0.880
Geographical location, n (%)			
Eastern region	92 (26.9)	3387 (35.2)	0.006
Central region	129 (37.7)	3138 (32.6)	
Western region	121 (35.4)	3106 (32.3)	
Married and living together	263 (76.9)	7870 (81.7)	0.024

A $p < 0.05$ was defined as statistically significant.
EQ-5D, EuroQol five-dimension questionnaire; SD, standard deviation.

Table 2 Glaucoma incidence of total population and total self-reported glaucoma between 2011 and 2018, stratified by age group

Age group (years)	Cases	Glaucoma incidence, among entire population (%)	Among those with self-reported incident glaucoma (%)
50–59	134	2.67	39.18
60–69	132	3.76	38.6
70–79	67	5.35	19.59
80+	9	4.79	2.63
Total	342	3.43	100

be female (60.8% vs 51.0%, $p<0.001$), to self-report poor distance vision at baseline (38.6% vs 22.1%, $p<0.001$), to have diabetes (14.3% vs 7.6%, $p<0.001$), to report currently drinking alcohol and drinking in the past year (64.9% vs 57.8%, $p<0.05$), currently smoking and smoking in the past (75.2% vs 67.4%, $p=0.01$), to be illiterate (45.3% vs 30.4%, $p<0.001$), have a lower health utility quality of life score (0.72 ± 0.23 vs 0.79 ± 0.25 , $p<0.001$), higher depression score (10.5 ± 7.08 vs 8.49 ± 6.32 , $p<0.001$) and to be unmarried (23.1% vs 18.3%) (table 1). The proportion of persons living in the different geographical regions of China also differed between those with versus without glaucoma (Eastern region 26.9% vs 35.2%, Central region 37.7% vs 32.6% and Western region 35.4% vs 32.3%, $p<0.01$).

Self-reported 7-year glaucoma incidence was 2.67% among the cohort aged 50–59 years, 3.76% at 60–69 years, 5.35% for 70–79 years and 4.79% for 80+ years (table 2).

Table 3 shows the glaucoma incidence between 2011 and 2018 stratified by geographic region and province. Central China had the highest incidence (3.95%) and Eastern China the lowest (2.64%).

In regression models of potential predictors of incidence of glaucoma among persons aged 50 years and above at baseline, older age (beta coefficient: 0.050, 95% CI: 0.001, 0.001, $p<0.001$), having hypertension (beta coefficient: 0.019, 95% CI: 0.006, 0.008, $p<0.001$) or diabetes (beta coefficient: 0.049, 95% CI: 0.028, 0.032, $p<0.001$), being female (beta coefficient: 0.036, 95% CI: 0.006, 0.020, $p<0.001$), recent smoking (beta coefficient: 0.029, 95% CI: 0.004, 0.020, $p=0.004$) and consumption of alcohol (beta coefficient: 0.026, 95% CI: 0.002, 0.017, $p<0.009$) and illiteracy (beta coefficient: -0.057 , 95% CI: -0.030 to -0.015 , $p<0.001$), were all significantly associated with increased incidence in the univariate model, and all factors except female sex remained significant in the multivariate model (table 4).

Incidence of glaucoma is 3.43% in the general study population, 3.94% among those with baseline hypertension, 6.52% among those with diabetes, 3.53% among those with obesity, 11.54% among those with low quality of life, 4.40% among those with depression and 3.35%

Table 3 Glaucoma incidence between 2011 and 2018, stratified by geographical region and province

Province	Self-reported glaucoma incidence, n (%)
Eastern China	
Tianjin	1 (1.30)
Beijing	1 (3.03)
Shanghai	1 (4.17)
Jiangsu	16 (3.16)
Zhejiang	11 (2.69)
Guangdong	8 (1.61%)
Liaoning	13 (4.38)
Fujian	7 (2.49)
Shandong	13 (1.48)
Jilin	13 (5.78)
Hebei	21 (4.45)
Heilongjiang	5 (3.29)
Total	92 (2.64)
Central China	
Hubei	7 (2.22)
Shanxi	11 (3.44)
Hunan	13 (2.80)
Henan	34 (4.32)
Jiangxi	15 (2.92)
Anhui	31 (6.34)
Total	129 (3.95)
Western China	
Inner Mongolia	19 (4.39)
Chongqing	15 (2.92)
Shaanxi	11 (3.56)
Xinjiang	3 (7.14)
Qinghai	4 (4.88)
Sichuan	40 (4.38)
Guangxi	8 (2.22)
Gansu	8 (3.56)
Yunnan	15 (2.55)
Guizhou	3 (2.44)
TOTAL	121 (3.75)

among those with below average household per capita expenditure.

Logistic regression models showed significant association between incidence of the following characteristics and baseline glaucoma: poor self-reported distance vision (beta coefficient: 1.106, 95% CI: 0.701, 1.511, $p<0.001$), having hypertension (beta coefficient: 0.545, 95% CI: 0.496, 0.593, $p<0.001$), having diabetes (beta coefficient: 0.388, 95% CI: 0.326, 0.449, $p<0.001$), not having obesity (beta coefficient: -0.184 , 95% CI: -0.239 , -0.129 , $p<0.001$) and lower mean value of health utility score of

Table 4 Regression models of potential predictors of self-reported incident glaucoma diagnosed by any physician between 2011 and 2018 among a nationally representative sample of Chinese persons aged ≥ 50 years who did not have glaucoma at baseline (n=9973)

Baseline characteristic	Univariate model		Multivariate model	
	β (95% CI)	P value	β (95% CI)	P value
Age, years	0.050 (0.001, 0.002)	<0.001	0.035 (0.001, 0.001)	<0.001
Hypertension present (adjusted for age)	0.019 (0.006, 0.008)	<0.001	0.005 (0.001, 0.003)	0.001
Diabetes present (adjusted for age)	0.049 (0.028, 0.032)	<0.001	0.052 (0.030, 0.034)	<0.001
Female	0.036 (0.006, 0.020)	<0.001	-0.001 (-0.002, 0.001)	0.611
Currently smoking and also smoking in the past	0.029 (0.004, 0.020)	0.004	0.020 (0.007, 0.010)	<0.001
Drinking alcohol currently and also drinking in the past year	0.026 (0.002, 0.017)	0.009	0.013 (0.004, 0.006)	<0.001
Literate	-0.057 (-0.030, 0.015)	<0.001	-0.044 (-0.019, 0.016)	<0.001
Rural (vs non-rural) hukou	0.006 (-0.006, 0.012)	0.544		

Bold type indicates values significant at $p < 0.05$.

residents' quality of life (beta coefficient: -0.040, 95% CI: -0.006, 0.776, $p < 0.001$) (table 5).

DISCUSSION

Glaucoma poses a substantial public health challenge globally, especially in ageing populations. A previous study reported the prevalence of glaucoma in the same population; however, glaucoma incidence has not

been reported in this population.¹⁰ Our study aimed to comprehensively assess nationwide glaucoma incidence rates and associated determinants among older Chinese persons, addressing a critical gap in understanding this condition's epidemiology.

Using data from the CHARLS from 2011 to 2018, we identified significant factors associated with incident glaucoma among adults aged 50 and older. Our findings demonstrated a self-reported glaucoma incidence of 3.4%, revealing demographic, socioeconomic and health-related disparities among those diagnosed with glaucoma versus those without. Affected individuals were older, more likely to report poor vision and exhibit comorbidities like diabetes and experienced lower quality of life scores and higher depression levels.

Regional variations were evident in glaucoma incidence, with Central China reporting the highest incidence rates. Glaucoma development can be attributed to a complex interplay of genetic, environmental and socioeconomic factors.¹¹ In Central China, for instance, neovascularisation and trauma were identified as leading causes for secondary glaucoma.¹² This elevated rate of trauma-related glaucoma may be linked to specific regional activities or occupational hazards prevalent in Central China.¹² Our findings showed that older age, hypertension, diabetes, illiteracy, smoking and alcohol consumption were associated with increased glaucoma incidence, highlighting the multifactorial nature of its risk factors. Our longitudinal findings are consistent with the cross-sectional baseline paper¹⁰ showing associations between glaucoma and hypertension, diabetes, smoking and alcohol use. Interestingly, our study found no association between incident glaucoma and socioeconomic or urban-rural status, despite a significant link between illiteracy and incident glaucoma. This contrasts with previous prevalence studies that reported an association between lower socioeconomic status and higher glaucoma prevalence.¹³ These findings highlight the need for more longitudinal studies on glaucoma

Table 5 Logistic regression models of potential subsequent clinical, mental health and socioeconomic associations with baseline (2011) self-reported glaucoma diagnosed by any physician, among a nationally representative sample of Chinese persons aged ≥ 50 years (n=9973)

2018 status	β (95% CI)	P value
Poor self-reported distance vision	1.106 (0.701, 1.511)	<0.001
Hypertension present (adjusted for age)	0.545 (0.496, 0.593)	<0.001
Diabetes present (adjusted for age)	0.388 (0.326, 0.449)	<0.001
Obesity 2015 (adjusted for 2011 age)*	-0.184 (-0.239, 0.129)	<0.001
Mean value of health utility score of residents' quality of life (EQ-5D)	-0.040 (-0.134, 0.045)	<0.001
Mean depression score (adjusted for age)	0.018 (-0.006, 0.776)	0.051
Logarithm of household per capita expenditure	-0.012 (-0.336, 0.072)	0.204

Bold type indicates values significant at $p < 0.05$ (Independent variable=2011 glaucoma status, dependent variable=subsequent 2018 clinical, mental health and socioeconomic associations.

*Wave 4 (2018) did not undergo physical examinations or blood tests; therefore, the BMI is based on the 2015 data.

BMI, body mass index; CI, confidence interval; EQ-5D, EuroQol five-dimension questionnaire.

incidence and risk factors, beyond cross-sectional prevalence studies.

Importantly, our study also underscored the impact of incident glaucoma on individuals' subsequent health and socioeconomic status. Those diagnosed with glaucoma at baseline showed increased incident comorbidities, lower quality of life scores and financial vulnerabilities, reflecting the broader repercussions of this condition beyond vision impairment. These findings not only highlight the importance of early prevention of risk factors, but also emphasise the full-cycle management of glaucoma in alleviating the associated disease burden in terms of health system and society.

Our findings align with previous localised studies in China, highlighting the significance of older age and lower education level as key risk factors for glaucoma.^{4 5} However, our nationwide scope underscores the need for targeted interventions at the national level addressing these risk factors.

The strengths of this study include use of a large, nationally representative sample, drawn from most of China's administrative units and selected using standardised protocols. Study personnel were trained in standard fashion to ensure all procedures were carried out as described. Our findings provide a basis for comparison with other low-income and middle-income countries and especially those undergoing rapid economic transitions. Limitations include the self-reported nature of glaucoma diagnosis and potential under-reporting or misdiagnosis, though evidence has suggested that self-reported medical diagnosis is reasonably accurate. A study showed that of the 200 subjects with medical record information indicating glaucoma, 165 (77.0%) correctly self-reported their glaucoma diagnosis.¹⁴ Among the 130 subjects with medical record information indicating glaucoma suspect or ocular hypertension, 109 (83.9%) correctly self-reported no glaucoma diagnosis.¹⁴ However, further studies validating self-reported diagnoses against medical records are warranted. An additional limitation is that the results may have been influenced by patients (42.2%) who did not attend the follow-up examination, or who failed to provide information on their glaucoma history. This might induce attrition bias in the results.

CONCLUSIONS

Our study provides critical insights into glaucoma incidence rates, risk factors and their association with long-term disease and psychosocial outcomes on the Chinese population. These findings underscore the need for national-level public health strategies focusing on early detection, improved vision care and targeted interventions to mitigate the burden of glaucoma and enhance overall well-being. This research lays the foundation for further investigations and policy formulations aimed at addressing the multifaceted challenges posed by glaucoma in China, ultimately contributing to enhanced eye health and improved quality of life.

Author affiliations

- ¹Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne, Victoria, Australia
- ²Ophthalmology, Department of Surgery, The University of Melbourne, Melbourne, Victoria, Australia
- ³Institute of Population Research, Peking University, Beijing, China
- ⁴Peking University Library, Peking University, Beijing, China
- ⁵Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing Key Laboratory of Ophthalmology and Visual Sciences, Beijing, People's Republic of China
- ⁶School of Optometry, The Hong Kong Polytechnic University, Kowloon, Hong Kong, People's Republic of China
- ⁷Centre for Public Health, Queen's University Belfast, Belfast, UK
- ⁸State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-Sen University, Guangzhou, China
- ⁹ORBIS International, New York, New York, USA

Acknowledgements The authors would like to thank the CHARLS team for the data that enabled these analyses. This research is supported by the Wellcome funding (R2806CPH).

Contributors CJ conceptualised the study, did literature review, participated in data collection and data analysis, drafted the manuscript and is responsible for the overall content as a guarantor. NC conceptualised the study, interpreted data and revised the manuscript. XJ participated in data collection and data analysis, interpreted data, created the figure and revised the manuscript. MK, JL, WH, RC, LL and MH participated in data interpretation and manuscript revision. All authors read and approved the final draft.

Funding This research is supported by the Wellcome funding (R2806CPH). CJ is supported by the Research Training Program Scholarship from the Commonwealth Government of Australia.

Disclaimer The funding bodies had no role in any aspect of the study or the decision to submit this manuscript for publication.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. The CHARLS was approved by the Ethical Review Committee of Peking University, and all participants gave written informed consent at the time of participation. This study used public data and, therefore, does not require additional ethics approval. The original CHARLS from which the data were collected was carried out in Beijing and has obtained ethics approval from the Ethical Review Committee of Beijing University; the ethical approval number is IRB00001052-11015. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Catherine Jan <http://orcid.org/0000-0001-7383-8208>
Xin Jin <http://orcid.org/0000-0001-8712-4513>
Nathan Congdon <http://orcid.org/0000-0001-9866-3416>

REFERENCES

- 1 Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol* 2012;96:614–8.
- 2 Song P, Wang J, Bucan K, et al. National and subnational prevalence and burden of glaucoma in China: A systematic analysis. *J Glob Health* 2017;7:020705.
- 3 Sun Y, Chen A, Zou M, et al. Disease Burden of Glaucoma in China: Findings from the Global Burden of Disease 2019 Study. *Clin Epidemiol* 2022;14:827–34.

- 4 Pan C-W, Yang W-Y, Hu D-N, *et al.* Longitudinal Cohort Study on the Incidence of Primary Open-Angle Glaucoma in Bai Chinese. *Am J Ophthalmol* 2017;176:127–33.
- 5 Zhang Y, Hao J, Zhang Q, *et al.* Five-year incidence of primary glaucoma and related risk factors - The Handan eye study. *Acta Ophthalmol* 2023;101:e50–60.
- 6 Chiu S-L, Chu C-L, Muo C-H, *et al.* The Prevalence and the Incidence of Diagnosed Open-Angle Glaucoma and Diagnosed Angle-Closure Glaucoma: Changes From 2001 to 2010. *J Glaucoma* 2016;25:e514–9.
- 7 Jan C, Xin J, Dong Y, *et al.* Patterns and determinants of incident cataract surgery in China from 2011 to 2015 using a nationally representative longitudinal database. *BMJ Open* 2023;13:e069702.
- 8 Zhao Y, Hu Y, Smith JP, *et al.* Cohort profile: the China Health and Retirement Longitudinal Study (CHARLS). *Int J Epidemiol* 2014;43:61–8.
- 9 Lewinsohn PM, Seeley JR, Roberts RE, *et al.* Center for Epidemiologic Studies Depression Scale (CES-D) as a screening instrument for depression among community-residing older adults. *Psychol Aging* 1997;12:277–87.
- 10 Sun J, Li T, Zhao X, *et al.* Prevalence and Risk Factors of Glaucoma Among Chinese People From the China Health and Retirement Longitudinal Study. *J Glaucoma* 2022;31:789–95.
- 11 Jayaram H, Kolko M, Friedman DS, *et al.* Glaucoma: now and beyond. *The Lancet* 2023;402:1788–801.
- 12 Liu Q, Liu C, Cheng W, *et al.* Clinical analysis of secondary glaucoma in Central China. *Sci Rep* 2023;13:8439.
- 13 Shweikh Y, Ko F, Chan MPY, *et al.* Measures of socioeconomic status and self-reported glaucoma in the U.K. Biobank cohort. *Eye (Lond)* 2015;29:1360–7.
- 14 MacLennan PA, McGwin G Jr, Searcey K, *et al.* Medical Record Validation of Self-Reported Eye Diseases and Eye Care Utilization Among Older Adults. *Curr Eye Res* 2013;38:1–8.