

BMJ Open Factors associated with declining under-five mortality rates from 2000 to 2013: an ecological analysis of 46 African countries

Aaron M Kipp,^{1,2} Meridith Blevins,^{1,3} Connie A Haley,^{1,2} Kasonde Mwinga,⁴ Phanuel Habimana,⁴ Bryan E Shepherd,^{1,3} Muktar H Aliyu,^{1,5} Tigist Ketsela,⁴ Sten H Vermund^{1,6}

To cite: Kipp AM, Blevins M, Haley CA, *et al.* Factors associated with declining under-five mortality rates from 2000 to 2013: an ecological analysis of 46 African countries. *BMJ Open* 2016;**6**:e007675. doi:10.1136/bmjopen-2015-007675

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-007675>).

Received 3 February 2015
Revised 17 August 2015
Accepted 21 September 2015



CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Aaron M Kipp;
aaron.kipp@vanderbilt.edu

ABSTRACT

Objective: Inadequate overall progress has been made towards the 4th Millennium Development Goal of reducing under-five mortality rates by two-thirds between 1990 and 2015. Progress has been variable across African countries. We examined health, economic and social factors potentially associated with reductions in under-five mortality (U5M) from 2000 to 2013.

Setting: Ecological analysis using publicly available data from the 46 nations within the WHO African Region.

Outcome measures: We assessed the annual rate of change (ARC) of 70 different factors and their association with the annual rate of reduction (ARR) of U5M rates using robust linear regression models.

Results: Most factors improved over the study period for most countries, with the largest increases seen for economic or technological development and external financing factors. The median (IQR) U5M ARR was 3.6% (2.8 to 5.1%). Only 4 of 70 factors demonstrated a strong and significant association with U5M ARRs, adjusting for potential confounders. Higher ARRs were associated with more rapidly increasing coverage of seeking treatment for acute respiratory infection ($\beta=0.22$ (ie, a 1% increase in the ARC was associated with a 0.22% increase in ARR); 90% CI 0.09 to 0.35; $p=0.01$), increasing health expenditure relative to gross domestic product ($\beta=0.26$; 95% CI 0.11 to 0.41; $p=0.02$), increasing fertility rate ($\beta=0.54$; 95% CI 0.07 to 1.02; $p=0.07$) and decreasing maternal mortality ratio ($\beta=-0.47$; 95% CI -0.69 to -0.24 ; $p<0.01$). The majority of factors showed no association or raised validity concerns due to missing data from a large number of countries.

Conclusions: Improvements in sociodemographic, maternal health and governance and financing factors were more likely associated with U5M ARR. These underscore the essential role of contextual factors facilitating child health interventions and services. Surveillance of these factors could help monitor which countries need additional support in reducing U5M.

Strengths and limitations of this study

- This study represents a comprehensive analysis of health factors and other determinants of health associated with declining under-five mortality in Africa.
- Study findings highlight the role of financing, economy and the broader health system in more rapidly reducing an already declining, albeit slowly, under-five mortality rate above and beyond the contribution of individual interventions.
- Inferences of causality may be limited due to the ecological study design and that the time periods for change in coverage of the factors did not always fully overlap with time periods for change in under-five mortality.
- The small sample size (at most $n=46$ countries, but often smaller due to missing data) limits the validity of some associations and prevents adjustment for too covariates.

INTRODUCTION

Under-five mortality (U5M) rates have declined worldwide from an estimated 90 deaths per 1000 live births in 1990 to 46 deaths per 1000 in 2013, representing an average annual rate of reduction (ARR) of 3%.¹ Higher income regions had an estimated U5M rate of 6 deaths per 1000 live births in 2012 (3.8% ARR); they account for only 1.4% of all under-five deaths. In contrast, the WHO African Region (including 46 countries in 2012; [figure 1](#) and [table 1](#)) experienced 90 under-five deaths per 1000 live births in 2013 and a 1990–2013 ARR of 2.9%.¹ Thus, African children are nearly 16 times more likely to die under age five than children from high income nations, though progress in Africa is being made.

In 2000, the 4th Millennium Development Goal (MDG4)² proposed the reduction of the U5M rate by two-thirds between 1990

and 2015, requiring a global average ARR of 4.4% to reach this goal. Africa has more than tripled its U5M ARR from 1.3% in 1990–2000 to 4.2% in 2000–2013, but the ARR achievements are inadequate to meet the MDG4 targets.¹ Proven cost-effective child survival interventions and proposed delivery channels for those interventions need to be scaled up.^{3–7} The *Countdown to 2015 for Maternal, Newborn and Child Survival* initiative was created in 2005 to monitor the use of these evidence-based interventions for maternal, neonatal and child health.⁸ In 2012, the Countdown's 75 priority countries accounted for >95% of all maternal and child deaths.^{9 10} Of note, 41 (55%) of the 75 Countdown priority countries come from the 46 WHO African Region countries (the exceptions were Algeria, Cape Verde, Mauritius, Namibia and Seychelles).

In 2006, WHO, the United Nations Children's Fund, and the World Bank developed Child Survival Strategy for the African Region aimed to support countries' efforts to reduce child deaths from preventable and treatable conditions by scaling up coverage of effective interventions.⁶ These include antenatal care, postnatal care, infant and young child feeding, the expanded programme on immunisations, integrated management of common childhood illness, prevention of mother-to-child transmission of HIV and use of insecticide treated nets, among others. As of 2013, 38 of the 46 African countries had developed or updated national child survival policies, strategies and plans. Population coverage in the African Region has increased with time, leading to high coverage of measles vaccination, neonatal tetanus protection, antiretroviral drugs for prevention of mother-to-child transmission of HIV and at least one antenatal care visit for every pregnant woman.¹¹ Coverage of other interventions remain low, however, in spite of recent increases in coverage, including exclusive breastfeeding, pneumococcus and *Haemophilus influenzae* vaccination, treatment of childhood diarrhoea, pneumonia and malaria, and use of insecticide treated nets. Despite progress, Africa is not on track to reach MDG4 due to inconsistent and suboptimal implementation and scale-up of evidence-based maternal, neonatal and child health interventions.¹⁰ With variable progress across countries in improving child survival in Africa, understanding factors associated with the most rapid changes in U5M reductions could assist regional and country efforts to improve child survival in the post-2015 era.

We sought to identify proximal (eg, interventions and services) and distal (eg, socioeconomic) factors influencing how rapidly countries reduce U5M, as evidenced by their ARR. An ecological analysis of country-level data was used to determine which factors previously reported to reduce U5M are also the factors most strongly associated with higher ARRs in Africa.

METHODS

The study was approved by the Institutional Review Boards at Vanderbilt University.

Mortality data were obtained for the period 2000–2013, while data on all factors of interest were obtained for the period 1998–2011. These are described in detail below.

Under-five mortality data

We obtained country-specific U5M rates from annual estimates provided by <http://www.childmortality.org>, the data used in the United Nations Children's Fund (UNICEF) Report on *Levels and Trends in Child Mortality*. For each of the 46 WHO African Region countries, the estimated U5M rates for 2000 and 2013 (the latest estimates available at the time of the analysis (accessed June 2015)) were obtained and the ARR from 2000 to 2013 was calculated. Use of the ARR as an outcome facilitates interpretation of results in the context of MDG4 progress metrics.

The U5M ARR reflects a constant rate of change in the U5M rate between two time periods and is calculated using the following equation (equation 1):

$$ARR = \left(\frac{\ln[y_{t+n}/y_t]}{n} \right) \times (-100)$$

where y_t is the mortality rate for a given year (eg, 2000) and n is the number of years between the two rates (eg, 13 years when calculating ARR from 2000 and 2013 rates). Consistent with how it is calculated and reported by UNICEF¹ and others,¹² the ARR is expressed as a per cent and will have a positive value when a country is reducing its mortality rate. As an example, an ARR of 4.4% or greater is needed for a country to meet MDG4 of reducing U5M by two-thirds between 1990 and 2015. In the years leading up to 2015, expressing the reduction in U5M as an ARR made it possible to monitor progress across countries and over different time periods.

Factors potentially associated with child mortality

Identification

Since we wanted to assess a broad range of factors potentially associated with U5M ARR, factors to be used in the analysis included those monitored by Countdown to 2015 as well as others identified through a comprehensive literature search of the PubMed database. We searched for studies on under-five, infant or neonatal mortality within any of the 46 countries in the WHO African Region that were published between 2002 and 2012. Abstracts were reviewed to identify factors that were (1) associated with under-five, infant or neonatal mortality, (2) not already reported by Countdown to 2015 and (3) had aggregate country-level data available for the analysis. Thirty-four factors met these criteria. These were combined with 20 Countdown 2015 intervention indicators (or closely related) and 16 non-intervention indicators reported in the Countdown 2015 country profiles (or were closely related). The final list used in the analysis included 70 factors from the

following categories: sociodemographics (18 factors), access to healthcare (16), governance and financing (11), maternal health (6), child survival interventions (7), clinical and health conditions (7), and other country infrastructure (5) (see online supplemental table S1).

Data sources and inclusion criteria

Data for 58 (83%) of the 70 indicators were obtained from the World Bank Data Catalogue.¹³ The World Bank Data Catalogue is a repository of national, regional and global indicator data that have been compiled from officially recognised international sources. In many instances, a single indicator may be derived from multiple data sources using modelling or aggregation techniques. Data for the remaining 12 factors not available through the World Bank Data Catalogue were obtained directly from each country's Demographic and Health Surveys¹⁴ (10 factors) or Countdown 2015 country profiles (2 factors).^{8 10} In total, 26 (37%) of 70 factors were obtained entirely or in part from country Demographic and Health Surveys or other household survey data, including the majority of the maternal health, access to healthcare and child survival intervention factors. Other sources included data collected and maintained by WHO, various UN divisions, UNAIDS, the Organisation for Economic Co-operation and Development, and the World Bank (see online supplemental tables S2 and S3 for further details).

For each of the 46 WHO African Region countries, data on the 70 factors were obtained that corresponded as close as possible to the years 2000 and 2011. To be considered sufficient for inclusion in the analyses, data for each factor had to meet the following three criteria: (1) reported for one of the years between 1998 and 2003, termed *2000 data*, (2) reported for one of the years between 2006 and 2011, termed *2011 data* and (3) the pair of data points for each factor had to be at least 5 years apart in order to minimise incorrect extrapolation when calculating changes in the indicator. If any of these criteria were not met, the change over time for that particular factor was not calculated and was deemed missing.

Indicator annual rate of change calculation

The annual rate of change (ARC) for each indicator is conceptually similar to the ARR for U5M and was calculated using the same ARR equation shown above in equation 1 but with one difference: the rate of change is multiplied by *positive* 100 instead of *negative* 100 so that the ARC has a negative value when the indicator decreases over time (ie, $\text{Coverage}_{2011} < \text{Coverage}_{2000}$) and a positive value when the indicator increases over time (equation 2):

$$\text{ARC} = \left(\frac{\ln[y_{t+n}/y_t]}{n} \right) \times (100)$$

This is in contrast to the U5M ARR which has a positive value when mortality is decreasing over time (ie, $\text{U5M}_{2013} < \text{U5M}_{2000}$).

Statistical analysis

The dependent variable of interest was the U5M ARR for 2000–2013. Each indicator ARC was an independent variable of interest. Both indicator ARCs and U5M ARR were analysed as continuous variables and no transformations were performed. The distribution of ARRs was inspected visually and was confirmed to be approximately normal by the Shapiro-Wilk test for normality ($p=0.33$). Associations were explored using linear regression. Given the sample size ($n=46$ countries), results from traditional linear regression methods may be overly influenced by outliers. These outliers, however, likely represent true data rather than data errors and exclusion would unnecessarily decrease the sample size. Hence, robust linear regression was used to minimise the influence of outlying observations, without excluding them^{15–17} using iteratively reweighted least squares (M-estimation with Huber weighting).

Multivariable robust linear regression models were fit for each factor of interest, resulting in 70 different regression models (one for each factor). A standard set of factors was identified a priori to be included in each model as covariates to adjust for potential confounding of the specific factor association being analysed. Given the sample size of 46 countries, at most, a decision was also made a priori to include no more than five covariates in the multivariable analyses to avoid overfitting. We selected the following factor ARCs for inclusion as covariates in each model based on previous ecological studies, consideration of what macro-level or system-level factors would influence the majority of the factors, and having nearly complete data: (1) access to improved water source, (2) health expenditure (relative to gross domestic product (GDP)), (3) adult HIV prevalence, (4) urban population prevalence and (5) receipt of antenatal care (when applicable). All regression models included these core factor ARCs as covariates unless expected to be highly correlated with the primary indicator of interest (eg, improved water sources was excluded from the model for the association between improved sanitation facilities; health expenditure relative to GDP was excluded from the model for the association between health expenditure relative to government expenditure). Changes in the receipt of antenatal care was only included when the indicator being analysed would occur following the pregnancy period (eg, maternal mortality ratio, births delivered at a health facility, measles immunisation, wasting prevalence) and was not included for sociodemographic factors, macro-level factors such as health expenditure and system-level factors such as physician density.

The estimated robust linear regression β coefficient for each indicator ARC and U5M ARR association can be interpreted as the change in ARR associated with

every 1% increase in the indicator ARC. For example, a β coefficient of 0.20 indicates that for every 1% increase in the indicator ARC there is a corresponding 0.2% increase in the ARR. Stated differently, a 5% increase in the indicator ARC (eg, 6% ARC compared to 1% ARC) corresponds to a 1% increase in the ARR (eg, 4% ARR vs 3% ARR). To help avoid type II errors, which can occur with small samples, we reported all associations when the p values were <0.10 , acknowledging that some of these associations may be due to chance alone, especially with higher p values. We report an indicator to be strongly associated with ARR when the adjusted β coefficient is ≥ 0.20 or ≤ -0.20 .

A preliminary analysis showed significantly different ARRs between countries reporting a specific indicator and countries not reporting a specific indicator when the indicator was reported by $<50\%$ of the countries. Since these differences suggest selection bias, we only present results for those factors for which at least 23 countries ($\geq 50\%$) have sufficient data (ie, non-missing indicator data for both time periods). Results for all factors are available in online supplemental tables S4 and S5.

All analyses were conducted using R-software V.2.15.2 (<http://www.r-project.org>).

RESULTS

Country-specific progress in U5M

As of 2013, U5M rates and corresponding ARRs for each of the 46 WHO African countries vary widely, with a median (IQR) of 3.6% (2.8% to 5.1%; [figure 1](#); [table 1](#)). Eleven countries (Eritrea, Ethiopia, Liberia, Madagascar, Malawi, Mozambique, Niger, Rwanda, Senegal, Tanzania and Uganda) have met or are on track to meet their MDG4 target. An additional four countries (Algeria, Cape Verde, Mauritius and Seychelles) are considered on track due to U5M rates <40 per 1000. Eight countries were making very little progress with ARRs $\leq 2\%$: Angola, Central African Republic, Chad, Comoros, Ghana, Lesotho, Mauritania and Zimbabwe. Of the remaining 23 countries, some were making good progress but will fall just short of their MDG4 target (eg, Congo, Mali and Zambia), while others were making some progress but still have a considerable ways to go before reaching MDG4 (eg, Cameroon, Kenya and South Africa; [table 1](#)).

Descriptive results of selected factors

Of the 70 factors identified, 60 (86%) had sufficient data from at least one country for the period 2000–2011. Of these, 41 (68%) factors had data for at least 50% of the countries and are shown in [table 2](#). The largest ARCs were generally observed for factors related to economic or technological development and external financing, ranging from an ARC of 4.2% (GDP per capita) to 29% (internet users) and including, for example, external resources for health (ARC of 6.2%). An extremely

large ARC also occurred for use of insecticide-treated bednets (ARC of 29.0%), an intervention which can be rapidly scaled up and which also had the lowest coverage at the start of the study period (2% in 2000). In contrast, vitamin A supplementation had very high coverage at the start of the study period (91%) and did not change dramatically over time (ARC of -0.3%). Coverage of other child survival interventions or access to healthcare generally improved over the study period with ARCs from 1.2% to 4.0%. Some countries, however, had worsening coverage over time as indicated by negative ARCs in the IQR. Adverse health conditions improved slowly over the study period with ARCs from -0.3% to -1.9% , and the largest improvement was seen for maternal mortality ratio (ARC of -3.7%). Similar to the adverse health conditions, sociodemographic factors improved over the study period for nearly all countries, with ARCs ranging from 0.1% to 3.2%. Details of all indicator ARCs are in online supplemental table S4.

Associations between indicator ARCs and U5M ARRs

There was evidence that ARRs were higher for countries that had more complete data reported for use in the analyses ([figure 2](#)). However, of the results presented below, all but one factor was reported by >30 countries.

Four indicator ARCs were strongly and statistically associated with U5M ARR after adjusting for potential confounders ([table 3](#), in bold). With three of these, larger ARCs were associated with larger U5M ARRs: fertility rate ($\beta=0.54$; $p=0.07$), health expenditure relative to GDP ($\beta=0.26$; $p=0.01$) and seeking treatment for acute respiratory infection (ARI) ($\beta=0.22$; $p=0.01$). With the fourth, larger ARCs for maternal mortality ratio were inversely associated with unfavourable and larger decreases in U5M ARR ($\beta=-0.47$; $p<0.01$).

An additional four factors had large ($\beta \geq 0.20$ or $\beta \leq -0.20$) but statistically non-significant ($p \geq 0.10$) associations with U5M ARR, while another four showed small but statistically significant associations ($\beta < 0.20$ with $p < 0.10$; [table 3](#)). Of the remaining 58 factors considered in this analysis, 27 had sufficient data from at least 23 of the 46 countries, but there was no association with U5M ARR ($\beta < 0.20$ and $p \geq 0.10$; seven are shown in [table 3](#)). The final 31 factors had no data available for analysis ($n=10$) or were reported by <23 countries and not presented here due to concerns about selection bias. Results for all 70 factors are available in online supplemental table S5.

DISCUSSION

Our study represents a comprehensive analysis of health-related factors and other determinants of health associated with declining U5M in Africa. From 70 factors considered, only four were found to have a strong and statistically significant association with higher rates of reduction in U5M between 2000 and 2013: more rapid increases in fertility rate, health expenditure relative to




Figure 1 Annual rate of reduction (ARR) in under-five mortality for the 46 countries in the WHO African Region, 2000 to 2013 (Higher ARR indicates more progress, represented by a greener colour on the map). The WHO African Region excludes Djibouti, Egypt, Libya, Morocco, Somalia, Sudan and Tunisia; all are included in the WHO Eastern Mediterranean Region. From May 2012, South Sudan is now officially part of the African Region but was not an independent WHO member nation during the study period.

GDP and seeking treatment for ARI, and a more rapid decrease in maternal mortality ratio. Maternal mortality ratio showed a particularly strong association ($\beta = -0.47$), indicating that a 1% increase in the annual reduction of the maternal mortality ratio was associated with a 0.47% increase in the U5M ARR. The associations for health expenditure relative to GDP and seeking treatment for ARI were half this magnitude ($\beta = 0.26$ to $\beta = 0.22$, respectively).

The strong and statistically significant association between more rapid declines in the maternal mortality ratio and U5M ARR may well represent the direct negative effect of maternal death on subsequent child survival. It may also be indicative of health systems that do poorly in preventing both maternal and child deaths. More rapid increases in treatment-seeking behaviour for ARI was the only health-related access or intervention

factor found to associate with a larger U5M ARR. Validation studies suggest that factors of access to care or behaviour (eg, seeking ARI treatment) may be more reliably measured than actual treatment (eg, children with ARI receiving treatment; we had insufficient data to analyse this), since quality of care and ability to recall symptoms may bias the coverage estimates.^{18 19} Similar concerns exist about reliability of vaccination coverage estimates due to information bias on the vaccination card or recall bias from maternal reports.²⁰ This may explain, in part, why some of these factors were not associated with ARR in our study.

The association between fertility rate and higher U5M ARR is at first counterintuitive. However, as shown in table 2, fertility rates are declining (negative ARC) in nearly all 46 of the countries. Thus, slower declines in fertility rates are associated with more rapid reductions

Table 1 Under-five mortality rates and annual rate of reduction (ARR) for the 46 countries in the WHO African Region, 2000–2013

Country	Under-five mortality rate (per 1000) in the given year*			ARR (%) 2000 to 2013	ARR (%) in 2000–2015 needed to meet MDG
	2000	2013	MDG for 2015		
Algeria†	39.6	25.2	15.7	3.5	6.2
Angola	216.7	167.4	75.3	2.0	7.0
Benin	146	85.3	59.8	4.1	6.0
Botswana	85.1	46.6	16.5	4.6	10.9
Burkina Faso	185.8	97.6	67.4	5.0	6.8
Burundi	148.9	82.9	56.9	4.5	6.4
Cameroon	151.2	94.5	45.5	3.6	8.0
Cape Verde†	35.3	26	21.0	2.4	3.5
Central African Republic	174.1	139.2	59.0	1.7	7.2
Chad	190.7	147.5	71.6	2.0	6.5
Comoros	101.3	77.9	41.8	2.0	5.9
Congo	121.4	49.1	30.7	7.0	9.2
Côte d'Ivoire	146.1	100	50.5	2.9	7.1
Dem. Rep. of the Congo	175.9	118.5	58.7	3.0	7.3
Equatorial Guinea	142.4	95.8	61.3	3.0	5.6
Eritrea	89.3	49.9	50.2	4.5	3.8
Ethiopia	145.5	64.4	68.3	6.3	5.0
Gabon	84.6	56.1	30.9	3.2	6.7
Gambia	119	73.8	56.6	3.7	5.0
Ghana	101.3	78.4	42.7	2.0	5.8
Guinea	170.2	100.7	79.2	4.0	5.1
Guinea-Bissau	180.8	123.9	74.9	2.9	5.9
Kenya	110.9	70.7	32.9	3.5	8.1
Lesotho	114.6	98	28.8	1.2	9.2
Liberia	175.2	71.1	82.7	6.9	5.0
Madagascar	110.6	56	53.6	5.2	4.8
Malawi	174.2	67.9	81.8	7.2	5.0
Mali	219.9	122.7	84.7	4.5	6.4
Mauritania	113.1	90.1	39.3	1.7	7.0
Mauritius†	18.6	14.3	7.7	2.0	5.9
Mozambique	168.5	87.2	79.0	5.1	5.0
Namibia	75.5	49.8	24.5	3.2	7.5
Niger	226.9	104.2	109.1	6.0	4.9
Nigeria	187.7	117.4	71.1	3.6	6.5
Rwanda	181.9	52	50.6	9.6	8.5
Sao Tome and Principe	89.3	51	36.8	4.3	5.9
Senegal	137	55.3	47.0	7.0	7.1
Seychelles†	14.2	14.2	5.5	0.0	6.3
Sierra Leone	231.5	160.6	89.2	2.8	6.4
South Africa	74.3	43.9	20.3	4.0	8.6
Swaziland	122.5	80	24.6	3.3	10.7
Togo	121.8	84.7	48.8	2.8	6.1
Uganda	147	66.1	59.6	6.1	6.0
United Rep. of Tanzania	131.5	51.8	55.7	7.2	5.7
Zambia	168.8	87.4	64.2	5.1	6.4
Zimbabwe	102.6	88.5	24.9	1.1	9.4

*U5M rate data obtained from <http://www.childmortality.org>, accessed June 2015.

†Countries with U5M rates <40 per 1000 in 2013 are considered to be on track despite having 1990–2013 ARR of less than 4%. ARR, annual rate of reduction; MDG, Millennium Development Goal.

in U5M—something that becomes more intuitive in the context of demographic transition, Birth rates begin to decline typically only after mortality rates have begun to decline. The observed association may therefore reflect that much of Africa is in the early and middle stages of

the demographic transition. Countries with the most rapid reductions in U5M have just begun to reduce their fertility rates, while countries that previously had substantial reductions in mortality (and therefore smaller ARRs in the recent past) have the most rapid reductions

Table 2 Median 2000 indicator values and corresponding indicator annual rate of change (ARC) for the 2000–2011 period

Indicator	N	Median 2000 value (IQR)	Median 2000–2011 ARC (IQR)
Sociodemographics			
Adult female literacy rate (%)	35	54 (37 to 72)	1.2 (0.8 to 2.7)
Female labour participation rate (%)	45	46 (43 to 49)	0.1 (–0.1 to 0.3)
Fertility rate (births per woman)	46	6 (5 to 6)	–1.4 (–2.1 to –0.9)
Gross domestic product (GDP) per capita (PPP int'l \$)	45	990 (680 to 1900)	4.2 (2.8 to 5.9)
Human development index (0–1)	39	0.38 (0.31 to 0.48)	1.1 (0.7 to 1.9)
Improved sanitation facilities (%)	45	25 (13 to 45)	1.6 (0.4 to 2.6)
Improved water source (%)	45	62 (51 to 79)	1.1 (0.2 to 1.8)
Labour force participation rate among women (%)	45	66 (50 to 76)	0.3 (–0.1 to 0.6)
Urban population prevalence (%)	46	35 (22 to 44)	1.3 (0.7 to 1.6)
Women with at least some secondary education (%)	23	16 (10 to 30)	3.2 (2.5 to 5.2)
Maternal health			
Pregnant women receiving prenatal care (%)	38	77 (64 to 88)	1.3 (0.6 to 2.7)
Adult female mortality rate (rate per 1000 persons)	46	360 (280 to 420)	–1.2 (–1.8 to –0.6)
Maternal mortality ratio (rate per 100 000 births)	45	550 (400 to 840)	–3.7 (–4.6 to –2.0)
Access to health care			
Seeking ARI treatment (%)	33	40 (27 to 55)	1.7 (–0.3 to 4.9)
Births attended by skilled health staff (%)	37	47 (39 to 60)	2.0 (0.3 to 3.3)
Children with fever receiving antimalarial drugs (%)	30	53 (32 to 61)	–4.9 (–7.8 to –0.2)
Diarrhoea treatment (%)	25	39 (28 to 45)	2.2 (–1.4 to 5.5)
Physicians (rate per 100 000 persons)	23	5.7 (2.9 to 12.0)	1.2 (–0.6 to 5.1)
Child survival interventions			
Children 1 year old receiving all basic vaccinations (%)	26	41 (29 to 65)	2.7 (1.1 to 5.1)
Exclusive breastfeeding (%)	36	25 (14 to 37)	4.0 (–0.2 to 8.2)
Measles immunisation (%)	46	69 (49 to 76)	1.3 (0.3 to 2.5)
Use of insecticide-treated bed nets (%)	30	2 (1 to 4)	29.0 (24.0 to 35.0)
Vitamin A supplementation coverage (%)	37	91 (78 to 96)	–0.3 (–2.0 to 0.7)
Clinical and health conditions			
Adult prevalence of HIV (%)	43	4 (1 to 8)	–0.8 (–3.6 to 1.7)
Low-birthweight babies (%)	30	13 (11 to 17)	–0.3 (–3.7 to 1.0)
Malnutrition prevalence (height for age) (%)	34	40 (34 to 48)	–1.1 (–2.4 to 0.2)
Malnutrition prevalence (weight for age) (%)	34	22 (18 to 27)	–1.8 (–3.3 to –1.0)
Prevalence of wasting (%)	34	9 (7 to 12)	–1.9 (–5.3 to 0.5)
Governance and financing			
External resources for health (%)	45	13 (4.6 to 20)	6.2 (–1.1 to 9.8)
Health expenditure (% of GDP) (%)	45	5 (4 to 6)	1.4 (0.5 to 3.2)
Health expenditure (% of government expenditures) (%)	45	9 (7 to 10)	2.5 (–0.4 to 3.6)
Net bilateral aid flows from DAC donors (US\$ in millions)	46	120 (40 to 260)	8.7 (4.9 to 15.0)
Net ODA received (% of GNI) (%)	46	9 (4 to 13)	–0.9 (–5.6 to 5.3)
Net ODA received per capita (US\$)	46	23 (15 to 41)	7.1 (2.4 to 13.0)
Out-of-pocket health expenditure (%)	45	43 (26 to 58)	–1.7 (–4.0 to –0.6)
Public spending on education (% of GDP) (%)	34	4 (3 to 5)	2.0 (0.2 to 3.8)
Public spending on education (% of government expenditure) (%)	25	16 (14 to 19)	1.6 (–0.2 to 3.0)
Safety and Rule of Law (0 to 100)	46	54 (43 to 66)	–0.0 (–0.9 to 0.4)
Other factors			
Households with television (%)	24	18 (6 to 29)	6.5 (4.0 to 10.0)
Internet users (rate per 100 persons)	46	0.2 (0.1 to 0.5)	29.0 (22.0 to 38.0)
Population coverage of cell phones (%)	26	37 (17 to 82)	8.2 (1.3 to 15.0)

ARC, annual rate of change; ARI, acute respiratory infection; DAC, Development Assistance Committee; GNI, gross national income; ODA, Official Development Assistance; PPP int'l \$, purchasing power parity of GDP converted to international dollars.

in fertility rate as the transition enters the final stages such the fertility rates once again come to parallel mortality rates.

Few other studies have explored the association between changes in health, economic and financing

factors and ARRs in child mortality. Fotso *et al*²¹ studied U5M ARRs from 22 African countries using data from 1988 to 2004. Larger increases in access to clean water and receipt of basic vaccinations (BCG); diphtheria, pertussis, tetanus; measles; and polio) were associated with




Figure 2 Correlation between under-five mortality ARR and completeness of indicator data for 46 countries in the WHO African Region for the period 2000–2013. The coefficient of 0.09 suggests that for every additional 11 factors reported by a country, there is a corresponding 1% increase in the ARR.

larger ARRs, while higher increases in urban populations were associated with lower ARRs. Our analyses found increases in access to clean water and urban population to be related to ARR, but these were not statistically significant. Findings from Fotso *et al.*²¹ are not directly comparable to ours, however, given their earlier time period under study, inclusion of fewer countries and focus on urban areas and no multivariable analyses.

Binkin *et al.*¹² studied U5M ARRs from 19 African countries using data primarily from the late 1990s to mid-2000s and grouping specific interventions into composite indices representing antenatal care, access to delivery services, seeking treatment, immunisations, breastfeeding and nutritional status. Of these, antenatal care, access to delivery services (including a skilled attendant at delivery, delivery in a health facility and/or caesarean section), and improved nutritional status (including reductions in underweight, stunting and/or wasting) were associated with higher ARRs. We did not find these associations in our analysis. And unlike our finding of increased coverage of seeking ARI treatment to be associated with higher ARRs, Binkin *et al.*¹² did not find seeking health treatment (including ARI treatment and diarrhoea treatment) to be associated with ARR. Again, however, their analyses differed from ours in terms of time period of study and number of countries included.

Lawn *et al.*²² assessed how annual rates of change in 11 factors representing context (gross national income, general and adolescent fertility rate, female literacy, total health expenditure, government effectiveness, political

stability and female HIV prevalence) and coverage (skilled birth attendance, tetanus prevention at birth and diphtheria, pertussis, tetanus vaccine coverage) were associated with neonatal mortality (not U5M) ARR between 2000 and 2010. Of these, only improved coverage of tetanus protection at birth was found to be associated with lower neonatal mortality ARR, a counter-intuitive finding. The authors argued that the paradox arose from the fact that the nations investing the most in scaling up tetanus coverage were those with the weakest health systems, high neonatal mortality and lower coverage of interventions. Many of the contextual factors analysed by Lawn *et al.* were found to have small and statistically non-significant associations in our analysis of U5M ARR.

Other ecological studies of country-level factors and under-five or infant mortality rates (not ARRs) have found the following to be associated with lower mortality in Africa: having a skilled attendant at delivery,²³ exclusive breast feeding,²⁴ prevention of mother-to-child transmission of HIV and antiretroviral therapy for children,²⁵ diphtheria, pertussis, tetanus or measles vaccination,²⁵ use of insecticide treated nets,²⁵ HIV,^{26 27} health expenditures,²⁶ income,^{23 27} urban population prevalence,²⁸ access to clean water,^{25 29} female education,^{23 27 28} fertility rate²⁸ and maternal mortality.²⁹ Stunting²⁵ and underweight²⁴ were associated with increased mortality. While our study found associations between U5M ARRs and changes in only some of these factors, we note that a lack of association between a factor's ARC and mortality ARR does not imply that the intervention or other factor is not associated with reduced mortality. Rather, it means that more rapid scale-up or change in a particular factor does not appear to be associated with a more rapid decline in mortality based on the ARR. During the study period, nearly every African country had declining U5M. A non-significant ARC-ARR association could present if such a decline was relatively constant, regardless of whether there was a slow or rapid change in the factor over the corresponding period. It also highlights the need for further research on delivery and subsequent measurement, including assessing quality, of these proven interventions.^{19 30}

Our findings highlight the role of financing, economy, and the broader health system to more rapidly reduce an already declining, albeit slowly, U5M rate above and beyond the contribution of individual interventions. As the Millennium Development Goals give way to Sustainable Development Goals for 2030, these factors must be addressed to enable further increases in maternal, neonatal and child health interventions.³¹ Model projections suggest that 3.8 million children will still die in 2030 if current U5M rate ARRs remain steady, and that 2.4 million will die even if ARRs increase substantially.²⁷ As U5M rates continue to decline, increasing attention must be paid to neonatal mortality which now accounts for 44% of all deaths in children under 5 years

Table 3 Crude and adjusted associations for select factors* and under-five mortality annual rate of reduction, 2000–2013

		Crude			Adjusted†		
	N‡	β	90% CI	p Value	β	90% CI	p Value
Sociodemographics							
Adult female literacy rate (%)	33	0.02	(−0.26 to 0.31)	0.39	0.11	(−0.19 to 0.41)	0.32
Female labour participation rate (%)	42	−1.07	(−2.42 to 0.29)	0.17	−0.32	(−1.56 to 0.92)	0.36
Fertility rate (births per woman)	42	0.78	(0.26 to 1.31)	0.02	0.54	(0.07 to 1.02)	0.07
Human development index (0–1)§	35	0.12	(0.04 to 0.20)	0.02	0.09	(0.01 to 0.17)	0.07
Improved water source (%)	42	0.40	(−0.05 to 0.85)	0.14	0.36	(−0.07 to 0.78)	0.16
Labour force participation rate among women (%)	42	−0.33	(−1.19 to 0.53)	0.33	0.19	(−0.57 to 0.96)	0.36
Urban population prevalence (%)	42	0.53	(−0.03 to 1.09)	0.12	0.39	(−0.15 to 0.93)	0.20
Maternal Health							
Adult female mortality rate (rate per 1000 persons)	42	−0.49	(−0.83 to −0.16)	0.03	−0.25	(−0.61 to 0.10)	0.20
Maternal mortality ratio (rate per 100 000 births)¶	35	−0.57	(−0.80 to −0.35)	<0.01	−0.47	(−0.69 to −0.24)	<0.01
Pregnant women receiving prenatal care (%)	35	0.09	(−0.30, 0.48)	0.37	0.09	(−0.24 to 0.41)	0.36
Access to Health Care							
Seeking ARI treatment (%)¶	31	0.26	(0.10 to 0.42)	0.02	0.22	(0.09 to 0.35)	0.01
Births attended by skilled health staff (%)¶	34	0.21	(−0.03 to 0.45)	0.14	0.18	(−0.07 to 0.43)	0.19
Clinical and Health Conditions							
Adult prevalence of HIV (%)	42	−0.13	(−0.25 to −0.01)	0.07	−0.09	(−0.20 to 0.02)	0.17
Malnutrition prevalence (height for age) (%)¶	32	0.11	(−0.15 to 0.38)	0.31	0.15	(−0.10 to 0.40)	0.24
Governance and Financing							
Health expenditure (% of GDP) (%)	42	0.24	(0.07 to 0.40)	0.03	0.26	(0.11 to 0.41)	0.01
Health expenditure (% of government expenditures) (%)**	42	0.11	(−0.01 to 0.23)	0.12	0.08	(−0.04 to 0.19)	0.22
Net ODA received per capita (US\$)	42	−0.01	(−0.09 to 0.07)	0.39	−0.08	(−0.15 to −0.01)	0.07
Out-of-pocket health expenditure (%)**	42	−0.19	(−0.37 to −0.02)	0.08	−0.19	(−0.34 to −0.03)	0.06
Other							
Households with television (%)	23	0.29	(0.13 to 0.46)	0.01	0.19	(0.05 to 0.33)	0.04

*Includes all factors for which (1) $\beta > 0.20$ or $\beta < -0.20$, or (2) $p < 0.10$. Factors having both are shown in bold. Additional factors of interest are shown if $N \geq 23$. (Results for all factors available in online supplemental table S5).

†All results adjusted for improved water source, health expenditure (% of GDP), adult HIV prevalence, and urban population prevalence unless otherwise noted.

‡N=sample size of the adjusted analyses and is <46 due to missing country data on some of the factors, including those included as covariates.

§Since the range of the HDI is only from 0 to 1, the β represents the change in ARR for each 0.10 increase in HDI ARC.

¶further adjusted for receipt of prenatal care.

** not adjusted for health expenditure (% of GDP).

ARI, acute respiratory infection; GDP, gross domestic product; ODA, Official Development Assistance.

of age (34% in Sub-Saharan African).^{1 32} Facility-based interventions and improving quality care for pregnant women and newborns are needed.³² The same financial, economic and broader health system factors that would increase U5M ARR would also increase reductions in neonatal mortality.

Limitations should be considered when interpreting the findings from this report. First, mortality and indicator data were obtained from secondary sources and some may have suboptimal methods of data collection and model-based estimation, though the World Bank, United Nations Children's Fund, the Interagency Group for Child Mortality Estimation, Demographic and Health Surveys, Multiple Indicator Cluster Surveys and Countdown to 2015 seek to be as authoritative sources as they can. A recent *PLoS Medicine* Collection highlighted the challenges in obtaining valid coverage data from community-based household surveys.¹⁹ Second, the study design was an ecological, study. As such, changes in the factors of interest were occurring simultaneously with changes in mortality, and temporality and causality inferences are limited. Ecological studies are prone to bias and associations at the aggregate level (eg, country) may not reflect the cause–effect relationship at the individual or local level. Nevertheless, we felt this design was appropriate given the interest in achieving MDG4, which is monitored at a country level. Third, the time period for calculating the ARCs did not always fully overlap with the time period for calculating mortality ARRs (eg, ARCs for 2000–2011 may have been calculated from data for 2003 and 2010). We attempted to minimise any potential bias by only considering data from factors for which at least a 5-year period of the 11-year time period were available for calculating ARCs. Finally, the small sample size (at most n=46 countries) limits the validity of many associations and prevents adjustment for many covariates. We cannot exclude residual confounding and low power to detect certain associations. In spite of these limitations, the unique strengths of our study include our broad identification of factors potentially relevant to child survival, beyond those typically used from Demographic and Health Surveys and Countdown to 2015. We also made statistical efforts to minimise bias or distortion from outliers, without excluding them.

There are numerous interventions, health factors and other determinants of health that influence child mortality rates. However, there are very few studies assessing which of these are associated with higher ARR reflecting more rapid declines in U5M. Policy makers now have evidence that can back decisions to propose increasing access to care for children under age five (extrapolating from our ARI access finding), and increasing health investments, notably towards both child and maternal health.

Author affiliations

¹Vanderbilt Institute for Global Health, Nashville, Tennessee, USA

²Department of Medicine, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

³Department of Biostatistics, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

⁴World Health Organization/Regional Office for Africa, Brazzaville, Congo

⁵Department of Health Policy, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

⁶Pediatrics Vanderbilt University School of Medicine, Nashville, Tennessee, USA

Acknowledgements The authors thank Dr Matshidiso Moeti who inspired the conceptualisation of this study and Jessica Islam for her assistance in reviewing country Demographic and Health Surveys and assisting with data entry.

Contributors AMK helped with the study design, acquisition of data, assisted with data analysis, interpretation of results and drafting the manuscript. He had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. MB assisted with acquisition of data, conducted the analysis and critically revised manuscript. CAH contributed to the study design, interpretation of results and drafting the manuscript. KM helped with conception and design of the study and critically revising the manuscript. PH helped with conception and design of the study and critically revising the manuscript. BES assisted with data analysis and critically revising the manuscript. MHA assisted with the study design and interpretation and critically revising the manuscript. TK helped with conception and design of the study and critically revising the manuscript. SHV helped with conception and design of the study and drafting the manuscript. All authors have participated sufficiently in the work to take public responsibility for its content.

Funding Funding for this project was provided by the WHO. Support for data management came from the Vanderbilt Institute for Clinical and Translational Research (grant UL1 TR000445 from the National Center for Advancing Translational Sciences at the National Institutes of Health). At the time of the study, Dr. Kipp was a Scholar with the HIV/AIDS, Substance Abuse, and Trauma Training Program (HA-STTP), at the University of California, Los Angeles; supported through an award from the National Institute on Drug Abuse (R25 DA035692).

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Additional data is available at <http://biostat.mc.vanderbilt.edu/wiki/Main/WHOUnder5Mortality>, or by emailing AMK at aaron.kipp@vanderbilt.edu.

Open Access This is an open access article distributed under the terms of the Creative Commons Attribution IGO License (<https://creativecommons.org/licenses/by-nc/3.0/igo/>), which permits use, distribution, and reproduction for non-commercial purposes in any medium, provided the original work is properly cited. In any reproduction of this article there should not be any suggestion that WHO or this article endorse any specific organisation or products. The use of the WHO logo is not permitted. This notice should be preserved along with the article's original URL.

REFERENCES

1. UNICEF, WHO, World Bank, UNDESA. Levels and Trends in Child Mortality: Report 2014—Estimates Developed by the United Nations Inter-agency Group for Child Mortality Estimation. New York: UNICEF, 2014.
2. UN. *United Nations Millennium Declaration*. United Nations, 2000.
3. Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet* 2003;361:2226–34.
4. Jones G, Steketee RW, Black RE, *et al*. How many child deaths can we prevent this year? *Lancet* 2003;362:65–71.
5. Bryce J, el Arifeen S, Pariyo G, *et al*. Reducing child mortality: can public health deliver? *Lancet* 2003;362:159–64.
6. Victora CG, Wagstaff A, Schellenberg JA, *et al*. Applying an equity lens to child health and mortality: more of the same is not enough. *Lancet* 2003;362:233–41.
7. Claeson M, Gillespie D, Mshinda H, *et al*. Knowledge into action for child survival. *Lancet* 2003;362:323–7.
8. Countdown to 2015. Tracking Progress in Child Survival: the 2005 Report. Geneva: World Health Organization, 2005.

9. Bryce J, Tetteri N, Victora CG, *et al.* Countdown to 2015: tracking intervention coverage for child survival. *Lancet* 2006;368: 1067–76.
10. Countdown to 2015. Building a Future for Women and Children: the 2012 Report. Geneva: World Health Organization, 2012.
11. WHO. *Atlas of health statistics of the African region 2012: health situation analysis of the African region*. Brazzaville, Republic of Congo: World Health Organization Regional Office for Africa, 2012.
12. Binkin N, Chopra M, Simen-Kapeu A, *et al.* Do improvements in outreach, clinical, and family and community-based services predict improvements in child survival? An analysis of serial cross-sectional national surveys. *BMC Public Health* 2011;11:456.
13. The World Bank Data Catalog. <http://datacatalog.worldbank.org/>
14. Measure DHS country reports. <http://dhsprogram.com/Where-We-Work/Country-List>
15. Huber PJ. *Robust statistics*. New York: Wiley, 1981.
16. Hampel FR, Ronchetti EM, Rousseeuw PJ, *et al.* *Robust statistics : the approach based on influence functions*. New York: Wiley, 1986.
17. Venables WN, Ripley BD. *Modern applied statistics with S*. New York: Springer, 2002.
18. Campbell H, El Arifeen S, Hazir T, *et al.* Measuring coverage in MNCH: challenges in monitoring the proportion of young children with pneumonia who receive antibiotic treatment. *PLoS Med* 2013;10:e1001421.
19. Bryce J, Arnold F, Blanc A, *et al.* Measuring coverage in MNCH: new findings, new strategies, and recommendations for action. *PLoS Med* 2013;10:e1001423.
20. Cutts FT, Izurieta HS, Rhoda DA. Measuring coverage in MNCH: design, implementation, and interpretation challenges associated with tracking vaccination coverage using household surveys. *PLoS Med* 2013;10:e1001404.
21. Fotso JC, Ezeh AC, Madise NJ, *et al.* Progress towards the child mortality millennium development goal in urban Sub-Saharan Africa: the dynamics of population growth, immunization, and access to clean water. *BMC Public Health* 2007;7:218.
22. Lawn JE, Kinney MV, Black RE, *et al.* Newborn survival: a multi-country analysis of a decade of change. *Health Policy Plan* 2012;27(Suppl 3):iii6–28.
23. Olafsdottir AE, Reidpath DD, Pokhrel S, *et al.* Health systems performance in Sub-Saharan Africa: governance, outcome and equity. *BMC Public Health* 2011;11:237.
24. Abrahams Z, McHiza Z, Steyn NP. Diet and mortality rates in Sub-Saharan Africa: stages in the nutrition transition. *BMC Public Health* 2011;11:801.
25. Akachi Y, Atun R. Effect of investment in malaria control on child mortality in Sub-Saharan Africa in 2002–2008. *PLoS ONE* 2011;6:e21309.
26. Novignon J, Olakojo SA, Nonvignon J. The effects of public and private health care expenditure on health status in Sub-Saharan Africa: new evidence from panel data analysis. *Health Econ Rev* 2012;2:22.
27. Wang H, Liddell CA, Coates MM, *et al.* Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:957–79.
28. Chuang YC, Sung PW, Chao HJ, *et al.* A longitudinal ecological study of the influences of political, economic, and health services characteristics on under-five mortality in less-developed countries. *Health Place* 2013;23:111–21.
29. Sartorius BKD, Sartorius K. Global infant mortality trends and attributable determinants—an ecological study using data from 192 countries for the period 1990–2011. *Popul Health Metrics* 2014;12:29.
30. Wazny K, Zipursky A, Black R, *et al.* Setting research priorities to reduce mortality and morbidity of childhood diarrhoeal disease in the next 15 years. *PLoS Med* 2013;10:e1001446.
31. Requejo JH, Bhutta ZA. The post-2015 agenda: staying the course in maternal and child survival. *Arch Dis Child* 2015;100(Suppl 1): S76–81.
32. Wardlaw T, You D, Hug L, *et al.* UNICEF Report: enormous progress in child survival but greater focus on newborns urgently needed. *Reprod Health* 2014;11:82.

SUPPLEMENTAL MATERIAL

Table S1. Factors assessed in the ecological analysis.

Category	Indicator name ^a
Socio-demographic (18 indicators listed)	
Age	<ul style="list-style-type: none"> • Age at first marriage, female • Teenage mothers (% of women ages 15-19 who have had children or are currently pregnant) • Age at first birth of women 20-49 yrs
Education	<ul style="list-style-type: none"> • Years of education completed among women 15-49 yrs • Percent of women 15-49 yrs old with at least some secondary education
Literacy	<ul style="list-style-type: none"> • Literacy rate, females (% of females ages 15 and above)
Children	<ul style="list-style-type: none"> • Number of living children per household • Fertility rate, total (births per woman)^c
Employment	<ul style="list-style-type: none"> • Labor force participation rate, female (% of female population ages 15-64) • Labor force, female (% of total labor force)
Water and Sanitation	<ul style="list-style-type: none"> • Access to improved water source (% of population with access)^b • Access to improved sanitation facilities (% of population with access)^b
Socio-economic status	<ul style="list-style-type: none"> • Gini index^d • Poverty headcount ratio at national poverty line (% of population) • Households with television (%) • Human development index (HDI)^e • Gross Domestic Product (GDP) per capita (PPP Int'l \$)
Rural/urban residence	<ul style="list-style-type: none"> • Urban population (% of total)
Maternal health (6 indicators listed)	
Family planning	<ul style="list-style-type: none"> • Unmet need for contraception (% of married women ages 15-49)^b
Antenatal care	<ul style="list-style-type: none"> • Pregnant women receiving prenatal care (%)^b • Pregnant women receiving prenatal care of at least four visits (%)^b • Percent of women with live births who received antenatal care during first 5 months of pregnancy
Maternal mortality	<ul style="list-style-type: none"> • Maternal mortality ratio (per 100,000 live births)^c • Mortality rate, adult, female (per 1,000 female adults)
Access to health care (16 indicators listed)	
Availability	<ul style="list-style-type: none"> • Physicians (per 1,000 persons)^c • Nurses and midwives (per 1,000 persons)^c • Nurses (per 1,000 persons) • Community health workers (per 1,000 persons)

	<ul style="list-style-type: none"> • Hospital beds (per 1,000 persons) • Births attended by skilled health staff (% of total)^b • % of births delivered by caesarean section^b
Utilization	<ul style="list-style-type: none"> • % of births delivered at a health facility • % of births with a postnatal visit within the first 2 days after delivery^b • Seeking ARI treatment (% of children under 5 taken to a health provider)^b
Treatment	<ul style="list-style-type: none"> • % of children 0-59 months of age with ARI who received antibiotics^b • Diarrhea treatment (% of children under 5 receiving oral rehydration and continued feeding)^b • Children with fever receiving antimalarial drugs (% of children under age 5 with fever)^b • Use of preventive treatment for malaria (% of pregnant women)^b • Adoption of low osmolarity Oral Rehydration Salts and zinc for management of diarrhea^c • Community treatment of pneumonia with antibiotics^c
Child survival interventions (7 indicators listed)	
PMTCT	<ul style="list-style-type: none"> • HIV positive pregnant women receiving antiretrovirals, using WHO/UNAIDS methodology (%)^b • Antiretroviral therapy coverage (% of persons with advanced HIV infection)^b
Breastfeeding	<ul style="list-style-type: none"> • Exclusive breastfeeding (% of children under 6 months)^b
Vitamin A supplementation	<ul style="list-style-type: none"> • Vitamin A supplementation coverage rate (% of children ages 6-59 months)^b
Immunizations	<ul style="list-style-type: none"> • Immunization, measles (% of children ages 12-23 months)^b • % of 12-23 month old children receiving all basic vaccinations^{b,f}
Bed nets	<ul style="list-style-type: none"> • Use of insecticide-treated bed nets (% of under-5 population)^b
Clinical and health conditions (7 indicators listed)	
HIV	<ul style="list-style-type: none"> • Prevalence of HIV, total (% of population ages 15-49) • Female adults with HIV (% of population ages 15+ with HIV) • Prevalence of HIV, female (% ages 15-24)
Preterm/birth weight	<ul style="list-style-type: none"> • Low-birth weight babies (% of births)^c
Malnutrition	<ul style="list-style-type: none"> • Malnutrition prevalence, height for age (% of children under 5)^c • Malnutrition prevalence, weight for age (% of children under 5)^c • Prevalence of wasting (% of children under 5)^c
Governance and financing (11 indicators listed)	
Health financing	<ul style="list-style-type: none"> • Health expenditure, total (% of GDP)^c • Health expenditure, public (% of government expenditure)^c • Out-of-pocket health expenditure (% of total expenditure on health)^c
Education financing	<ul style="list-style-type: none"> • Public spending on education, total (% of GDP) • Public spending on education, total (% of government expenditure)
Donor funding	<ul style="list-style-type: none"> • External resources for health (% of total expenditure on health) • Net bilateral aid flows from DAC donors, Total (current US\$)^g • Net official development aid (ODA) received (% of Gross National Income)^{c,h}

	<ul style="list-style-type: none"> • Net ODA received per capita (current US\$)^{c,h} • Net ODA received (constant 2010 US\$)^{c,h}
Quality of governance	<ul style="list-style-type: none"> • Safety and Rule of Law (Ibrahim index with 4 indicators)ⁱ
Other (5 indicators listed)	
Country infrastructure	<ul style="list-style-type: none"> • Road density (km of road per 100 sq. km of land area) • Roads, paved (% of total roads) • Access to electricity (% of population) • Population coverage of mobile cellular telephony (%) • Internet users (per 100 persons)

^a 10 indicators in bold were not included in the database due to 100% insufficient data (see methods); analysis dataset has 61 indicators.

^b one of the 27 Countdown to 2015 intervention indicators (or closely related)

^c one of the Countdown to 2015 country profile indicators (or closely related)

^d The GINI index is a measure of income inequality, where 0 represents perfect equality and 100 represents perfect inequality

^e The Human Development Index is a composite measure of development incorporating life expectancy at birth, mean years of schooling and expected years of schooling, and gross national income per capita, where 0 represents very low development and 1 represents very high development

^f Includes BCG, measles and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

^g DAC, Development Assistance Committee. DAC members are Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Japan, Republic of Korea, Luxembourg, Netherlands, New Zealand, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States, and European Union Institutions

^h ODA, Official Development Assistance. ODA includes loans and grants made by DAC members, non-DAC countries, and multilateral institutions

ⁱ Ibrahim index is a composite measure of a countries provision of political, social and economic public goods and services. It is comprised of four indicates: Personal safety, Rule of law, Accountability and corruption and, National security. The index ranges from 0 to 100, with higher values representing better governance.

Table S2. Detailed description of data sources for each of the 70 factors assessed in the ecological study.

Indicator ^a	Source for data analysis ^j	Primary source reported by World Bank Data Catalogue (if applicable) ^k	Additional notes ^k
Socio-demographic (18 indicators listed)			
Age at first marriage, female	WB-HNPS	UN, Department of Economic and Social Affairs, Population Division. World Marriage Data.	
Teenage mothers (% of women ages 15-19 who have had children or are currently pregnant)	WB-WDI	DHS (Macro Int.)	
Age at first birth of women 20-49 yrs	Country DHS		
Years of education completed among women 15-49 yrs	Country DHS		
Percent of women 15-49 yrs old with at least some secondary education	Country DHS		
Literacy rate, females (% of females ages 15 and above)	WB-WDI	UN Educational, Scientific, and Cultural Organization (UNESCO) Institute for Statistics.	
Number of living children per household	Country DHS		
Fertility rate, total (births per woman) ^c	WB-WDI	(1) UN Population Division. World Population Prospects, (2) United Nations Statistical Division. Population and Vital Statistics Report (various years), (3) Census reports and other statistical publications from national statistical offices	Annual data series are interpolated from 5-year period data; for countries without vital registration systems fertility rates are extrapolated from earlier censuses or other surveys. If no empirical information available, a model is used to estimate the share of births to adolescents.
Labor force participation rate, female (% of female population ages 15-64)	WB-WDI	International Labour Organization, Key Indicators of the Labour Market database.	ILO modeled estimates using both nationally reported and imputed data.
Labor force, female (% of total labor force)	WB-WDI	International Labour Organization, using World Bank population estimates.	
Improved water source (% of population with access) ^b	WB-WDI	WHO and UN Children's Fund, Joint Measurement Programme http://www.wssinfo.org	Linear trend line through data points to provide estimates for all years between 1990 and 2015
Improved sanitation facilities (% of population with access) ^b	WB-WDI		
GINI index ^d	WB-WDI	World Bank, Development Research Group.	Data based on household survey data obtained from government statistical agencies and World Bank country

			departments. (http://iresearch.worldbank.org/PovcalNet/index.htm).
Poverty headcount ratio at national poverty line (% of population)	WB-WDI	World Bank, Global Poverty Working Group.	Data are based on World Bank's country poverty assessments and country Poverty Reduction Strategies.
Households with television (%)	WB-ADI	International Telecommunication Union	
Human development index (HDI) ^e	WB-ADI	UN Development Program, Human Development Reports http://hdr.undp.org/en/statistics/data/	
Gross Domestic Product (GDP) per capita (PPP Int'l \$)	WB-WDI	World Bank, International Comparison Program database.	
Urban population (% of total)	WB-WDI	UN, World Urbanization Prospects.	The indicator is calculated using World Bank population estimates and urban ratios from the UN World Urbanization Prospects. Annual data series are interpolated from census and other national population data using demographic modeling
Maternal health (6 indicators listed)			
Unmet need for contraception (% of married women ages 15-49) ^b	WB-WDI	Household surveys, including DHS (Macro Int.) and MICS (UNICEF)	
Pregnant women receiving prenatal care (%) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	
Pregnant women receiving prenatal care of at least four visits (%) ^b	WB-HNPS		
Percent of women with live births who received antenatal care during first 5 months of pregnancy	Country DHS		
Maternal mortality ratio (per 100,000 live births) ^c	WB-WDI	WHO, UNICEF, UNFPA, The World Bank, and the United Nations Population Division. Trends in Maternal Mortality: 1990 to 2013. Geneva, World Health Organization, 2014	Modeled estimates are based on the Maternal Mortality Estimation Inter-Agency Group (WHO, UNICEF, UNFPA, and World Bank) and use a multilevel regression model utilizing national maternal mortality data and socioeconomic information, including fertility, birth attendants, and GDP.
Mortality rate, adult, female (per 1,000 female adults)	WB-WDI	(1) UN Population Division. World Population Prospects. New York, UN, Department of Economic and Social Affairs	Annual data series are linear interpolations from 5-year period data; for countries without vital registration

		http://esa.un.org/wpp/unpp/panel_population.htm , (2) University of California, Berkeley, and Max Planck Institute for Demographic Research. Human Mortality Database. www.mortality.org or www.humanmortality.de	systems fertility rates are extrapolated from earlier censuses or other surveys.
Access to health care (16 indicators listed)			
Physicians (per 1,000 people) ^c	WB-WDI	WHO, Global Atlas of the Health Workforce http://apps.who.int/globalatlas/ .	
Nurses and midwives (per 1,000 people) ^c	WB-HNPS		
Nurses (per 1,000 people)	WB-WDI		100% insufficient for analysis
Community health workers (per 1,000 people)	WB-WDI	WHO, OECD, supplemented by country data.	
Hospital beds (per 1,000 people)	WB-WDI	Data after 2005 are extracted from the World Health Statistics Table 6 published by WHO. WHS data is based on PAHO basic indicators (2011) www.paho.org/English/SHA/coredata/tabulator/newTabulator.htm	
Births attended by skilled health staff (% of total) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	For many countries, includes MICS data reanalyzed by UNICEF
% of births delivered by caesarean section ^b	Country DHS		
Adoption of low osmolarity Oral Rehydration Salts and zinc for management of diarrhea^c	Countdown 2015 reports		100% insufficient for analysis
Community treatment of pneumonia with antibiotics^c	Countdown 2015 reports		100% insufficient for analysis
% of births delivered at a health facility	Country DHS		
% of births with a postnatal visit within the first 2 days after delivery ^b	Country DHS		
Seeking ARI treatment (% of children under 5 taken to a health provider) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	For many countries, includes DHS or MICS data reanalyzed by UNICEF
% of children 0-59 months of age with ARI who received antibiotics ^b	Country DHS		100% insufficient for analysis
Diarrhea treatment (% of children under 5 receiving oral rehydration and continued feeding) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	For many countries, includes DHS or MICS data reanalyzed by UNICEF
Children with fever receiving	WB-WDI		

antimalarial drugs (% of children under age 5 with fever) ^b			
Use of preventive treatment for malaria (% of pregnant women) ^b	WB-HNPS	DHS (Macro Int.)	
Child survival interventions (7 indicators listed)			
HIV positive pregnant women receiving antiretrovirals, using WHO/UNAIDS methodology (%) ^b	WB-ADI	UNAIDS and the WHO's Report on the Global AIDS Epidemic. http://data.unaids.org/pub/GlobalReport/2008/	100% insufficient for analysis
Antiretroviral therapy coverage (% of people with advanced HIV infection) ^b	WB-WDI	UNAIDS and the WHO's Report on the Global AIDS Epidemic.	100% insufficient for analysis
Exclusive breastfeeding (% of children under 6 months) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	For some countries, includes MICS data reanalyzed by UNICEF
Vitamin A supplementation coverage rate (% of children ages 6-59 months) ^b	WB-WDI	UN Children's Fund, State of the World's Children.	
Immunization, measles (% of children ages 12-23 months) ^b	WB-WDI	WHO and UNICEF http://www.who.int/immunization_monitoring/routine/en/	
% of 12-23 month old children receiving all basic vaccinations ^{b,f}	Country DHS		
Use of insecticide-treated bed nets (% of under-5 population) ^b	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	
Clinical and health conditions (7 indicators listed)			
Prevalence of HIV, total (% of population ages 15-49)	WB-WDI	UNAIDS and the WHO's Report on the Global AIDS Epidemic.	All years are model based estimates from 2012 report
Female adults with HIV (% of population ages 15+ with HIV)	WB-HNPS		100% insufficient for analysis
Prevalence of HIV, female (% ages 15-24)	WB-HNPS		100% insufficient for analysis
Low-birth weight babies (% of births) ^c	WB-WDI	UNICEF, State of the World's Children, Childinfo, and DHS (Macro Int.)	For many countries, includes DHS or MICS data reanalyzed by UNICEF
Malnutrition prevalence, height for age (% of children under 5) ^c	WB-WDI	WHO, Global Database on Child Growth and Malnutrition. Aggregation is based on UNICEF/WHO Joint Global Malnutrition Analysis Date Set 2011 and additional analysis by UNICEF.	MICS and other UNICEF surveys, DHS, national nutrition surveys
Malnutrition prevalence, weight for age (% of children under 5) ^c	WB-WDI		
Prevalence of wasting (% of children under 5) ^c	WB-WDI		
Governance and financing (11 indicators listed)			

Health expenditure, total (% of GDP) ^c	WB-WDI	WHO National Health Account database http://apps.who.int/nha/database	Many low-income countries use DHS or MICS funded by donors to obtain health system data.
Health expenditure, public (% of government expenditure) ^c	WB-WDI	WHO Global Health Expenditure database http://apps.who.int/nha/database	
Out-of-pocket health expenditure (% of total expenditure on health) ^c	WB-HNPS	WHO National Health Account database http://apps.who.int/nha/database/DataExplore/rRegime.aspx	Many low-income countries use DHS or MICS funded by donors to obtain health system data.
Public spending on education, total (% of GDP)	WB-WDI	UN Educational, Scientific, and Cultural Organization (UNESCO) Institute for Statistics.	
Public spending on education, total (% of government expenditure)	WB-WDI		
External resources for health (% of total expenditure on health)	WB-WDI	WHO National Health Account database http://apps.who.int/nha/database for the most recent updates	Many low-income countries use DHS or MICS funded by donors to obtain health system data.
Net bilateral aid flows from DAC donors, Total (current US\$) ^g	WB-WDI	DAC of the Organisation for Economic Co-operation and Development, Geographical Distribution of Financial Flows to Developing Countries, Development Co-operation Report, and International Development Statistics database. www.oecd.org/dac/stats/idsonline	
Net ODA received (constant 2010 US\$)^{c,h}	WB-WDI		100% insufficient for analysis
Net official development aid (ODA) received (% of Gross National Income) ^{c,h}	WB-WDI		
Net ODA received per capita (current US\$) ^{c,h}	WB-WDI	DAC of the Organisation for Economic Co-operation and Development, Geographical Distribution of Financial Flows to Developing Countries, Development Co-operation Report, and International Development Statistics database. www.oecd.org/dac/stats/idsonline . World Bank estimates are used for the denominator.	
Safety and Rule of Law (Ibrahim index with 4 indicators) ⁱ	WB-ADI	Mo Ibrahim Foundation, electronic files and web site.	
Road density (km of road per 100 sq. km of land area)	WB-WDI	International Road Federation, World Road Statistics and electronic files	
Roads, paved (% of total roads)	WB-WDI		
Access to electricity (% of population)	WB-WDI	International Energy Agency, World Energy Outlook 2010.	100% insufficient for analysis
Population coverage of mobile cellular telephony (%)	WB-ADI	International Telecommunication Union	
Internet users (per 100 people)	WB-WDI		

^a 10 indicators in bold were not included in the database due to 100% insufficient data (see methods); analysis dataset has 60 indicators.

^b one of the 27 Countdown to 2015 intervention indicators (or closely related)

^c one of the Countdown to 2015 country profile indicators (or closely related)

^d The GINI index is a measure of income inequality, where 0 represents perfect equality and 100 represents perfect inequality

^e The Human Development Index is a composite measure of development incorporating life expectancy at birth, mean years of schooling and expected years of schooling, and gross national income per capita, where 0 represents very low development and 1 represents very high development

^f Includes BCG, measles and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

^g DAC, Development Assistance Committee. DAC members are Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Japan, Republic of Korea, Luxembourg, Netherlands, New Zealand, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States, and European Union Institutions

^h ODA, Official Development Assistance. ODA includes loans and grants made by DAC members, non-DAC countries, and multilateral institutions

ⁱ Ibrahim index is a composite measure of a countries provision of political, social and economic public goods and services. It is comprised of four indicates: Personal safety, Rule of law, Accountability and corruption and, National security. The index ranges from 0 to 100, with higher values representing better governance.

^j Data source for the current analysis: WB, World Bank Data Catalogue; WDI, World Development Index; ADI, Africa Development Index; HNPS, Health Nutrition & Population Statistics; Country DHS indicates data were obtained directly from each country's Demographic and Health Survey; Countdown to 2015 indicates data were obtained directly from each country's Country Profile in the Countdown to 2015 reports.

^k Summarized from the indicator/country metadata provided in the World Bank Data Catalogue. See metadata in World Bank Data Catalogue for more details (Data Catalogue code listed in Supplemental Table S3). <http://databank.worldbank.org>.

Table S3. World Bank Data Catalogue source and code for each of the 70 factors assessed in the ecological study.

Indicator ^a	Source for Data analysis ^j	World Bank Data Catalogue Code ^k
Socio-demographic (18 indicators listed)		
Age at first marriage, female	WB-HNPS	SP.DYN.SMAM.FE
Teenage mothers (% of women ages 15-19 who have had children or are currently pregnant)	WB-WDI	SP.MTR.1519.ZS
Age at first birth of women 20-49 yrs	Country DHS	
Years of education completed among women 15-49 yrs	Country DHS	
Percent of women 15-49 yrs old with at least some secondary education	Country DHS	
Literacy rate, females (% of females ages 15 and above)	WB-WDI	SE.ADT.LITR.FE.ZS
Number of living children per household	Country DHS	
Fertility rate, total (births per woman) ^c	WB-WDI	SP.DYN.TFRT.IN
Labor force participation rate, female (% of female population ages 15-64)	WB-WDI	SL.TLF.CACT.FE.ZS
Labor force, female (% of total labor force)	WB-WDI	SL.TLF.TOTL.FE.ZS
Improved water source (% of population with access) ^b	WB-WDI	SH.H2O.SAFE.ZS
Improved sanitation facilities (% of population with access) ^b	WB-WDI	SH.STA.ACSN
GINI index ^d	WB-WDI	SI.POV.GINI
Poverty headcount ratio at national poverty line (% of population)	WB-WDI	SI.POV.NAHC
Households with television (%)	WB-ADI	IT.TVS.HOUS.ZS
Human development index (HDI) ^e	WB-ADI	UNDP.HDI.XD
Gross Domestic Product (GDP) per capita (PPP Int'l \$)	WB-WDI	NY.GDP.PCAP.PP.CD
Urban population (% of total)	WB-WDI	SP.URB.TOTL.IN.ZS
Maternal health (6 indicators listed)		
Unmet need for contraception (% of married women ages 15-49) ^b	WB-WDI	SP.UWT.TFRT
Pregnant women receiving prenatal care (%) ^b	WB-WDI	SH.STA.ANVC.ZS
Pregnant women receiving prenatal care of at least four visits (%) ^b	WB-HNPS	SH.STA.ANV4.ZS
Percent of women with live births who received antenatal care during first 5 months of pregnancy	Country DHS	
Maternal mortality ratio (per 100,000 live births) ^c	WB-WDI	SH.STA.MMRT
Mortality rate, adult, female (per 1,000 female adults)	WB-WDI	SP.DYN.AMRT.FE
Access to health care (16 indicators listed)		
Physicians (per 1,000 people) ^c	WB-WDI	SH.MED.PHYS.ZS
Nurses and midwives (per 1,000 people) ^c	WB-HNPS	SH.MED.NURS.P3
Nurses (per 1,000 people)		
Community health workers (per 1,000 people)	WB-WDI	SH.MED.NUMW.P3
Hospital beds (per 1,000 people)	WB-WDI	SH.MED.BEDS.ZS
Births attended by skilled health staff (% of total) ^b	WB-WDI	SH.STA.BRTC.ZS
% of births delivered by caesarean section ^b	Country DHS	
Adoption of low osmolarity Oral Rehydration Salts and zinc for management of diarrhea^c	Countdown 2015	

Community treatment of pneumonia with antibiotics^c	Countdown 2015	
% of births delivered at a health facility	Country DHS	
% of births with a postnatal visit within the first 2 days after delivery ^b	Country DHS	
Seeking ARI treatment (% of children under 5 taken to a health provider) ^b	WB-WDI	SH.STA.ARIC.ZS
% of children 0-59 months of age with ARI who received antibiotics ^b	Country DHS	
Diarrhea treatment (% of children under 5 receiving oral rehydration and continued feeding) ^b	WB-WDI	SH.STA.ORCF.ZS
Children with fever receiving antimalarial drugs (% of children under age 5 with fever) ^b	WB-WDI	SH.MLR.TRET.ZS
Use of preventive treatment for malaria (% of pregnant women) ^b	WB-HNPS	SH.MLR.PREG.ZS
Child survival interventions (7 indicators listed)		
HIV positive pregnant women receiving antiretrovirals, using WHO/UNAIDS methodology (%)^b	WB-ADI	SH.HIV.PREG.VIRAL S.ZS
Antiretroviral therapy coverage (% of people with advanced HIV infection)^b	WB-WDI	SH.HIV.ARTC.ZS
Exclusive breastfeeding (% of children under 6 months) ^b	WB-WDI	SH.STA.BFED.ZS
Vitamin A supplementation coverage rate (% of children ages 6-59 months) ^b	WB-WDI	SN.ITK.VITA.ZS
Immunization, measles (% of children ages 12-23 months) ^b	WB-WDI	SH.IMM.MEAS
% of 12-23 month old children receiving all basic vaccinations ^{b,f}	Country DHS	
Use of insecticide-treated bed nets (% of under-5 population) ^b	WB-WDI	SH.MLR.NETS.ZS
Clinical and health conditions (7 indicators listed)		
Prevalence of HIV, total (% of population ages 15-49)	WB-WDI	SH.DYN.AIDS.ZS
Female adults with HIV (% of population ages 15+ with HIV)	WB-HNPS	SH.DYN.AIDS.FE.ZS
Prevalence of HIV, female (% ages 15-24)	WB-HNPS	SH.HIV.1524.FE.ZS
Low-birth weight babies (% of births) ^c	WB-WDI	SH.STA.BRTW.ZS
Malnutrition prevalence, height for age (% of children under 5) ^c	WB-WDI	SH.STA.STNT.ZS
Malnutrition prevalence, weight for age (% of children under 5) ^c	WB-WDI	SH.STA.MALN.ZS
Prevalence of wasting (% of children under 5) ^c	WB-WDI	SH.STA.WAST.ZS
Governance and financing (11 indicators listed)		
Health expenditure, total (% of GDP) ^c	WB-WDI	SH.XPD.TOTL.ZS
Health expenditure, public (% of government expenditure) ^c	WB-WDI	SH.XPD.PUBL.GX.ZS
Out-of-pocket health expenditure (% of total expenditure on health) ^c	WB-HNPS	SH.XPD.OOPC.TO.ZS
Public spending on education, total (% of GDP)	WB-WDI	SE.XPD.TOTL.GD.ZS
Public spending on education, total (% of government expenditure)	WB-WDI	SE.XPD.TOTL.GB.ZS
External resources for health (% of total expenditure on health)	WB-WDI	SH.XPD.EXTR.ZS
Net bilateral aid flows from DAC donors, Total (current US\$) ^g	WB-WDI	DC.DAC.TOTL.CD
Net official development aid (ODA) received (% of Gross National Income) ^{c,h}	WB-WDI	DT.ODA.ODAT.GN.ZS
Net ODA received per capita (current US\$) ^{c,h}	WB-WDI	DT.ODA.ODAT.PC.ZS
Net ODA received (constant 2010 US\$)^{c,h}	WB-WDI	DT.ODA.OATL.CD
Safety and Rule of Law (Ibrahim index with 4 indicators) ⁱ	WB-ADI	MO.INDEX.SRLW.XQ
Other (5 indicators listed)		
Road density (km of road per 100 sq. km of land area)	WB-WDI	IS.ROD.DNST.K2

Roads, paved (% of total roads)	WB-WDI	IS.ROD.PAVE.ZS
Access to electricity (% of population)	WB-WDI	EG.ELC.ACCS.ZS
Population coverage of mobile cellular telephony (%)	WB-ADI	IT.MOB.COV.ZS
Internet users (per 100 people)	WB-WDI	IT.NET.USER.P2

^a 10 indicators in bold were not included in the database due to 100% insufficient data (see methods); analysis dataset has 60 indicators.

^b one of the 27 Countdown to 2015 intervention indicators (or closely related)

^c one of the Countdown to 2015 country profile indicators (or closely related)

^d The GINI index is a measure of income inequality, where 0 represents perfect equality and 100 represents perfect inequality

^e The Human Development Index is a composite measure of development incorporating life expectancy at birth, mean years of schooling and expected years of schooling, and gross national income per capita, where 0 represents very low development and 1 represents very high development

^f Includes BCG, measles and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

^g DAC, Development Assistance Committee. DAC members are Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Japan, Republic of Korea, Luxembourg, Netherlands, New Zealand, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom, United States, and European Union Institutions

^h ODA, Official Development Assistance. ODA includes loans and grants made by DAC members, non-DAC countries, and multilateral institutions

ⁱ Ibrahim index is a composite measure of a countries provision of political, social and economic public goods and services. It is comprised of four indicators: Personal safety, Rule of law, Accountability and corruption and, National security. The index ranges from 0 to 100, with higher values representing better governance.

^j Data source for the current analysis: WB, World Bank Data Catalogue; WDI, World Development Index; ADI, Africa Development Index; HNPS, Health Nutrition & Population Statistics; Country DHS indicates data were obtained directly from each country's Demographic and Health Survey; Countdown to 2015 indicates data were obtained directly from each country's Country Profile in the Countdown to 2015 reports.

^k Indicator code as documented by the World Bank Data Catalogue; <http://databank.worldbank.org>.

Table S4. Median baseline indicator values and corresponding ARC for the 2000-2011 period.

Indicator	N	Median 2000 value (IQR)	Median 2000-2011 ARC (IQR)
<i>Socio-demographics</i>			
Access to improved sanitation facilities (%)	45	25 (13 - 45)	1.6 (0.4 - 2.7)
Access to improved water source (%)	45	62 (51 - 79)	1.1 (0.2 - 1.8)
Adult female literacy rate (%)	35	54 (37 - 72)	1.2 (0.8 - 2.8)
Female age at first marriage (years)	17	21 (20 - 22)	0.2 (-0.1 - 0.6)
Female labor participation rate (%)	45	46 (43 - 49)	0.1 (-0.1 - 0.3)
Fertility rate (births per woman)	46	6 (5 - 6)	-1.4 (-2.0 - -0.9)
GDP per capita (PPP int'l \$)	45	990 (680 - 1,900)	4.3 (2.8 - 6.1)
Gini index (0 to 100)	22	43 (41 - 50)	0.0 (-1.0 - 0.6)
Human development index (0 to 1)	39	0.38 (0.31 - 0.48)	1.1 (0.7 - 1.9)
Labor force participation rate among women (%)	45	66 (50 - 76)	0.3 (-0.1 - 0.6)
Median age at pregnancy (or first birth) (years)	18	19 (19 - 20)	0.2 (0.1 - 0.2)
Median number of living children per household (N)	18	2 (2 - 3)	0.1 (-0.6 - 0.4)
Median years of maternal education (years)	6	4 (3 - 6)	2.4 (0.9 - 4.2)
Poverty headcount ratio at national poverty line (%)	15	46 (39 - 55)	-1.2 (-3.0 - -0.5)
Teenage mothers (%)	15	22 (19 - 32)	-1.1 (-2.5 - -0.6)
Urban population prevalence (%)	46	35 (22 - 44)	1.3 (0.7 - 1.7)
Women with at least some secondary education (%)	23	16 (10 - 30)	3.2 (2.5 - 5.3)
<i>Maternal health</i>			
Adult female mortality rate (rate per 1,000 persons)	46	360 (280 - 420)	-1.2 (-1.8 - -0.6)
Antenatal care during first 5 months of pregnancy (%)	19	57 (45 - 67)	1.4 (0.5 - 3.4)
Maternal mortality ratio (rate per 100,000 births)	45	550 (400 - 840)	-3.6 (-4.5 - -2.0)
Pregnant women receiving ≥4 prenatal care visits (%)	21	46 (23 - 62)	1.7 (-0.3 - 4.0)
Pregnant women receiving prenatal care (%)	38	77 (64 - 88)	1.3 (0.6 - 2.7)
Unmet need for contraception (%)	18	27 (22 - 32)	0.6 (-1.2 - 1.9)
<i>Access to health care</i>			
Births attended by skilled health staff (%)	37	47 (39 - 60)	2.0 (0.3 - 3.4)
Births delivered at a health facility (%)	19	43 (34 - 55)	1.8 (0.1 - 3.7)
Births delivered by caesarean section (%)	18	3 (1 - 4)	5.3 (1.6 - 7.6)
Births with a postnatal visit ≤2 days after delivery (%)	10	8 (5 - 12)	16.0 (8.0 - 21.0)
Children with fever receiving antimalarial drugs (%)	30	53 (32 - 61)	-4.8 (-7.5 - -0.2)
Community health workers (rate per 1,000 persons)	2	0.2 (0.2 - 0.3)	15.0 (13.0 - 17.0)
Diarrhea treatment (%)	25	39 (28 - 45)	2.3 (-1.3 - 5.7)
Hospital beds (rate per 1,000 persons)	8	1.2 (0.4 - 2.0)	-2.3 (-4.6 - -1.0)
Nurses and midwives (rate per 1,000 persons)	6	1.2 (0.6 - 1.6)	-3.0 (-5.0 - -1.0)

Physicians (rate per 100,000 persons)	23	5.7 (2.9 – 12.0)	1.2 (-0.6 - 5.2)
Pregnant women using malaria preventive treatment (%)	3	36 (29 - 37)	10.0 (-1.8 - 15.0)
Seeking ARI treatment (%)	33	40 (27 - 55)	1.7 (-0.3 - 5.0)
<i>Child survival interventions</i>			
Children 1 year old receiving all basic vaccinations (%)	26	41 (29 - 65)	2.8 (1.1 - 5.3)
Exclusive breastfeeding (%)	36	25 (14 - 37)	4.1 (-0.2 - 8.6)
Measles immunization (%)	46	69 (49 - 76)	1.3 (0.3 - 2.5)
Use of insecticide-treated bed nets (%)	30	2 (1 - 4)	34.0 (28.0 - 42.0)
Vitamin A supplementation coverage (%)	37	91 (78 - 96)	-0.3 (-2.0 - 0.7)
<i>Clinical and health conditions</i>			
Adult prevalence of HIV (%)	43	4 (1 - 8)	-0.8 (-3.5 - 1.7)
Low-birthweight babies (%)	30	13 (11 - 17)	-0.3 (-3.6 - 1.0)
Malnutrition prevalence (height for age) (%)	34	40 (34 - 48)	-1.1 (-2.3 - 0.2)
Malnutrition prevalence (weight for age) (%)	34	22 (18 - 27)	-1.7 (-3.2 - -0.9)
Prevalence of wasting (%)	34	9 (7 - 12)	-1.9 (-5.2 - 0.5)
<i>Governance and financing</i>			
External resources for health (%)	45	13 (4.6 - 20)	6.4 (-1.0 - 10.0)
Health expenditure (relative to GDP) (%)	45	5 (4 - 6)	1.4 (0.5 - 3.3)
Health expenditure (relative to government) (%)	45	9 (7 - 10)	2.5 (-0.4 - 3.7)
Net bilateral aid flows from DAC donors (US \$ in millions)	46	120 (40 - 260)	9.1 (5.0 - 16.0)
Net ODA received (% of GNI) (%)	46	9 (4 - 13)	-0.9 (-5.4 - 5.4)
Net ODA received per capita (US\$)	46	23 (15 - 41)	7.4 (2.4 - 13.0)
Out-of-pocket health expenditure (%)	45	43 (26 - 58)	-1.7 (-3.9 - -0.6)
Public spending on education (% of GDP) (%)	34	4 (3 - 5)	2.1 (0.2 - 3.9)
Public spending on education (% of gov. expenditure) (%)	25	16 (14 - 19)	1.6 (-0.2 - 3.0)
Safety and Rule of Law (0 to 100)	46	54 (43 - 66)	-0.0 (-0.9 - 0.4)
<i>Other</i>			
Households with television (%)	24	18 (6 - 29)	6.7 (4.1 – 10.0)
Internet users (rate per 100 persons)	46	0.2 (0.1 - 0.5)	34.0 (25.0 - 46.0)
Paved roads (%)	19	12 (10 - 29)	1.2 (-1.4 - 3.1)
Population coverage of cell phones (%)	26	37 (17 - 82)	8.5 (1.3 - 16.0)
Road density (km of road per 100 sq km)	8	8 (4 - 20)	1.4 (0.3 - 6.2)

Table S5. Crude and adjusted associations between indicators and under-five mortality ARR, 2000-2013.

Indicator ^a	Crude				Adjusted ^{b,c}		
	N	β	90% CI	p-value	β	90% CI	p-value
<i>Socio-demographics</i>							
Access to improved sanitation facilities (%) ^e	42	0.08	(-0.20, 0.37)	0.350	0.04	(-0.23, 0.31)	0.385
Access to improved water source (%)	42	0.40	(-0.05, 0.85)	0.139	0.36	(-0.07, 0.78)	0.155
Adult female literacy rate (%)	33	0.02	(-0.26, 0.31)	0.392	0.11	(-0.19, 0.41)	0.324
Female age at first marriage (years)	16	0.07	(-0.83, 0.98)	0.388	-0.16	(-0.66, 0.34)	0.334
Female labor participation rate (%)	42	-1.07	(-2.42, 0.29)	0.171	-0.32	(-1.56, 0.92)	0.362
Fertility rate (births per woman)	42	0.78	(0.26, 1.31)	0.023	0.54	(0.07, 1.02)	0.069
GDP per capita (PPP int'l \$) ^f	42	0.00	(-0.21, 0.21)	0.396	0.06	(-0.14, 0.26)	0.352
GINI index (0 to 100)	21	-0.04	(-0.46, 0.39)	0.389	0.22	(-0.16, 0.60)	0.248
Human development index (0 to 1) ^g	35	0.12	(0.04, 0.20)	0.023	0.09	(0.01, 0.17)	0.066
Labor force participation rate among women (%)	42	-0.33	(-1.19, 0.53)	0.325	0.19	(-0.57, 0.96)	0.363
Median age at pregnancy (or first birth) (years)	17	-0.43	(-5.16, 4.30)	0.388	-0.45	(-3.71, 2.82)	0.379
Median number of living children per household (N)	17	0.78	(-0.06, 1.62)	0.123	-0.21	(-0.84, 0.42)	0.334
Median years of maternal education (years)	6	0.01	(-0.23, 0.26)	0.373	-0.08	(NA, NA)	
Poverty headcount ratio at national poverty line (%)	15	-0.08	(-0.46, 0.30)	0.367	-0.25	(-0.63, 0.13)	0.215
Teenage mothers (%)	14	-0.41	(-1.09, 0.26)	0.230	-0.44	(-0.72, -0.17)	0.022
Urban population prevalence (%)	42	0.53	(-0.03, 1.09)	0.117	0.39	(-0.15, 0.93)	0.196
Women with at least some secondary education (%)	21	-0.11	(-0.26, 0.04)	0.187	-0.05	(-0.17, 0.07)	0.303
<i>Maternal health</i>							
Adult female mortality rate (rate per 1,000 persons)	42	-0.49	(-0.83, -0.16)	0.025	-0.25	(-0.61, 0.10)	0.199
Antenatal care during first 5 months of pregnancy (%)	18	0.35	(0.12, 0.58)	0.022	0.29	(0.10, 0.47)	0.024
Maternal mortality ratio (rate per 100,000 births) ^d	35	-0.57	(-0.80, -0.35)	<0.001	-0.47	(-0.69, -0.24)	0.003
Pregnant women receiving ≥ 4 prenatal care visits (%)	20	0.22	(0.06, 0.37)	0.034	0.07	(-0.01, 0.16)	0.134
Pregnant women receiving prenatal care (%)	35	0.09	(-0.30, 0.48)	0.369	0.09	(-0.24, 0.41)	0.359
Unmet need for contraception (%)	18	0.02	(-0.10, 0.14)	0.373	0.03	(-0.06, 0.11)	0.331
<i>Access to health care</i>							
Seeking ARI treatment (%) ^d	31	0.26	(0.10, 0.42)	0.016	0.22	(0.09, 0.35)	0.012
Births attended by skilled health staff (%) ^d	34	0.21	(-0.03, 0.45)	0.142	0.18	(-0.07, 0.43)	0.192
Births delivered at a health facility (%) ^d	18	0.17	(-0.01, 0.36)	0.112	0.05	(-0.10, 0.20)	0.335
Births delivered by caesarean section (%) ^d	17	0.18	(0.04, 0.32)	0.052	0.13	(0.05, 0.20)	0.017
Births with a postnatal visit ≤ 2 days after delivery (%) ^d	9	-0.10	(-0.22, 0.01)	0.134	-0.14	(-0.14, -0.14)	<0.001
Children with fever receiving antimalarial drugs (%) ^d	28	0.05	(-0.02, 0.12)	0.176	0.06	(-0.01, 0.12)	0.143

Community health workers (rate per 1,000 persons)	2						
Diarrhea treatment (%) ^d	24	-0.10	(-0.18, -0.01)	0.072	-0.09	(-0.18, 0.00)	0.111
Hospital beds (rate per 1,000 persons)	8	-0.09	(-0.29, 0.11)	0.283	-0.32	(-0.45, -0.19)	0.013
Nurses and midwives (rate per 1,000 persons)	5						
Physicians (rate per 100,000 persons)	21	0.03	(-0.09, 0.15)	0.360	0.06	(0.00, 0.13)	0.089
Pregnant women using malaria preventive treatment (%)	3						
<i>Child survival interventions</i>							
Children 1 year old receiving all basic vaccinations (%) ^d	23	0.09	(-0.09, 0.28)	0.278	-0.14	(-0.39, 0.10)	0.236
Exclusive breastfeeding (%) ^d	33	-0.01	(-0.08, 0.06)	0.388	-0.03	(-0.09, 0.04)	0.308
Measles immunization (%) ^d	35	0.03	(-0.19, 0.26)	0.382	0.11	(-0.08, 0.30)	0.254
Use of insecticide-treated bed nets (%) ^d	28	0.02	(-0.02, 0.07)	0.277	0.01	(-0.04, 0.05)	0.381
Vitamin A supplementation coverage (%) ^d	31	0.11	(0.01, 0.21)	0.085	0.06	(-0.04, 0.16)	0.241
<i>Clinical and health conditions</i>							
Adult prevalence of HIV (%)	42	-0.13	(-0.25, -0.01)	0.073	-0.09	(-0.20, 0.02)	0.171
Low-birthweight babies (%) ^d	27	0.04	(-0.10, 0.17)	0.359	0.05	(-0.07, 0.17)	0.303
Malnutrition prevalence (height for age) (%) ^d	32	0.11	(-0.15, 0.38)	0.306	0.15	(-0.10, 0.40)	0.244
Malnutrition prevalence (weight for age) (%) ^d	32	-0.07	(-0.28, 0.15)	0.344	-0.02	(-0.23, 0.19)	0.390
Prevalence of wasting (%) ^d	32	0.01	(-0.12, 0.14)	0.390	0.03	(-0.09, 0.15)	0.357
<i>Governance and financing</i>							
External resources for health (%)	42	0.01	(-0.05, 0.07)	0.373	-0.01	(-0.06, 0.05)	0.390
Health expenditure (% of GDP) (%)	42	0.24	(0.07, 0.40)	0.028	0.26	(0.11, 0.41)	0.010
Health expenditure (% of government expenditure) (%) ^f	42	0.11	(-0.01, 0.23)	0.120	0.08	(-0.04, 0.19)	0.217
Net bilateral aid flows from DAC donors (US\$ in millions)	42	0.01	(-0.06, 0.07)	0.394	-0.05	(-0.12, 0.03)	0.224
Net ODA received (relative to GNI) (%)	42	-0.01	(-0.09, 0.07)	0.392	-0.08	(-0.15, -0.01)	0.074
Net ODA received per capita (US\$)	42	0.02	(-0.03, 0.08)	0.301	-0.01	(-0.06, 0.05)	0.388
Out-of-pocket health expenditure (%) ^f	42	-0.19	(-0.37, -0.02)	0.079	-0.19	(-0.34, -0.03)	0.061
Public spending on education (% of GDP) (%)	33	0.03	(-0.13, 0.19)	0.379	-0.06	(-0.21, 0.10)	0.329
Public spending on education (% of gov. expenditure) (%)	24	0.19	(0.00, 0.38)	0.106	0.06	(-0.14, 0.27)	0.345
Safety and Rule of Law (0 to 100)	42	0.10	(-0.14, 0.34)	0.316	-0.07	(-0.30, 0.15)	0.342
<i>Other</i>							
Households with television (%)	23	0.29	(0.13, 0.46)	0.009	0.19	(0.05, 0.33)	0.044
Internet users (rate per 100 persons)	42	0.05	(0.01, 0.09)	0.048	0.03	(-0.02, 0.07)	0.221
Paved roads (%)	17	0.07	(-0.10, 0.24)	0.309	0.06	(-0.07, 0.19)	0.294
Population coverage of cell phones (%)	24	0.01	(-0.07, 0.09)	0.387	0.01	(-0.05, 0.07)	0.383
Road density (km of road per 100 sq. km)	7	-0.05	(-0.37, 0.26)	0.364	-0.12	(-0.46, 0.22)	0.240

^a 10 of 70 indicators not shown in table because sufficient data were missing for all 46 countries

^b N>2 required to run crude robust linear regression models; N>6 required to run adjusted robust linear regression models which include a minimum of four covariates.

^c All results adjusted for improved water source, health expenditure (relative to GDP), adult HIV prevalence, and urban population prevalence unless otherwise noted below:

^d further adjusted for receipt of prenatal care

^e not adjusted for improved water source

^f not adjusted for health expenditure (relative to GDP)

^g Because the range of the HDI is only from 0 to 1, the β represents the change in ARR for each 0.10 increase in HDI ARC.